

CLASS 11th – BIOLOGY

BANK OF BIOLOGY

NCERT BASED SHORT NOTES

1. THE LIVING WORLD

Life is a unique, complex organization of molecules, expressing through chemical reactions which lead to growth, development, responsiveness, adaptation & reproduction.

A **living organism** is self-replicating, evolving and self-regulating interactive system capable of responding to external stimuli.

PROPERTIES OF LIVING ORGANISMS

1. Growth

- It is the increase in number & mass of cells by cell division.
- In plants, growth continues throughout their lifespan.
- In animals, growth is only up to a certain age. However, cell division occurs to replace lost cells.
- Basically, growth is the increase in mass & size. Thus non-living objects also grow (surface accumulation of material). So growth is not a defining property of living organisms.
- In living organisms, growth is from inside.

2. Reproduction

- It is the production of progeny having features similar to those of parents.
- Organisms reproduce asexually and sexually.
- In unicellular organisms, growth & reproduction are same because they reproduce by cell division.
- Many organisms do not reproduce (e.g. mules, worker bees, infertile human couples, etc). Hence, reproduction is not a perfect defining property of living organisms.

3. Metabolism

- It is the sum total of all biochemical reactions taking place inside a living system.
- It is the defining feature of living organisms.
- Metabolic reactions can be demonstrated outside the body in cell-free systems. Isolated metabolic reactions *in vitro* are not living things but are living reactions.

4. Cellular organization

- Organisms are made up of one or more cells.
- It is the defining feature of living organisms.

5. Consciousness

- It is the ability of organisms to sense their environment and respond to environmental stimuli (like light, water, temperature, other organisms, chemicals, pollutants, etc).
- All organisms are 'aware' of their surroundings. So, it is the defining property of living organisms.
- Human is the only organism having **self-consciousness**.

DIVERSITY IN THE LIVING WORLD

- The number and types of organisms present on earth refer to **biodiversity**.
 - Number of species described is **1.7-1.8 million**.
 - **Taxonomy** is the study of **identification, classification & nomenclature** of organisms.
- Systematics** (Latin 'systema' = systematic arrangement) deals with evolutionary relationships among organisms.
- *Systema Naturae* is the book written by **Linnaeus**.

Basic processes of taxonomy

- **Characterization:** It is the understanding of characters of organisms such as external and internal structure, structure of cell, development process, ecological information etc.
- **Identification:** It is the correct description of the organism so that the naming is possible.
- **Classification:** It is the grouping of organisms into convenient categories (**taxa**) based on characters.
- **Nomenclature (naming):** It is the standardization of names of the organisms such that an organism is known by the same name all over the world.

The system of naming with two components is called **Binomial nomenclature**. It is proposed by **Linnaeus**.

Botanical names are based on the rules in **International Code for Botanical Nomenclature (ICBN)**.

Zoological names are based on **International Code for Zoological Nomenclature (ICZN)**.


Universal rules of Binomial nomenclature

- Scientific names are in **Latin** or Latinised and written in **italics**. When handwritten, they are underlined separately.

- The first word is genus name (**Generic name**) and second word is the species name (**specific epithet**).
E.g. *Homo sapiens*- *Homo* represents the genus name and *sapiens* represents the species name.
- The Genus name starts with capital letter and the species name starts with small letter.
- Name of the author (in abbreviated form) appears at the end of the biological name.
E.g., *Mangifera indica* Linn. It indicates that this species was first described by Linnaeus.

TAXONOMIC CATEGORIES

- Classification involves hierarchy of steps in which each step represents a **taxonomic category (rank)**.
- All categories together constitute a **taxonomic hierarchy**.
- A group of organisms occupying a particular category is called a **taxon (pl. taxa)**. E.g. Class Mammalia.
- Each **category** or **taxon** represents a unit of classification.



Species: It is a group of closely related organisms capable of interbreeding to produce fertile offspring.

It is the lowest category. E.g.

Common name	Generic name	Specific epithet
Mango	<i>Mangifera</i>	<i>indica</i>
Potato	<i>Solanum</i>	<i>tuberosum</i>
Nightshade	<i>Solanum</i>	<i>nigrum</i>
Tomato	<i>Solanum</i>	<i>lycopersicum</i>
Brinjal	<i>Solanum</i>	<i>melongena</i>
Lion	<i>Panthera</i>	<i>leo</i>
Tiger	<i>Panthera</i>	<i>tigris</i>
Leopard	<i>Panthera</i>	<i>pardus</i>
Modern man	<i>Homo</i>	<i>sapiens</i>

Genus: It is the aggregates of closely related species.

E.g. Potato, tomato & brinjal are species of genus *Solanum*.

Lion, leopard & tiger are species of genus *Panthera*. This genus differs from genus *Felis* (genus of cats).

Family: It is a group of closely related genera. E.g.

Family Solanaceae includes Genus *Solanum*, Genus *Petunia* and Genus *Datura*.

Family Felidae includes Genus *Panthera* and Genus *Felis*.

Order: It is the assemblage of related families. E.g.

Order *Polymoniales* includes Family *Convolvulaceae* and Family *Solanaceae*.

Order *Carnivora* includes Family *Felidae* & Family *Canidae*.

Class: It is the assemblage of related orders. E.g.

Order *Primata*, *Carnivora* etc. is placed in class *Mammalia*.

Phylum (Division in case of plants): It is the assemblage of related classes.

E.g. Classes *Amphibia*, *Reptilia*, *Aves*, *Mammalia* etc. come under phylum Chordata.

Kingdom: The assemblage of related phyla. It is the highest category. E.g. Kingdom *Plantae*, Kingdom *Animalia* etc.

Organisms with their taxonomic categories

Common name	Man	Housefly	Mango	Wheat
Biological name	<i>Homo sapiens</i>	<i>Musca domestica</i>	<i>Mangifera indica</i>	<i>Triticum aestivum</i>
Species	<i>sapiens</i>	<i>domestica</i>	<i>indica</i>	<i>aestivum</i>
Genus	Homo	Musca	Mangifera	Triticum
Family	Hominidae	Muscidae	Anacardiaceae	Poaceae
Order	Primata	Diptera	Sapindales	Poales
Class	Mammalia	Insecta	Dicotyledonae	Monocotyledonae
Phylum/Division	Chordata	Arthropoda	Angiospermae	Angiospermae
Kingdom	Animalia	Animalia	Plantae	Plantae

TAXONOMICAL AIDS

a. Herbarium

- It is a store house (repository) of plant specimens that are dried, pressed and preserved on sheets and are arranged according to universally accepted classification.
- Herbarium sheets are labelled with information about date and place of collection, English, local and botanical names, family, collector's name etc.

b. Botanical gardens

- These are specialized gardens having collections of living plants for reference and identification.
- Each plant is labelled with its botanical name and family.
- **Famous botanical gardens:**
 - o Royal Botanical Garden at Kew (England).
 - o Indian Botanical Garden, Howrah (India).
 - o At National Botanical Research Institute, Lucknow (India).

c. Biological Museum

- It is a collection of *preserved plants and animals* for study and reference.
- A museum contains
 - Specimens preserved in preservative solutions in containers or jars.
 - Preserved dry specimens of plants and animals.
 - Insects preserved in insect boxes after collecting, killing and pinning.

- Stuffed larger animals like birds and mammals.
- Collections of animal skeletons.

d. Zoological Parks (Zoos)

- These are the places where *live wild animals* are kept in protected environments under human care.
- It helps to learn about their food habits and behaviour.

e. Key

- It is an analytical method of identification of organisms based on similarities and dissimilarities.
- It is based on the contrasting characters generally in a pair called **couplet**.
- Each couplet has two opposite options. Of these, only relevant option is accepted and other is rejected.
- Each statement in the key is called a **lead**.

Flora, manuals, monographs & catalogues

- **Flora:** Actual account of habitat and distribution of plant species of a given area.
- **Manuals:** The record that contains information for identification of names of species found in an area.
- **Monographs:** The records that contain information on any one taxon.
- **Catalogue:** Alphabetical list of species.

MODEL QUESTIONS

1. "Consciousness is a defining property of living organisms". Comment.
2. Note the relationship between first two words and fill up the fourth place
 - a. Monera: Kingdom Primata:
 - b. Botanical nomenclature: ICBN Zoological nomenclature:
 - c. Highest taxonomic rank: Kingdom Lowest taxonomic rank:
3. Re-arrange the following in the descending order:
Genus – order – kingdom – family – phylum – species – class
4. Analyze the following scientific names
 - i. panthera Tigris ii. Mangifera Indica
 - a. If there any mistake, correct them.
 - b. Mention their common name.
5. Fill up the table

Common name	Housefly	Mango	Wheat
Family	A	Anacardiaceae	B
Order	Diptera	C	Poales
Class	D	Dicotyledonae	E
Phylum/Division	Arthropoda	F	Angiospermae

6. Compare museum and zoological park.
7. Complete the table

Taxonomical aid	Plant/Animal	Live/Dead
Herbarium		
Botanical garden		
Zoological park		
Biological museum		

8. Match the following

A	B
a. Herbarium	i. Contrast pair or couplet
b. Key	ii. Information on any one taxon
c. Monographs	iii. Habitat and distribution of plant species
d. Flora	iv. Storage of collected dried plant specimens

2. BIOLOGICAL CLASSIFICATION

Aristotle's classification

- Aristotle was the earliest to attempt a more scientific basis for classification of organisms.
- He classified plants to **trees, shrubs & herbs** and animals into 2 groups- those **with red blood & without red blood**.

Linnaeus's Two-kingdom classification

- **Linnaeus** (1758) classified organisms into **Two Kingdoms**- Kingdom **Plantae** & Kingdom **Animalia**.

Drawbacks of 2-kingdom classification:

- **Prokaryotes** (Bacteria, cyanobacteria) and **eukaryotes** (fungi, mosses, ferns, gymnosperms & angiosperms) were included under 'Plants' based on the presence of cell wall. But they are widely differed in other characteristics.

- It included the unicellular and the multicellular organisms in same group. E.g. *Chlamydomonas* and *Spirogyra* were placed under algae.
- It did not differentiate between the heterotrophic fungi and the autotrophic green plants. Fungi have chitinous cell wall while the green plants have cellulosic cell wall.

Five Kingdom Classification

- It is proposed by **R.H. Whittaker (1969)**.
- It includes Monera, Protista, Fungi, Plantae & Animalia.
- This is based on cell structure, thallus organization, mode of nutrition, reproduction and phylogenetic relationships.


Three-domain system: It divides Kingdom Monera into two domains. Eukaryotic kingdoms are included in third domain. Thus it is **six-kingdom classification**.

Characteristics of the five kingdoms

Characters	Monera	Protista	Fungi	Plantae	Animalia
Cell type	Prokaryotic	Eukaryotic	Eukaryotic	Eukaryotic	Eukaryotic
Cell wall	Non-cellulosic (poly-saccharide + amino acid)	Present in some	Present (Chitin & polysaccharides)	Present (Cellulose)	Absent
Nuclear membrane	Absent	Present	Present	Present	Present
Body organisation	Cellular	Cellular	Multicellular, loose tissue	Tissue/organ	Tissue/organ/organ system
Mode of nutrition	Autotrophic (photosynthetic & chemosynthetic) and heterotrophic (saprophyte/parasite)	Autotrophic (photosynthetic) and heterotrophic	Heterotrophic (saprophytic or parasitic)	Autotrophic (photosynthetic)	Heterotrophic (holozoic, saprophytic etc.)

1. KINGDOM MONERA (BACTERIA)

- Bacteria are the most abundant microorganisms.
- Hundreds of bacteria are present in a handful of soil.
- They also live in extreme habitats such as hot springs, deserts, snow & deep oceans. Many are parasites.
- Based on shape, bacteria are 4 types: **Coccus** (Spherical), **Bacillus** (Rod-shaped), **Vibrium** (Comma-shaped) & **Spirillum** (Spiral).



- Some bacteria are **autotrophic** (synthesize food from inorganic substrates). Majority are **heterotrophs** (they do not synthesize food but depend on other organisms or on dead organic matter for food).

I. Archaeobacteria


- They live in harshest habitats such as extreme salty areas (**halophiles**), hot springs (**thermoacidophiles**) and marshy areas (**methanogens**).
- Archaeobacteria have a different cell wall structure for their survival in extreme conditions.
- **Methanogens** are present in the guts of ruminant animals (cows, buffaloes etc). They produce **methane (biogas)** from the dung of these animals.

II. Eubacteria ('true bacteria')

- They have a **rigid cell wall** and a **flagellum** (if motile).
- They include **Autotrophs** (photosynthetic and chemosynthetic) and **Heterotrophs**.

a. Photosynthetic autotrophs (E.g. Cyanobacteria):

- They have **chlorophyll a** similar to green plants.
- Cyanobacteria (blue-green algae) are unicellular, colonial or filamentous, marine or terrestrial algae.
- The colonies are generally surrounded by **gelatinous sheath**.
- They often form blooms in polluted water bodies.
- Some of them fix atmospheric nitrogen in specialized cells (**heterocysts**). E.g., *Nostoc* & *Anabaena*.



b. Chemosynthetic autotrophs:

- They oxidize inorganic substances such as nitrates, nitrites & ammonia and use the released energy for ATP production.
- They help in recycling nutrients like nitrogen, phosphorous, iron and sulphur.

c. Heterotrophic bacteria:

- They are the **most abundant** in nature.

- The majority are important **decomposers**.


Impacts of Heterotrophic bacteria on human affairs:

- They are used to make curd from milk.
- Production of antibiotics.
- Fixing nitrogen in legume roots etc.
- Some are pathogens causing diseases. E.g. Cholera, typhoid, tetanus and citrus canker.

Reproduction in Bacteria:

- Bacteria reproduce mainly by **fission**.
- Under unfavourable conditions, they produce **spores**.

- They also reproduce by a sort of sexual reproduction (DNA transfer from one bacterium to other).



A dividing bacterium

Mycoplasmas are organisms without a cell wall. They are the **smallest living cells**. They can survive without oxygen. Many are pathogenic in animals and plants.

2. KINGDOM PROTISTA

- It includes **single-celled eukaryotes**.
- The cell contains a well-defined nucleus and other membrane-bound organelles. Some have flagella or cilia.
- Protists are primarily aquatic.
- It is a link with plants, animals and fungi.
- They reproduce asexually and sexually (cell fusion and zygote formation).
- Protista includes **Chrysophytes, Dinoflagellates, Euglenoids, Slime moulds** and **Protozoans**.

I. Chrysophytes

- Found in fresh water and marine environments.
- Microscopic and float passively in water currents (plankton).
- Most of them are photosynthetic.
- It includes **diatoms & golden algae (desmids)**.
- **Diatoms:** They have siliceous cell walls forming two thin overlapping shells, which fit together as in a soap box. The cell wall deposit of diatoms over billions of years in their habitat is known as '**diatomaceous earth**'. This is used in polishing, filtration of oils and syrups.
- Diatoms are the chief 'producers' in the oceans.

II. Dinoflagellates

- Mostly marine and photosynthetic.
- They appear yellow, green, brown, blue or red based on the main pigments present in their cells.
- The cell wall has stiff cellulose plates on the outer surface.
- Most of them have **2 flagella**; one lies longitudinally and the other transversely in a furrow between the wall plates.
- **Red dinoflagellates** (E.g. *Gonyaulax*) undergo rapid multiplication so that the sea appears red (**red tides**). They release toxins that kill marine animals like fishes.

III. Euglenoids

- Mainly fresh water organisms found in stagnant water.
- Instead of a cell wall, they have a protein rich layer called **pellicle**. It makes their body flexible.

- They have **two flagella**, a short and a long one.
- They are **photosynthetic** in the presence of sunlight. In the absence of sunlight, they behave like **heterotrophs** by preying on smaller organisms.
- The pigments are identical to those in higher plants.
- E.g. *Euglena*.

IV. Slime Moulds

- They are saprophytic protists.
- The body moves along decaying twigs and leaves engulfing organic material.
- Under suitable conditions, they form an aggregation called **plasmodium**. It may spread over several feet.
- Under unfavourable conditions, plasmodium differentiates and forms fruiting bodies bearing **spores** at their tips. Spores have true walls. They are highly resistant and survive for many years. Spores are dispersed by air.

V. Protozoans

They are heterotrophs (predators or parasites).

They are the primitive relatives of animals.

There are 4 major groups of protozoans:

- **Amoeboid protozoans:** They live in fresh water, sea water or moist soil. They move and capture prey by putting out **pseudopodia** (false feet). E.g. *Amoeba*. Marine forms have silica shells on their surface. Some of them are parasites. E.g. *Entamoeba*.
- **Flagellated protozoans:** They are free-living or parasitic. They have **flagella**. The parasitic forms cause diseases like **sleeping sickness**. E.g. *Trypanosoma*.
- **Ciliated protozoans:** They are aquatic, actively moving organisms using thousands of **cilia**. They have a cavity (**gullet**) that opens to outside. By the movement of cilia, the water with food enters gullet. E.g. *Paramecium*.
- **Sporozoans:** They have an infectious spore-like stage in their life cycle. E.g. *Plasmodium* (malarial parasite).

3. KINGDOM FUNGI

- It is a unique kingdom of heterotrophic organisms.
- Fungi are cosmopolitan.
- They grow in warm and humid places.
- E.g. mould on bread & rotten fruits, mushroom, toadstools.
- White spots on mustard leaves are due to a parasitic fungus.
- Some fungi are the source of antibiotics, e.g., *Penicillium*.

- Some unicellular fungi (e.g. yeast) are used to make bread and beer.
- Other fungi cause diseases in plants and animals. E.g. wheat rust-causing *Puccinia*.
- Except yeasts, fungi are filamentous. Their bodies consist of thread-like structures called **hyphae**.

- The network of hyphae is known as **mycelium**.
- Hyphae are 2 types:
 - o **Coenocytic hyphae**: They are continuous tubes filled with multinucleated cytoplasm.
 - o **Septate hyphae**: They have septae or cross walls.
- Fungal cell wall is made of **chitin & polysaccharides**.
- Most fungi are **saprophytes** (absorb soluble organic matter from dead substrates). Some are **parasites**.
- Some live as **symbionts**. E.g. **Lichens** (fungi+ algae), **mycorrhiza** (fungi + roots of higher plants).

Reproduction:

- **Vegetative propagation**: By fragmentation, fission & budding.
- **Asexual reproduction**: By spores such as **conidia**, **sporangiospores** and **zoospores**.
- **Sexual reproduction**: By **oospores**, **ascospores** and **basidiospores**. They are produced in distinct structures called **fruiting bodies**.
- The sexual cycle involves 3 steps:
 - a. Plasmogamy**: Fusion of protoplasm between two motile or non-motile gametes.
 - b. Karyogamy**: Fusion of two nuclei.
 - c. Meiosis** in zygote to give haploid spores.
- When a fungus reproduces sexually, **two haploid hyphae** of compatible mating types come together and fuse.
- In some fungi, the fusion of two haploid cells immediately results in **diploid cells (2n)**.
- In ascomycetes and basidiomycetes, a **dikaryotic stage** or **dikaryophase** ($n + n$ i.e. two nuclei per cell) occurs. Such a condition is called a **dikaryon**. Later, parental nuclei fuse and the cells become diploid.
- The fungi form fruiting bodies in which reduction division occurs, leading to formation of haploid spores.

Based on morphology of mycelium, mode of spore formation & fruiting bodies, Fungi are classified into different classes:

1. **Phycomycetes**
2. **Ascomycetes**
3. **Basidiomycetes**
4. **Deuteromycetes**

I. Phycomycetes (Lower Fungi)

- They occur in aquatic habitats and on decaying wood in moist and damp places or as obligate parasites on plants.
- The mycelium is **aseptate** and **coenocytic**.
- **Asexual reproduction**: By motile **zoospores** or by non-motile **aplanospores**. These are produced in sporangium.
- **Sexual reproduction**: **Zygosporangia** are formed by fusion of two gametes. These gametes are **isogamous** (similar in morphology) or **anisogamous or oogamous** (dissimilar).

- E.g. *Mucor*, *Rhizopus* (bread mould) and *Albugo* (parasitic fungi on mustard).

II. Ascomycetes (sac-fungi)

- They are unicellular (e.g., yeast, *Sacharomyces*) or multicellular (e.g., *Penicillium*).
- Mycelium is branched and septate.
- They are saprophytic, decomposers, parasitic or coprophilous (growing on dung).
- **Asexual reproduction**: By **conidia** produced exogenously on the special mycelium called **conidiophores**. Conidia germinate to produce mycelium.
- **Sexual reproduction**: By **ascospores** produced endogenously in sac like asci (sing. ascus). The asci are arranged to form fruiting bodies called **ascocarps**.
- E.g. *Aspergillus*, *Claviceps* and *Neurospora*.
- *Neurospora* is used in biochemical and genetic work.
- Morels & truffles are edible.

III. Basidiomycetes

- Includes **mushrooms, bracket fungi or puffballs**.
- They grow in soil, on logs and tree stumps and in living plant bodies as parasites (e.g. rusts and smuts).
- The mycelium is branched and septate.
- The asexual spores are generally not found, but **vegetative reproduction** by fragmentation is common.
- The sex organs are absent, but **plasmogamy** occurs by fusion of two vegetative or somatic cells of different strains or genotypes. The resultant structure is dikaryotic which gives rise to **basidium**. Karyogamy and meiosis take place in basidium producing four **basidiospores** exogenously. Basidia are arranged in fruiting bodies (**basidiocarps**).
- E.g. *Agaricus* (mushroom), *Ustilago* (smut) and *Puccinia* (rust fungus).

IV. Deuteromycetes

- Commonly known as **imperfect fungi** because only their asexual or vegetative phases are known.
- When perfect (sexual) stages were discovered, they were often moved to ascomycetes or basidiomycetes.
- It is also possible that **asexual and vegetative stage** have been given one name placing under **deuteromycetes** and the **sexual stage** another name placing under **another class**. When the linkages were established, the fungi were correctly identified and moved out of deuteromycetes.
- They reproduce only by asexual spores (**conidia**).
- The mycelium is septate and branched.
- Some are saprophytes or parasites. Majority are decomposers of litter and help in mineral cycling.
- E.g. *Alternaria*, *Colletotrichum* and *Trichoderma*.

4. KINGDOM PLANTAE (PLANT KINGDOM)

- Plants are **eukaryotic chlorophyll**-containing organisms with **cellulosic cell wall**.
- Some are partial heterotrophs (e.g. insectivorous plants like bladderwort & Venus flytrap) or parasites (e.g. *Cuscuta*).
- Plantae includes **algae, bryophytes, pteridophytes, gymnosperms and angiosperms**.


- Life cycle of plants has 2 phases: **Diploid sporophytic & haploid gametophytic**. These phases alternate with each other. This is called **alternation of generation**.
- Among different plant groups, length of the haploid & diploid phases is varied. Also, these phases are free living or dependent on others.

5. KINGDOM ANIMALIA (ANIMAL KINGDOM)

- Animals are **multicellular, heterotrophic, eukaryotic** organisms **without cell wall**.
- They directly or indirectly depend on plants for food.
- They digest their food in an internal cavity and store food reserves as glycogen or fat. Their mode of nutrition is **holozoic** (by ingestion of food).
- They have a definite growth pattern and grow into adults that have a definite shape and size.
- Higher forms show sensory and neuromotor mechanism.
- Most of them are capable of locomotion.
- The sexual reproduction is by copulation of male and female followed by embryological development.

VIRUSES, VIROIDS, PRIONS AND LICHENS

- In the five-kingdom classification, acellular organisms (viruses, viroids & prions) and lichens are not mentioned.
- Viruses are non-cellular and not truly 'living'. So they are not included in five-kingdom classification.
- Viruses have an inert crystalline structure outside the living cell.
- Viruses are **obligate parasites**.
- When they infect a cell, they take over the machinery of the host cell to replicate themselves, killing the host.
- **Louis Pasteur** gave the name **virus** (means venom or poisonous fluid).
- **D.J. Ivanowsky (1892)** discovered virus. He recognized certain microbes that cause mosaic disease of tobacco. They were smaller than bacteria because they passed through bacteria-proof filters.
- **M.W. Beijerinck (1898)** demonstrated that the extract of the infected tobacco plants cause infection in healthy plants and called the fluid as *Contagium vivum fluidum* (infectious living fluid).
- **W.M. Stanley (1935)** showed that viruses could be crystallized and crystals consist largely of proteins.
- A virus is a **nucleoprotein**, i.e., it has a **protein coat (capsid) & genetic material (RNA or DNA)**.
- The genetic material is infectious.
- No virus contains both RNA & DNA.
- Generally, plant viruses have single stranded RNA. Animal viruses have either single or double stranded RNA or double stranded DNA. **Bacteriophages** (viruses that infect bacteria) usually have double stranded DNA.
- The **capsid** made of small subunits (**capsomeres**) protects nucleic acid. Capsomeres are arranged in **helical or polyhedral geometric** forms.



- Viruses cause diseases like **mumps, small pox, herpes, influenza & AIDS**. In plants, the symptoms can be mosaic formation, leaf rolling and curling, yellowing and vein clearing, dwarfing and stunted growth.
- **Viroid**: It is an infectious agent with a free low molecular weight RNA and no protein coat. These are smaller than viruses. It is discovered by **T.O. Diener (1971)**. He found that it caused **potato spindle tuber disease**.
- **Prions**: These are **abnormally folded protein** that cause some infectious neurological diseases. These are similar in size to viruses. They cause **bovine spongiform encephalopathy (BSE) or mad cow disease** in cattle and its analogous variant **Cr-Jacob disease (CJD)** in humans.

LICHENS

- Lichens are symbiotic associations (mutually useful associations) between **algae & fungi**.
- The algal component is called **phycobiont** (autotrophic) and fungal component is **mycobiont** (heterotrophic).
- Algae prepare food for fungi and fungi provide shelter and absorb mineral nutrients and water for its partner.
- Lichens are very good **pollution indicators**. They do not grow in polluted areas.

MODEL QUESTIONS

- Based on the relationship, fill in the blanks.

a) Sac fungi: Ascomycetes	Imperfect fungi:
b) Ciliated Protozoans: Paramecium	Flagellated Protozoans:
c) Spherical shaped bacteria:	Rod shaped bacteria: Bacillus
- Louis Pasteur named virus which means venom or poisonous fluid.

a. Who crystallized virus for the first time?	b. Name one plant disease caused by virus.
---	--
- Arrange the organisms given in brackets under two categories, prokaryotes and Eukaryotes.
(*Rhizopus, Amoeba, Chlamydomonas, Nostoc, Bacteria, Yeast, Paramecium, Mycoplasma, Anabaena*)
- Select the odd man. Justify your answer.

a) Chrysophytes, Dinoflagellates, Mycoplasma, Euglenoids	b) Oscillatoria, Spirulina, Ascomycete, Nostoc
c) Oospores, ascospores, zoospores, basidiospores	
- Differentiate between ascomycetes and basidiomycetes.
- Viruses are not included in five-kingdom classification. Why?

3. PLANT KINGDOM

Systems of Biological classification

1. Artificial classification systems

- Earliest systems of classification.
- They were based on vegetative characters or superficial morphological characters such as habit, colour, number and shape of leaves, etc.
- **Linnaeus's artificial system** of classification was based on the androecium structure.

Drawbacks:

- They separated the closely related species since they were based on a few characteristics.
- Equal weightage to vegetative and sexual characteristics. This is not acceptable since the vegetative characters are more easily affected by environment.

2. Natural classification systems

- These are based on natural affinities among organisms.

- It considers external features and internal features (ultrastructure, anatomy, embryology & phytochemistry).
- E.g. Classification for flowering plants given by **George Bentham & Joseph Dalton Hooker**.

3. Phylogenetic classification systems

- It is based on evolutionary relationships among organisms.
- This assumes that organisms in the same taxa have a common ancestor.

Other sources to resolve the problems in classification:

- **Numerical Taxonomy:** It is based on all observable characteristics. It is easily carried out using computers. Number & codes are assigned to all the characters and the data are processed. Thus, hundreds of characters can be equally considered.
- **Cytotaxonomy:** It is based on cytological information like chromosome number, structure, behaviour etc.
- **Chemotaxonomy:** It uses chemical constituents of plants.

ALGAE

- Algae are simple, thalloid, autotrophic, chlorophyll-bearing and aquatic (fresh water & marine) organisms.
- They also occur in moist stones, soils and wood.
- Some occur in association with fungi (lichen) and animals (e.g., on sloth bear).
- The form and size of algae is highly variable.
 - Microscopic unicellular forms: E.g. *Chlamydomonas*.
 - Colonial forms: E.g. *Volvox*.
 - Filamentous forms: E.g. *Ulothrix* and *Spirogyra*.

Reproduction:

- **Vegetative reproduction:** By fragmentation. Each fragment develops into a thallus.
- **Asexual reproduction:** By the production of spores. E.g. **zoospores** (most common). They are flagellated (motile) and on germination gives rise to new plants.
- **Sexual reproduction:** Through fusion of two gametes. It is many types:
 - **Isogamous:** Fusion of gametes similar in size. They may be flagellated (e.g. *Ulothrix*) or non-flagellated (non-motile, e.g. *Spirogyra*).
 - **Anisogamous:** Fusion of two gametes dissimilar in size. E.g. Some species of *Eudorina*.
 - **Oogamous:** Fusion between one large, non-motile (static) female gamete and a smaller, motile male gamete. E.g. *Volvox*, *Fucus*.

Benefits of algae:


- ⊙ Through photosynthesis, they fix half of the total CO₂ on earth and increase the level of dissolved oxygen.
- ⊙ They are primary producers and the basis of the food cycles of all aquatic animals.
- ⊙ Many marine algae (70 species) are used as food. E.g. *Porphyra*, *Laminaria* and *Sargassum*.
- ⊙ **Agar** (from *Gelidium* & *Gracilaria*) is used to grow microbes and in ice-creams and jellies.

- ⊙ Some marine brown & red algae produce **hydrocolloids** (water holding substances). E.g. **algin** (brown algae) and **carrageen** (red algae). These are used commercially.
- ⊙ Protein-rich unicellular algae like *Chlorella* & *Spirulina* are used as food supplements by space travellers.

Algae include 3 classes: **Chlorophyceae**, **Phaeophyceae** and **Rhodophyceae**.

1. Chlorophyceae (green algae)

- Unicellular, colonial or filamentous.
- They are usually grass green due to the pigments **chlorophyll a** and **b** in chloroplasts.
- The chloroplasts may be discoid, plate-like, reticulate, cup-shaped, spiral or ribbon-shaped in different species.
- Most of them have one or more **pyrenoids** (storage bodies) located in the chloroplasts. Pyrenoids contain protein besides starch.
- Some algae store food as oil droplets.
- They have a rigid cell wall made of an inner layer of cellulose and an outer layer of pectose.
- E.g. *Chlamydomonas*, *Volvox*, *Ulothrix*, *Spirogyra* & *Chara*.




Reproduction:

- **Vegetative reproduction:** By fragmentation or by formation of different types of spores.
- **Asexual reproduction:** By flagellated zoospores produced in zoosporangia.
- **Sexual reproduction:** Isogamous, anisogamous or oogamous.

2. Phaeophyceae (brown algae)

- They are mostly marine forms.
- They show great variation in size & form. They range from simple branched, filamentous forms (E.g. *Ectocarpus*) to profusely branched forms (e.g. kelps- 100 m in height).
- They have chlorophyll *a*, *c*, carotenoids & xanthophylls.
- They vary in colour from olive green to brown depending upon the amount of a xanthophyll pigment, **fucoxanthin**.
- Food is stored as complex carbohydrates (**laminarin** or **mannitol**).
- The vegetative cells have a cellulosic wall covered by a gelatinous coating of **algin**.
- Protoplast contains plastids, central vacuole and nucleus.
- Plant body is attached to substratum by a **holdfast**, and has a stalk (**stipe**) and leaf like photosynthetic organ (**frond**).
- E.g. *Ectocarpus*, *Dictyota*, *Laminaria*, *Sargassum* & *Fucus*.




Reproduction:

- **Vegetative reproduction:** By fragmentation.
- **Asexual reproduction:** By pear-shaped **biflagellate zoospores** (have 2 unequal laterally attached flagella).

- **Sexual reproduction:** Isogamous, anisogamous or oogamous. Union of gametes occurs in water or within the oogonium (oogamous species). Gametes are pear-shaped (pyriform) bearing 2 laterally attached flagella.

3. Rhodophyceae (red algae)

- They have a red pigment, **r-phycoerythrin**.
- Majority are marine especially in the warmer areas.
- They occur in both well-lighted regions close to the surface of water and at great depths in oceans where relatively little light penetrates.
- The red thalli of most of the red algae are multicellular.
- Some of them have complex body organisation.
- The food is stored as **floridean starch** which is very similar to amylopectin and glycogen in structure.
- E.g. *Polysiphonia*, *Porphyra*, *Gracilaria* and *Gelidium*.



Reproduction:

- **Vegetative reproduction:** By fragmentation.
- **Asexual reproduction:** By non-motile spores.
- **Sexual reproduction:** Oogamous. By non-motile gametes. It has complex post fertilisation developments.

Classes	Chlorophyceae (Green algae)	Phaeophyceae (brown algae)	Rhodophyceae (Red algae)
Major pigments	Chlorophyll a, b	Chlorophyll a, c, Fucoxanthin	Chlorophyll a, d, Phycoerythrin
Stored food	Starch	Mannitol, laminarin	Floridean Starch
Cell wall	Cellulose	Cellulose and algin	Cellulose
Flagellar number & position of insertion	2-8, equal, apical	2, unequal, lateral	Absent
Habitat	Fresh water, salt water & brackish water	Fresh water (rare), salt water & brackish water	Fresh water (some), salt water (most) & brackish water

BRYOPHYTES

- They are called **amphibians of the plant kingdom** because they can live in soil but need water for sexual reproduction.
- They occur in damp, humid and shaded localities.
- Their body is more differentiated than that of algae. It is thallus-like and prostrate or erect, and attached to the substratum by unicellular or multicellular **rhizoids**.
- They lack true roots, stem or leaves. They may possess root-like, leaf-like or stem-like structures.
- The main plant body is haploid. It produces gametes, hence is called a **gametophyte**.
- The sex organs in bryophytes are multicellular.
- The male sex organ (**antheridium**) produces biflagellate **antherozoids**. The female sex organ (**archegonium**) is flask-shaped and produces a single **egg**.
- Antherozoids are released to water and meet archegonium. An antherozoid fuses with the egg to form **zygote**.

- Zygotes do not undergo meiosis immediately. They produce a multicellular body called a **sporophyte**.
- Sporophyte is not free-living but attached to the photosynthetic gametophyte and derives nourishment from it. Some cells of the sporophyte undergo meiosis to form haploid spores. They germinate to form gametophyte.

Importance of Bryophytes:


- ☉ Some mosses provide food for herbaceous mammals, birds and other animals.
- ☉ Species of *Sphagnum* (a moss) provide peat. It is used as fuel. It has water holding capacity so that used as packing material for trans-shipment of living material.
- ☉ They are ecologically important because of their role in **plant succession** on bare rocks/soil. Mosses along with lichens decompose rocks making the substrate suitable for the growth of higher plants.

☉ Since mosses form dense mats on the soil, they can prevent soil erosion.

The bryophytes are divided into **liverworts** and **mosses**.

Liverworts


- They grow usually in moist, shady habitats such as banks of streams, marshy ground, damp soil, bark of trees and deep in the woods.
- Their plant body is thalloid. E.g. *Marchantia*. Thallus is dorsi-ventral and closely appressed to the substrate. The leafy members have tiny leaf-like appendages in two rows on the stem-like structures.
- **Asexual reproduction:** By **fragmentation** of thalli, or by the formation of **gemmae** (sing. gemma). Gemmae are green, multicellular, asexual buds that develop in small receptacles (**gemma cups**) on the thalli. Gemmae are detached from the parent body and germinate to form new individuals.
- **Sexual reproduction:** Male and female sex organs are produced on the same or different thalli. Sporophyte is differentiated into a **foot**, **seta** and **capsule**. After meiosis, spores are produced within the capsule. These spores germinate to form free-living gametophytes.



A liverwort – *Marchantia*: Female thallus & Male thallus

Mosses

- The predominant stage of the life cycle of a moss is the **gametophyte**. It consists of two stages.
 - **Protonema stage:** The first stage which develops directly from a spore. It is a creeping, green, branched and frequently filamentous stage.
 - **Leafy stage:** The second stage which develops from the secondary protonema as a lateral bud. They consist of upright, slender axes bearing spirally arranged leaves. They are attached to soil through multicellular and branched rhizoids. This stage bears the sex organs.
- **Vegetative reproduction:** By **fragmentation** and **budding** in the secondary protonema.
- **Sexual reproduction:** The **antheridia** & **archegonia** are produced at the apex of leafy shoots. After fertilisation, zygote develops into a sporophyte, consisting of a foot, seta and capsule. The sporophyte in mosses is more elaborate than that in liverworts. The capsule contains spores. Spores are formed after meiosis. Mosses have an elaborate mechanism of spore dispersal.
- E.g. *Funaria*, *Polytrichum* and *Sphagnum*.



Funaria, gametophyte & sporophyte Sphagnum gametophyte

PTERIDOPHYTES

- They include **horsetails** and **ferns**.
- They are found in cool, damp, shady places. Some flourish well in sandy-soil conditions.
- Evolutionarily, they are the first terrestrial plants to possess vascular tissues (xylem & phloem).
- In bryophytes, the dominant phase in the life cycle is the gametophyte. In pteridophytes, the dominant phase (main plant body) is a **sporophyte**. It is differentiated to **true root, stem & leaves**. These organs have well-differentiated vascular tissues.
- The leaves in pteridophyta are small (**microphylls**) as in *Selaginella* or large (**macrophylls**) as in ferns.
- **Economic importance:** They are used for medicinal purposes and as soil-binders and ornamentals.


REPRODUCTION:

- The sporophytes bear **sporangia** that are subtended by leaf-like appendages called **sporophylls**. In some cases, sporophylls may form distinct compact structures called **strobili or cones** (E.g. *Selaginella*, *Equisetum*).
- Sporangia produce spores by meiosis in spore mother cells.

- The spores germinate to give inconspicuous, small, multicellular, free-living, mostly photosynthetic thalloid gametophytes called **prothallus**.
- Prothallus requires cool, damp, shady places to grow. Also, it needs water for fertilization. So, the spread of pteridophytes is limited and restricted to narrow geographical regions.
- The gametophytes (prothallus) bear male and female sex organs called **antheridia** and **archegonia**, respectively.
- Water is needed for transfer of **antherozoids** (male gametes from antheridia) to the mouth of archegonium.
- Antherozoid fuses with the egg in the archegonium to form zygote. Zygote develops to a multicellular well-differentiated **sporophyte**.
- Most of the pteridophytes produce similar kinds of spores (**homosporous plants**). Others produce two kinds of spores, **macro (mega)** & **micro spores**. They are **heterosporous**. E.g. *Selaginella* & *Salvinia*.
- The **megaspores** & **microspores** germinate and give rise to female and male gametophytes, respectively. The

female gametophytes are retained on the parent sporophytes for variable periods.

- Within female gametophytes, zygotes develop into young embryos. This event is a precursor to the **seed habit**. It is considered as an important step in evolution.
- The pteridophytes have 4 classes:
 1. **Psilopsida**: E.g. *Psilotum*
 2. **Lycopsidea**: E.g. *Selaginella*, *Lycopodium*
 3. **Sphenopsida**: E.g. *Equisetum*
 4. **Pteropsida**: E.g. *Dryopteris*, *Pteris*, *Adiantum*



GYMNOSPERMS

- Gymnosperms (*gymnos*: naked, *sperma*: seeds) are plants in which the **ovules are not enclosed by ovary wall** and remain exposed before and after fertilization. **Seeds** that develop post-fertilization are **not covered (naked)**.
- They include medium-sized trees or tall trees and shrubs. **Sequoia** (giant redwood) is the tallest tree species.
- The roots are generally **tap roots**.
- Roots in some genera have fungal association in the form of **mycorrhiza** (E.g. *Pinus*).
- In plants like *Cycas*, small specialized roots (**coralloid roots**) are associated with N_2 -fixing cyanobacteria.
- Stems are unbranched (*Cycas*) or branched (*Pinus*, *Cedrus*).
- Leaves are simple or compound. They are well-adapted to withstand extreme temperature, humidity and wind.
- In *Cycas*, the pinnate leaves persist for a few years.
- In conifers (*Pinus*, *Cedrus* etc.), the needle-like leaves reduce the surface area. Their thick cuticle and sunken stomata also help to reduce water loss.

REPRODUCTION:

- Gymnosperms are **heterosporous**. They produce haploid **microspores** and **megaspores**.
- Some leaves are modified into **sporophylls**. They are compactly and spirally arranged along an axis to form **lax** or **strobili** or **cones**.
- Sporophylls bear **sporangia** in which spores are produced.
- Sporophylls are 2 types:
 - o **Microsporophylls**: They are arranged to **male strobili (microsporangiate)**. They bear **microsporangia**. The

microspores develop into male gametophytes. It is highly reduced and confined to only a limited number of cells. This gametophyte is called a **pollen grain**. The pollen grains are developed within the microsporangia.

- o **Megasporophylls**: They are arranged to female strobili (**macrosporangiate**). They bear **megaspore** (ovules). Megasporangium mainly consists of a body called **nucellus**. It is protected by envelopes. The megaspore mother cell is differentiated from a cell of the nucellus. Megaspore mother cell undergoes meiosis to form four megaspores. One of the megaspores enclosed within the **Megasporangium (nucellus)** develops into a multicellular female gametophyte that bears two or more **archegonia**. The multicellular female gametophyte is also retained within megasporangium.
- The male or female cones may be borne on the same tree (*Pinus*) or on different trees (*Cycas*).
- Unlike bryophytes and pteridophytes, in gymnosperms, the male and the female gametophytes do not have an independent free-living existence. They remain within the sporangia retained on the sporophytes.
- The pollen grain released from the microsporangium are carried in air currents and meet the opening of the ovules. The pollen tube carrying the male gametes grows towards archegonia in the ovules and discharges their contents near the mouth of the archegonia.
- After fertilization, zygote develops into an embryo and the ovules into seeds.

ANGIOSPERMS (FLOWERING PLANTS)

- They are an exceptionally large group of plants.
- They range in size from tiny, almost microscopic *Wolffia* to tall trees of *Eucalyptus* (over 100 metres).
- They include 2 classes: Dicotyledons & Monocotyledons.
 - o **Dicotyledons**: Have 2 cotyledons in seeds, reticulate venations in leaves and tetramerous or pentamerous flowers (4 or 5 members in each floral whorl).
 - o **Monocotyledons**: Have only one cotyledon, parallel venation in leaves and trimerous flowers (3 members in each floral whorl).


REPRODUCTION:

- **Flower** is the reproductive structure.

- Male sex organ in a flower is the **stamen**. Each stamen consists of a **filament** with an **anther** at the tip. Within the anthers, the **pollen mother cell** divides by meiosis to produce **microspores** which matures into **pollen grains**.
- Female sex organ in a flower is the **pistil**. It consists of a swollen **ovary** at its base, a long slender **style & stigma**. Ovary contains **ovules**. An ovule has a **megaspore mother cell** that undergoes meiosis to form 4 haploid **megaspores**. 3 of them degenerate and one divides to form **embryo sac**.
- Each embryo-sac has a 3-celled **egg apparatus** (one **egg cell** & two **synergids**), 3 **antipodal cells** & 2 **polar nuclei**. The polar nuclei eventually fuse to produce a **diploid secondary nucleus**.

- Pollen grains dispersed from anthers are carried by wind or other agencies to the stigma of pistil. It is called **pollination**.
- Pollen grains germinate on the stigma and the resulting **pollen tubes** grow through the tissues of stigma and style and reach the ovule.
- Pollen tubes enter the embryo-sac where 2 male gametes are discharged. One male gamete fuses with egg cell to form **zygote (syngamy)**. The other male gamete fuses with diploid secondary nucleus to produce triploid **primary endosperm nucleus (PEN)**. Because of the involvement of two fusions, this event is called **double fertilisation**. It is an event unique to angiosperms.
- The zygote develops into an **embryo** (with one or two cotyledons). The PEN develops into **endosperm** which provides nourishment to the developing embryo.
- Synergids & antipodals degenerate after fertilization.
- During these events, the **ovules** develop into **seeds** and the **ovaries** develop into **fruit**.

- The seeds are enclosed by fruits.



PLANT LIFE CYCLES AND ALTERNATION OF GENERATIONS




- In plants, both haploid and diploid cells can divide by mitosis. This forms haploid and diploid plant bodies.
- Haploid plant body (**gametophyte**) produces gametes by mitosis.
- After fertilization, the zygote also divides by mitosis to produce a diploid plant body (**sporophyte**). This produces haploid **spores** by meiosis.
- Spores divide by mitosis to form a haploid plant body.
- Thus, during the life cycle of any sexually reproducing plant, there is an alternation of generations between gametophyte (n) and sporophyte (2n).

Zygote undergoes meiosis to form haploid spores. They divide mitotically to form gametophyte. The dominant, photosynthetic phase is the free-living gametophyte. E.g. Algae such as *Volvox*, *Spirogyra* and some species of *Chlamydomonas*.

- Diplontic:** In this, diploid sporophyte is the dominant, photosynthetic, independent phase. Gametophytic phase is represented by the single to few-celled haploid gametophyte. E.g. An alga, *Fucus* sp., all seed-bearing plants (gymnosperms & angiosperms - the gametophytic phase is few to multi-celled).
- Haplo-diplontic:** It is the intermediate condition between haplontic & diplontic. Both gametophyte & sporophyte are multicellular and often free-living. But they have different dominant phases. E.g. Bryophytes & Pteridophytes.

Patterns of Plant life cycles

- Haplontic:** In this, sporophytic generation is represented only by the zygote. There are no free-living sporophytes.



MODEL QUESTIONS

1. Match the names given in column I with the items in column II

Column I

- a) Adiantum
- b) Sargassum
- c) Sunflower
- d) Riccia

Column II

- Bryophyte
- Angiosperm
- Algae
- Pteridophyte

2. Bryophytes are called 'amphibians of the plant kingdom'. Justify the statement.
3. Match the following.

A

- a. Algae
- b. Pteridophytes
- c. Gymnosperms
- d. Angiosperms
- e. Bryophytes

B

- Naked seeded plants.
- Amphibians of plant kingdom
- Flowering plants
- Thalloid body, photosynthetic and mainly aquatic
- Independent sporophytic and gametophytic phase. Commonly called ferns.

4. The most characteristic feature of angiosperms is double fertilization.
a. What is double fertilization? b. Why is it called triple fusion?
5. Identify the plant group from the given data.
a. Plant body is not differentiated into root, stem and leaf.
b. Reproductive structures are strobili.
6. Some Pteridophytes produce two types of spores.
a. Name the phenomenon. b. This event is an important step in evolution. Explain.
7. Some features of Haplontic and Diplontic life cycles are given below. Arrange them correctly in a table.
- Sporophyte is a single celled zygote
 - Meiosis occurs in Zygote
 - Sporophyte dominant
 - No free-living Sporophyte
 - Gametophyte dominant
 - Meiosis occurs before zygote formation

4. ANIMAL KINGDOM

Animals are **multicellular** and **heterotrophic** organisms **without cell wall** and **chlorophyll**.

Kingdom Animalia includes 11 major phyla:

- | | |
|--------------------|------------------|
| 1. Porifera | 7. Arthropoda |
| 2. Cnidaria | 8. Mollusca |
| 3. Ctenophora | 9. Echinodermata |
| 4. Platyhelminthes | 10. Hemichordata |
| 5. Aschelminthes | 11. Chordata |
| 6. Annelida | |

BASIS OF CLASSIFICATION

1. Levels of organization

Based on this, animals are grouped into four levels:

- Cellular level of organization:** Here, the cells are arranged as loose cell aggregates. E.g. Porifera.
- Tissue level of organization:** Here, the cells are arranged into **tissues**. E.g. Cnidarians and Ctenophores.
- Organ level of organization:** Here, tissues are arranged into **organs**. E.g. Higher animals (Platyhelminthes to chordates).
- Organ system level of organization:** Here, organs are associated to **organ system**. Each system performs a specific physiological function. E.g. Higher animals.

Organ systems of various animals show complexities. E.g.

Digestive system is 2 types:

- **Incomplete:** It has only a single opening that acts as mouth & anus. Seen in Cnidaria and Platyhelminthes.
- **Complete:** It has 2 openings (mouth & anus).

Circulatory system is 2 types: **open & closed**.

2. Body symmetry

It is the arrangement of similar body parts on 2 sides of main axis of the body. Based on symmetry, animals are 2 types: Asymmetrical and Symmetrical.

- Asymmetrical:** Here, body cannot be divided into 2 equal halves. E.g. Most Poriferans, Snails etc.
- Symmetrical:** Here, body can be divided into 2 equal halves. It is 2 types.
 - **Radial symmetry:** Here, body can be divided into 2 equal halves in **any vertical plane** along **central axis** (oral-aboral axis) of the body. E.g. some Poriferans, Cnidarians, Ctenophores and Echinoderms (adult).
 - **Bilateral symmetry:** Here, body can be divided into **right & left halves** in only **one plane**. E.g. Platyhelminthes to Chordata (except adult Echinodermata).


The body of bilaterally symmetrical animal has a **dorsal side** (upper), a **ventral side** (lower), left & right **lateral sides**, **anterior** (cephalic) side and **posterior** (anal or tail) side.

3. Germinal layers (Embryonic layers)

These are layers of embryo from which all the body organs are formed. Based on the number of germ layers, animals are 2 types- Diploblastic and Triploblastic.

a. Diploblastic animals: 2 germ layers- outer ectoderm and inner endoderm. In between these layers, an undifferentiated jelly-like layer called **mesoglea** is present. E.g. Cnidaria & Ctenophora.

b. Triploblastic animals: 3 germ layers- Outer ectoderm, middle mesoderm and inner endoderm. E.g. Platyhelminthes to Chordata.




4. Coelom (body cavity)

It is the cavity lined by mesoderm. It is seen between body wall and gut wall. Coelom separates the muscles of gut and body wall.

Based on the nature of coelom, animals are 3 types:

- Acoelomate:** No coelom. The space between body wall and digestive cavity is filled with matrix (parenchyma). E.g. Porifera to Platyhelminthes.
- Pseudocoelomate:** False coelom. Here, the body cavity is not lined by mesoderm. Mesoderm is scattered pouches. E.g. Aschelminthes.
- Coelomate:** True coelom. Here, the coelom arises from the mesoderm. Coelom is lined by peritoneal layer and filled with coelomic fluid. E.g. Annelida to Chordata.



Functions of coelom:

- It accommodates visceral organs.
- Coelomic fluid reduces friction between visceral organs.
- It acts as shock absorber.

5. Metamerism (segmentation)

It is the phenomenon in which the body or organs is externally and internally divided into **repeated segments (metameres)**. E.g. Annelids (earthworm etc.), Arthropods.


6. Notochord

It is a **mesodermally derived supporting rod** formed on the dorsal side during embryonic development in some animals. Animals with notochord are called **chordates** and those without notochord are called **non-chordates**.

GENERAL CHARACTERS OF DIFFERENT PHYLA (NON-CHORDATES)

Features	Porifera (Sponges)	Cnidaria (Coelenterata)	Ctenophora (Comb jellies or Sea walnuts)
Grades of organization	Cellular	Tissue	Tissue
Symmetry	Asymmetrical. Some are radial.	Radial	Radial
Germ layers	-	Diploblastic	Diploblastic
Coelom	Acoelomate	Acoelomate	Acoelomate
Habit and habitat	Aquatic (mostly marine). Sedentary. Solitary/colonial.	Aquatic (mostly marine). Sessile/free swimming. Solitary/colonial.	Exclusively marine. Solitary & pelagic
Digestive system	Absent. Intracellular digestion.	Incomplete. Intracellular & extracellular digestion.	Incomplete. Intracellular and extracellular digestion.
Respiratory system	Absent	Absent	Absent
Circulatory system	Absent	Absent	Absent
Reproduction	Asexual (fragmentation) & Sexual. Hermaphrodite. Internal fertilization. Development is indirect.	Polyp reproduces asexually (budding) and medusa sexually. Most are separate sexes. External fertilization. Development is indirect.	Only Sexual. Hermaphrodite. External fertilization. Development is indirect.
Unique features	Water canal (water transport) system. Millions of ostia (pores) . Spongocoel & canals are lined with choanocytes (collar cells). Body is supported by spicules and spongin fibres.	Tentacles with cnidoblasts. Gastro-vascular cavity (coelenteron) with an opening (mouth) on hypostome . Polyp & Medusa forms are seen. Some shows alternation of generation . Corals have skeleton (CaCO ₃).	Locomotion is by 8 vertical external rows of ciliated comb plates . Tentacles present. Shows Bioluminescence .
Examples	<i>Sycon</i> (<i>Scypha</i>), <i>Spongilla</i> (fresh water sponge), <i>Euspongia</i> (Bath sponge)	<i>Hydra</i> , <i>Obelia</i> , <i>Aurelia</i> , <i>Physalia</i> (Portuguese man of war), <i>Adamsia</i> (Sea-anemone), <i>Pennatulata</i> (Sea pen), <i>Gorgonia</i> (Sea fan), <i>Meandrina</i> (Brain coral)	<i>Ctenoplana</i> , <i>Pleurobrachia</i>

- Water canal system:** Here, water enters through minute pores (**ostia**) in the body wall into a central cavity (**spongocoel**), from where it goes out through **osculum**. Canal system is used for food gathering, gas exchange and removal of wastes.
- Hermaphrodite (Monoecious):** Male and female sex organs are seen in same individual.
- Tentacles:** Finger-like structures which surrounds the mouth of coelenterates. Used for food capture & defense.
- Cnidoblasts (Cnidocytes):** These are stinging cells (present on the tentacles and the body) with a poison-filled capsule called **nematocyst**. Cnidoblast is used for **anchorage, defense** and to **capture prey**.
- Polyp & Medusa:** 2 types of body forms in cnidarians.
Polyp is tubular attached asexual form, with upwardly directed mouth & tentacles. E.g. *Hydra*, *Adamsia*.
Medusa is umbrella like, free-swimming sexual form, with downwardly directed mouth & tentacles. E.g. *Aurelia* (Jelly fish).
- Alternation of generation (Metagenesis):** The phenomenon in which polyps produce medusae asexually and medusae form the polyps sexually. E.g. *Obelia*.
- Bioluminescence:** It is the property of some animals to emit light from the body.



GENERAL CHARACTERS OF DIFFERENT PHyla (NON-CHORDATES)

Features	Platyhelminthes (Flatworms)	Aschelminthes (Roundworms)	Annelida (Segmented or Ringed worms)	Arthropoda (Joint-legged animals)
Grades of organization	Organ & Organ system	Organ system	Organ system	Organ system
Symmetry	Bilateral	Bilateral	Bilateral	Bilateral
Germ layers	Triploblastic	Triploblastic	Triploblastic	Triploblastic
Coelom	Acoelomate	Pseudocoelomate	Coelomate	Coelomate
Habit and habitat	Mainly aquatic. Endoparasites. Some are free-living.	Aquatic and terrestrial. Free living or parasitic in plants & animals.	Terrestrial, fresh water or marine. Free living or parasitic.	Cosmopolitan
Digestive system	Incomplete	Complete. Tubular alimentary canal with well-developed muscular pharynx.	Complete	Complete
Respiratory system	Absent	Absent	Cutaneous respiration. Some have branchial (gill) respiration.	Gills/ book gills/ trachea/book lungs
Circulatory system	Absent	Absent	Closed type	Open type
Reproduction	Asexual (fragmentation) and Sexual. Hermaphrodite. Internal Fertilization. Development is indirect. Many larval stages.	Dioecious. Sexual reproduction. Internal fertilization. Development is direct or indirect.	Sexual. Earthworms & leeches are monoecious. <i>Neries</i> is dioecious. Development is direct or indirect.	Mostly dioecious. Usually internal fertilization. Mostly oviparous . Development is direct or indirect.
Unique features	Unsegmented, dorso-ventrally flattened body (except tape worms). Excretion and osmo-regulation by Flame cells (protonephridia) . Parasites have Hooks & suckers . Some absorb nutrients from the host through their body surface.	Body is circular in cross section. Syncytial epidermis. Thick cuticle. An excretory tube to remove body waste through excretory pore. Sexual dimorphism (females are longer than males).	Segmentation like rings. Longitudinal and circular muscles help in locomotion. Locomotory organs are setae (in earthworm) or parapodia (in <i>Neries</i>). Excretion by Nephridia . Paired ganglia connected by lateral nerves to a double ventral nerve cord.	Jointed appendages . Body has 3 regions: head, thorax & abdomen . Body is covered by chitinous cuticle (exoskeleton) . Excretion by Malpighian tubules . Sensory organs are antennae, compound & simple eyes, statocysts (balance organs) .
Examples	<i>Taenia solium</i> (Tape worm), <i>Fasciola</i> (Liver fluke), <i>Planaria</i> (shows high regeneration capacity).	<i>Ascaris</i> (Roundworm), <i>Ancylostoma</i> (Hookworm), <i>Wuchereria</i> (Filarial worm).	<i>Pheretima</i> (earthworm), <i>Hirudinaria</i> (blood sucking Leech), <i>Neries</i> (aquatic). Parapodia for swimming).	Spider, Scorpion, Crab, Prawn, Insects etc. Economically important insects: <i>Apis, Bombyx, Laccifer</i> . Vectors: Mosquitoes (<i>Anopheles, Culex & Aedes</i>), Housefly etc. Gregarious pest: <i>Locusta</i> . Living fossil: <i>Limulus</i> (King crab)

1. **Dioecious:** Sexes are separate.
2. **Sexual dimorphism:** Morphological differences between male and female.
3. Arthropoda is the largest phylum. Over two-thirds of all named species are arthropods.

GENERAL CHARACTERS OF DIFFERENT PHYLA (NON-CHORDATES)

Features	Mollusca (Soft-bodied animals)	Echinodermata (Spiny-skinned animals)	Hemichordata
Grades of organization	Organ system	Organ system	Organ system
Symmetry	Bilateral	Adults radial. Larvae bilateral.	Bilateral
Germ layers	Triploblastic	Triploblastic	Triploblastic
Coelom	Coelomate	Coelomate	Coelomate
Habit and habitat	Aquatic. Few are terrestrial.	Exclusively marine.	Exclusively marine.
Digestive system	Complete	Complete. Ventral mouth and dorsal anus.	Complete
Respiratory system	Gills in aq. forms and pulmonary sac in terrestrial forms.	Dermal branchiae (skin gills or papulae) and tube feet.	Gills
Circulatory system	Open type	Open type	Open type
Reproduction	Dioecious. Oviparous. Development is direct or indirect.	Dioecious. External fertilization. Development is indirect. Ciliated free-swimming larva.	Dioecious. External fertilization. Development is indirect.
Unique features	Body has head, visceral mass (visceral hump) & muscular foot. Head has sensory tentacles. Calcareous shell. Feather-like gills for respiration & excretion. Mantle & radula are seen.	They have an endoskeleton of calcareous ossicles (Spiny bodied). Head absent. Water vascular system present. Excretory system absent. Shows autotomy & regeneration.	Worm-like cylindrical body formed of anterior proboscis, a collar and a long trunk. Collar bears stomochord (a rudimentary structure similar to notochord). Excretion by Proboscis gland.
Examples	<i>Pila</i> (Apple Snail), <i>Pinctada</i> (Pearl Oyster), <i>Sepia</i> (Cuttlefish), <i>Loligo</i> (Squid), <i>Octopus</i> (Devil fish), <i>Aplysia</i> (Sea Hare), <i>Dentalium</i> (Tusk shell), <i>Chaetopleura</i> (Chiton)	<i>Asterias</i> (Starfish), <i>Echinus</i> (Sea Urchin), <i>Echinocardium</i> , <i>Antedon</i> (Sea Lily), <i>Cucumaria</i> (Sea Cucumber), <i>Ophiura</i> (Brittle Star)	<i>Balanoglossus</i> (Tongue worm), <i>Saccoglossus</i>


- Mollusca** is the second largest phylum.
- Mantle (Pallium):** The membrane which covers visceral mass. Space between the hump and mantle is called *mantle cavity*.
- Radula:** File-like rasping organ present in the mouth of molluscs. It is used for feeding.
- Water vascular (ambulacral) system:** In this system, sea water enters through a porous plate called *madreporite* and reaches the radiating canals and *tube feet (podia)*. Its functions are locomotion, respiration, capture & transport of food and excretion.
- Hemichordata was earlier considered as a sub-phylum of Chordata. Like chordates, it has pharyngeal gill slits.

PHYLUM CHORDATA

It includes animals with notochord, dorsal tubular nerve cord and pharyngeal gill slits.

Notochord is a flexible rod located in the mid dorsal line between the alimentary canal and the nerve cord in the embryo.

Differences between Chordata and Non-Chordata




**Chordata characteristics
(Body plan)**

Chordata	Non-Chordata
1. Notochord is found in the embryonic stage	Absent
2. Central nervous system is dorsal, hollow and single	Ventral, solid and double
3. Pharyngeal gill slits present	Absent
4. Ventral heart	Dorsal heart (if present)
5. A post-anal part (tail) is present	Absent

Phylum Chordata is classified into 3 subphyla: **Urochordata, Cephalochordata & Vertebrata.**

PROTOCHORDATA (ACRANIATA)		VERTEBRATA (CRANIATA)
Urochordata (Tunicata)	Cephalochordata	
<ul style="list-style-type: none"> • Notochord present only in larval tail. • Body is covered by test made up of tunicin. • Exclusively marine. • Hermaphrodite. • E.g. <i>Ascidia</i>, <i>Salpa</i>, <i>Doliolum</i>. 	<ul style="list-style-type: none"> • Notochord from head to tail region and is persistent throughout the life. • Fish-like body. • Exclusively marine. • Sexes are separate. • E.g. <i>Branchiostoma</i> (Amphioxus or Lancelet). 	<ul style="list-style-type: none"> • Possess notochord during the embryonic period. • Notochord is replaced by a cartilaginous or bony vertebral column in the adult. • Ventral muscular heart. • Kidneys for excretion & osmoregulation • Paired appendages (fins or limbs).



CLASS CYCLOSTOMATA

- All are *ectoparasites* on some fishes.
- Elongated body without scales and paired fins.
- 6-15 pairs of *gill slits* for respiration.
- Sucking and circular mouth *without jaws*.
- *Cartilaginous cranium* and *vertebral column*.
- Circulation is *closed* type.
- Marine, but migrate for *spawning* to fresh water. After spawning, they die. Their larvae, after metamorphosis, return to ocean.
- E.g. *Petromyzon* (Lamprey) and *Myxine* (Hagfish).

SUPERCLASS PISCES (FISHES)

Class Chondrichthyes (Cartilaginous fishes)	Class Osteichthyes (Bony fishes)
Marine. Stream-lined body. Predaceous.	Marine & fresh water. Stream-lined body.
Cartilaginous endoskeleton. Notochord is persistent throughout life.	Bony endoskeleton.
Ventral mouth.	Terminal mouth.
Gill slits without operculum. Powerful jaws.	4 pairs of gills covered by operculum on each side.
Skin with placoid scales . Teeth are modified placoid scales which are backwardly directed.	Scales are Cycloid, ctenoid etc.
No air bladder . So, they have to swim constantly to avoid sinking.	Air bladder for buoyancy.
Poikilotherms (cold-blooded).	Poikilotherms (cold-blooded).
Two-chambered heart (one auricle and one ventricle).	Two-chambered heart (one auricle and one ventricle).
Sexes are separate. In males, pelvic fins bear claspers . Internal fertilization. Many of them viviparous .	Sexes are separate. External fertilisation. Mostly oviparous . Development is direct.
Examples <i>Scoliodon</i> (Dogfish), <i>Pristis</i> (Saw fish), <i>Carcharodon</i> (Great white shark), <i>Trygon</i> (Sting ray- has poison sting), <i>Torpedo</i> (Electric ray- has <i>electric organ</i>).	Examples Marine: <i>Exocoetus</i> (flying fish), <i>Hippocampus</i> (seahorse) Fresh water: <i>Labeo</i> (Rohu), <i>Catla</i> (Katla), <i>Clarias</i> (Magur). Aquarium: <i>Betta</i> (Fighting fish), <i>Pterophyllum</i> (Angel fish).


SUPERCLASS TETRAPODA

Class Amphibia	Class Reptilia	Class Aves (Birds)	Class Mammalia
They live in aquatic & terrestrial habitats and need water for breeding.	Dry & cornified skin, epidermal <i>scales</i> or <i>scutes</i> .	Presence of <i>feathers</i> and <i>beak</i> . Forelimbs are modified into <i>wings</i> .	Presence of <i>mammary glands</i> (milk producing glands).
Body has head & trunk. Some have tail. Moist skin without scales. Most have 2 pairs of limbs.	Snakes and lizards shed their scales as <i>skin cast</i> . Limbs- 2 pairs (if present). Crawling mode of locomotion.	Dry skin without glands except the <i>oil gland</i> at the base of tail. Hind limbs have <i>scales</i> and are modified for walking, swimming or clasp tree branches. Hollow & pneumatic long bones.	Skin with <i>hair</i> . 2 pairs of limbs for walking, running, climbing, burrowing, swimming or flying.
<i>Tympanum</i> represents ear.	<i>Tympanum</i> represents ear.	<i>Tympanum</i> represents ear.	External ear (<i>Pinnae</i>).
3-chambered heart (2 auricles + 1 ventricle).	3-chambered heart (but a septum partially separates ventricle). Heart is 4-chambered in crocodiles.	4-chambered heart.	4-chambered heart.
Poikilotherms	Poikilotherms	Homoiotherms	Homoiotherms
Alimentary canal, urinary & reproductive tracts open into a <i>Cloaca</i> which opens to exterior.	Well-developed alimentary canal.	Digestive tract has additional chambers, the <i>crop</i> & <i>gizzard</i> .	Well-developed alimentary canal. Dentition is <i>Heterodont</i> , <i>thecodont</i> & <i>diphyodont</i> .
Respiration is by <i>gills</i> (in larva), <i>lungs</i> & <i>skin</i>	Respiration by <i>lungs</i> .	Double respiration. <i>Air sacs</i> connected to lungs.	Respiration by <i>lungs</i> .
Sexes are separate. External fertilisation. Oviparous . Development is indirect.	Internal fertilisation. Oviparous . Development is direct.	Internal fertilisation. Oviparous . Development is direct.	Sexes are separate. Internal fertilisation. Viviparous (except <i>Echidna</i> and <i>Platypus</i>). Development is direct.
Examples	Examples	Examples	Examples
<i>Bufo</i> (Toad), <i>Rana</i> (Frog), <i>Hyla</i> (Tree frog), <i>Salamandra</i> (Salamander), <i>Ichthyophis</i> (Limbless amphibia)	<i>Chelone</i> (Turtle), <i>Testudo</i> (Tortoise), <i>Chameleon</i> (Tree lizard), <i>Calotes</i> (Garden lizard), <i>Crocodylus</i> (Crocodile), <i>Alligator</i> , <i>Hemidactylus</i> (Wall lizard). Poisonous snakes: <i>Naja</i> (Cobra), <i>Bangarus</i> (Krait), <i>Vipera</i> (Viper) etc. Non-poisonous snakes: <i>Python</i> etc.	<i>Corvus</i> (Crow), <i>Columba</i> (Pigeon), <i>Psittacula</i> (Parrot), <i>Struthio</i> (Ostrich), <i>Pavo</i> (Peacock), <i>Aptenodytes</i> (Penguin), <i>Neophron</i> (Vulture) etc.	<i>Ornithorhynchus</i> (Platypus), <i>Macropus</i> (Kangaroo), <i>Pteropus</i> (flying fox), <i>Camelus</i> (Camel), <i>Macaca</i> (Monkey), <i>Rattus</i> (Rat), <i>Canis</i> (dog), <i>Felis</i> (Cat), <i>Elephas</i> (Elephant), <i>Equus</i> (Horse), <i>Delphinus</i> (Common dolphin), <i>Balaenoptera</i> (blue whale), <i>Panthera tigris</i> (Tiger), <i>Panthera leo</i> (lion)

- **Poikilotherms (Cold-blooded animals):** Animals that lack the capacity to regulate their body temperature.
- **Homoiotherms (warm-blooded animals):** Animals having ability to maintain a constant body temperature.

MODEL QUESTIONS

1. Observe the diagram showing the water canal system of sponges and complete the given flow chart.



Surrounding sea water → A → B → C → Surrounding sea water

2. Observe the diagram and answer the following questions:



- Identify the two forms.
- Name the phylum in which this phenomenon is seen.
- Mention any two differences between them.
- Of these, which is more advanced? Give reason.

3. Note the relation between first two words and suggest a suitable word for the fourth place

- | | | | |
|--------------------|---------------|-----------------|---------|
| a. Annelida | : Nephridia | Platyhelminthes | : |
| b. Platyhelminthes | : Acoelomates | Pseudocoelomate | : |
| c. Fish | : Gills | Insects | : |

4. Odd man out and give reason

- Ascaris*, *Wuchereria*, *Ancylostoma*, *Limulus*
- Earthworm, hookworm, roundworm, filarial worm
- Flying fish, Sea horse, Hag fish, Angel fish


5. Match the following related things from B and C with column A

A	B	C
Jelly fish	Sepia	Annelida
Cuttlefish	Hirudinaria	Echinodermata
Sea urchin	Aurelia	Mollusca
Leech	Echinus	Cnidaria

6. Find out the corresponding phylum with an example from the following general characters

- They are multicellular with a tissue grade of organisation, all are aquatic, radially symmetrical, diploblastic and mouth is encircled by finger like tentacles with stinging cells.
- They are bilaterally symmetrical, vermiform animals, triploblastic, exhibit true metamerism.
- They are exclusively marine, triploblastic, spines on the skin and radially symmetrical in the adult and bilaterally symmetrical in the larval stage.

7. Observe the following diagram.



- Identify the diagram.
- Copy the diagram and label A, B, C & D.

8. Give reasons for the following statements

- All vertebrates are chordates but all chordates are not vertebrates.
- Shark has to swim continuously, otherwise it will sink down.

9. Study the following facts by connecting with the features of tetrapoda and answer the following

- | | | |
|--------------------------------------|------------------------|--------------------------|
| • Aq. larval life & terr. adult life | • Mammary glands | • Poikilotherms |
| • Bear limbs | • Presence of feathers | • Oviparous |
| • Heterodont dentition | • Caudal fin | • Skin with hair |
| • Heart is 4-chambered | • Bony endoskeleton | • Dry and cornified skin |
| • Pneumatic bone | • Operculum | • Notochord present |

- Select the common features of all tetrapods.
- Select the unique features of different tetrapods.
- Mention the features which are not suitable for tetrapods.

5. MORPHOLOGY OF FLOWERING PLANTS


- **Morphology** is the study of external forms of organisms.
- A flowering plant (Angiosperm) has 2 parts: **Root system** (underground part) & **Shoot system** (portion above the ground).

THE ROOT


It is the underground part formed from **radicle** of embryo.

Root systems are 3 types:

- **Tap root system:** It consists of **primary roots (tap root)** and its branches (lateral roots such as **secondary roots, tertiary roots**). Seen in dicots. Primary root is elongated from **radicle**. E.g. Mustard plant.
- **Fibrous root system:** In monocots, primary root is short lived and is replaced by many roots. They originate from the base of stem to form fibrous root system. E.g. Wheat.
- **Adventitious root system:** Roots that arise from parts other than radicle. E.g. Grass, *Monstera* and banyan tree.



- **Region of maturation:** It is proximal to elongation zone. Here, the cells differentiate and mature.
- **Root hairs:** Very fine, delicate, thread-like structures formed from epidermal cells in region behind region of elongation. They absorb water and minerals from the soil.



Modifications of Root

In some plants, roots are modified to perform functions other than absorption and conduction. E.g.

- **Swollen roots for food storage:** E.g. Tap roots of carrot, turnips and adventitious roots of sweet potato.
- **Prop roots:** Hanging structures that support banyan tree.
- **Stilt roots:** The supporting roots coming out of the lower nodes of the stem. E.g. maize & sugarcane.
- **Pneumatophores:** The roots that come out of the ground and grow vertically upwards to get oxygen for respiration. E.g. *Rhizophora* growing in swampy areas.

Functions of root

- ☉ Absorption of water and minerals from the soil.
- ☉ Provide a proper anchorage to the plant parts.
- ☉ Storage of reserve food material.
- ☉ Synthesis of plant growth regulators.

Regions of the Root

- **Root cap:** It is the covering at the apex of root. It protects the tender apex of the root.
- **Region of meristematic activity:** Seen above the root cap. Here, the cells are very small, thin-walled and with dense protoplasm. They divide repeatedly.
- **Region of elongation:** Region just above the meristematic region. Here, cells undergo rapid elongation and enlargement. Helps in growth of the root in length.

THE STEM

- It is the ascending part of the axis that develops from the **plumule** of the embryo of a germinating seed.
- It bears branches, leaves, flowers, fruits, buds (terminal or axillary), nodes and internodes.
- **Nodes** are the regions of the stem where leaves are born. **Internodes** are the portions between two nodes.
- Young stem is generally green and later often become woody and dark brown.

Functions of stem:

- ☉ Spreading out branches bearing leaves, flowers and fruits.
- ☉ It conducts water, minerals and photosynthates.
- ☉ Food storage, support, protection & vegetative propagation.

Modifications of Stem

- **For food storage:** E.g. underground stems of potato, ginger, turmeric, *zaminkand*, *Colocasia* etc. They also act as organs of perennation to tide over conditions unfavourable for growth.
- **Stem tendrils:** Slender and spirally coiled structures formed from axillary buds. They help plants to climb. E.g. Gourds (cucumber, pumpkins, watermelon) & grapevines.


- **Thorns:** Woody, straight and pointed structures developed from axillary buds. They protect plants from browsing animals. E.g. *Citrus*, *Bougainvillea*.
 - **Phylloclade:** It is a green, flattened or fleshy cylindrical stem containing chlorophyll for photosynthesis. Found in some plants of arid regions. E.g. *Opuntia* (flattened stem), *Euphorbia* (cylindrical stem).
 - **Stolon:** Slender lateral branch that arises from the base of the main axis and after growing aurally for some time arch downwards to touch the ground. E.g. mint & jasmine.
 - **Offset:** It is a lateral branch with short internodes and each node bearing a rosette of leaves and a tuft of roots. E.g. aquatic plants like *Pistia* and *Eichhornia*.
 - **Sucker:** The lateral branches that originate from the basal underground part of the main stem. It grows horizontally beneath the soil and come out obliquely upward giving rise to leafy shoots. E.g. Banana, Pineapple & *Chrysanthemum*.
- Underground stems of grass, strawberry etc. spread to new niches. When older parts die, new plants are formed.

THE LEAF

- It is a lateral, flattened structure borne on the stem.
- It develops at the node and bears a bud in its axil.
- The **axillary bud** later develops into a branch.
- Leaves originate from shoot apical meristems and are arranged in an acropetal order.
- They are important vegetative organs for photosynthesis.

A typical leaf has 3 main parts:

- **Leaf base:** With this, the leaf is attached to stem. It may bear two lateral small leaf-like structures called **stipules**. In monocots, the leaf base expands into a sheath covering the stem partially or wholly. In some leguminous plants, the leaf base may be swollen. It is called **pulvinus**.
- **Petiole:** It helps to hold the leaf blade to light. Long thin flexible petioles allow leaf blades to flutter in wind, thereby cooling leaf and bringing fresh air to leaf surface.
- **Lamina (leaf blade):** The green expanded part with veins & veinlets. The middle prominent vein is called **midrib**. Veins provide rigidity to lamina and act as channels of transport for water, minerals & food materials.




Venation

- It is the arrangement of veins and veinlets in leaf lamina.
- It is 2 types:
 - **Reticulate venation:** Here, the veinlets form a network. It is seen in dicotyledons.
 - **Parallel venation:** Here, the veins run parallel to each other within a lamina. It is seen in monocotyledons.

Types of Leaves

- **Simple leaf:** Here, leaf lamina is entire or when incised, the incisions do not touch the midrib.
- **Compound leaf:** Here, the incisions of the lamina reach up to the midrib breaking it into several leaflets. A bud is seen in the axil of petiole in simple & compound leaves, but not in the axil of leaflets of the compound leaf. The compound leaves are 2 types.
 - **Pinnately compound leaf:** In this, many leaflets are present on a common axis, the **rachis**, which represents the midrib of the leaf. E.g. neem.


- **Palmately compound leaf:** In this, leaflets are attached at a common point (at the tip of petiole). E.g. silk cotton.



Phyllotaxy

It is the pattern of arrangement of leaves on the stem or branch. It is 3 types:

- **Alternate:** In this, a single leaf arises at each node in alternate manner. E.g. China rose, mustard & sun flower.
- **Opposite:** In this, a pair of leaves arise at each node and lie opposite to each other. E.g. *Calotropis* and guava.
- **Whorled:** In this, more than two leaves arise at a node and form a whorl. E.g. *Alstonia*.




Modifications of Leaves

- Leaves are modified to perform functions other than photosynthesis. E.g.
 - **Tendrils:** For climbing. E.g. peas.
 - **Spines:** For defense. E.g. cacti.
 - **Fleshy leaves:** To store food. E.g. onion and garlic.
- In plants such as Australian acacia, the leaves are small and short-lived. The petioles in these plants expand, become green and synthesise food.
- Leaves of some insectivorous plants (e.g. pitcher plant, Venus-fly trap) are also modified leaves.

THE FLOWER AND THE INFLORESCENCE

- A flower is a modified shoot wherein the shoot apical meristem changes to floral meristem.
- Internodes do not elongate and the axis gets condensed.
- The apex produces different kinds of floral appendages laterally at successive nodes instead of leaves.
- When a shoot tip transforms into a flower, it is solitary.
- The arrangement of cluster of flowers on the floral axis is called **inflorescence**.
- Based on whether the apex gets converted into a flower or continues to grow, inflorescences are 2 types: Racemose and Cymose.



- **Racemose:** In this, the main axis continues to grow. Flowers are borne laterally in an acropetal succession.

- **Cymose:** In this, main axis terminates in a flower, hence is limited in growth. Flowers are borne in a basipetal order.

THE FLOWER

- It is the **reproductive unit** in the angiosperms.
- It is meant for sexual reproduction.
- A flower has a **stalk (pedicel)**. Its swollen end is called **thalamus (receptacle)**.
- Reduced leaf found at the base of the pedicel is called **bracts**. Flowers with bracts are called **bracteate** and those without bracts, **ebracteate**.
- A typical flower has 4 kinds of whorls arranged on thalamus- **calyx, corolla, androecium & gynoecium**.
- Calyx & corolla are accessory organs, while androecium and gynoecium are reproductive organs.
- In flowers like lily, the calyx and corolla are not distinct. It is termed as **perianth**.
- When a flower has both androecium and gynoecium, it is **bisexual**. A flower having either only androecium or only gynoecium is **unisexual**.

Based on symmetry, flowers are 3 types:


- **Actinomorphic** (radial symmetry): Here, a flower can be divided into 2 equal radial halves in any radial plane passing through the centre. E.g. mustard, *datura*, chilli.
- **Zygomorphic** (bilateral symmetry): Here, a flower can be divided into two similar halves only in a particular vertical plane. E.g. pea, gulmohur, bean, *Cassia*.
- **Asymmetric** (irregular): Here, a flower cannot be divided into two similar halves by any vertical plane passing through the centre. E.g. canna.

Based on number of floral appendages, flowers are classified as follows:

- **Trimerous:** Floral appendages are multiple of 3.
- **Tetramerous:** Floral appendages are multiple of 4.
- **Pentamerous:** Floral appendages are multiple of 5.

Based on the position of calyx, corolla and androecium in respect of the ovary on thalamus, the flowers are 3 types:

- **Hypogynous:** Here, gynoecium occupies the highest position while other parts are situated below it. The ovary is **superior**. E.g. mustard, China rose & brinjal.
- **Perigynous:** Here, gynoecium is situated in the centre and other parts are located on the rim of the thalamus at the same level. Ovary is **half inferior**. E.g. plum, rose, peach.
- **Epigynous:** Here, the margin of thalamus grows upward enclosing the ovary completely and getting fused with it. Other parts arise above the ovary. The ovary is **inferior**. E.g. Guava, cucumber, ray florets of sunflower.





Hypogynous Perigynous Perigynous Epigynous

Parts of a Flower

a) Calyx

- It is the outermost whorl of flower. It is made of **sepals**.

- Generally, sepals are green, leaf like and protect the flower in the bud stage.
- The calyx may be **gamosepalous** (sepals united) or **polysepalous** (sepals free).




b) Corolla

- It is the whorl inner to calyx. It is composed of **petals**.
- Petals have bright colour to attract insects for pollination.
- Corolla may be **gamopetalous** (petals united) or **polypetalous** (petals free).
- Shape and colour of corolla vary in plants. Corolla may be tubular, bell-shaped, funnel-shaped or wheel-shaped.

Aestivation: It is the mode of arrangement of sepals and petals in floral bud. It is many types:

- **Valvate:** Sepals or petals in a whorl just touch one another at the margin, without overlapping. E.g. *Calotropis*.
- **Twisted:** One margin of the appendage overlaps that of the next one and so on. E.g. China rose, lady's finger & cotton.
- **Imbricate:** Margins of sepals or petals overlap one another but not in any particular direction. E.g. *Cassia* & gulmohur.
- **Vexillary (papilionaceous):** In this, there are five petals; the largest (standard) overlaps the two lateral petals (wings) which in turn overlap the two smallest anterior petals (keel). E.g. pea & bean.



c) Androecium

- The male reproductive part composed of **stamens**.
- Each stamen represents the male reproductive organ. It consists of a **stalk (filament)** and an **anther**.
- Each anther is usually **bilobed**.
- Each lobe has 2 chambers called **pollen-sacs**.
- In pollen-sacs, **pollen grains** are produced.
- A sterile stamen is called **staminode**.
- When stamens are attached to petals, they are **epipetalous**. E.g. brinjal. When stamens are attached to perianth they are **epiphyllous**. E.g. lily.
- If the stamens are free, it is called **polyandrous**.

- If they are united, it is called **synandrous**. It is many types:
 - o **Monadelphous**: Stamens are united into one bunch or one bundle. E.g. China rose.
 - o **Diadelphous**: Stamens are united into two bundles. E.g. pea.
 - o **Polyadelphous**: Stamens are united into more than two bundles. E.g. citrus.
- There may be a variation in the length of filaments within a flower. E.g. *Salvia* and mustard.

d) Gynoecium (Pistil)

The female reproductive part made up of one or more **carpels**. A carpel has 3 parts:


- o **Stigma**: It is the receptive surface for pollen grains. It is usually at the tip of the **style**.
- o **Style**: An elongated tube that connects ovary to stigma.
- o **Ovary**: It is the enlarged basal part on which the style lies. Each ovary bears one or more **ovules** attached to a flattened, cushion-like **placenta**.

Polycarpellary pistils (pistil with many carpels) are 2 types:

- o **Apocarpous**: Carpels are free. E.g. lotus and rose.
- o **Syncarpous**: Carpels are fused. E.g. mustard and tomato.

Placentation: It is the arrangement of ovules on the placenta within the ovary. It is many types:

- o **Marginal**: Here, the placenta forms a ridge along the ventral suture of the ovary and the ovules are borne on this ridge forming two rows. E.g. pea.
- o **Axile**: Here, the placenta is axial and the ovules are attached to it in a multilocular ovary. E.g. China rose, tomato and lemon.
- o **Parietal**: Here, the ovules develop on the inner wall of the ovary or on peripheral part. Ovary is one-chambered but it becomes two-chambered due to the formation of the false septum. E.g. mustard and *Argemone*.
- o **Basal**: Here, placenta develops at the base of ovary and a single ovule is attached to it. E.g. sunflower, marigold.
- o **Free central**: Here, ovules are borne on central axis and septa are absent. E.g. *Dianthus* and *Primrose*.




After fertilisation, the ovules develop into seeds and the ovary matures into a fruit.

THE FRUIT

- It is a **ripened ovary** developed after fertilisation.
- It is a characteristic feature of the flowering plants.
- A fruit formed without fertilisation of the ovary is called **parthenocarpic** fruit.
- In mango & coconut, fruit is called a **drupe**. They are one seeded and develop from monocarpellary superior ovaries.
- A fruit consists of
 - o **Pericarp (fruit wall)**: It may be dry or fleshy. Thick and fleshy pericarp is differentiated into outer **epicarp**, middle **mesocarp** and inner **endocarp**.
 - o **Seeds**

- In mango, the pericarp is well differentiated into thin epicarp, fleshy edible mesocarp and stony hard endocarp.
- In coconut, the mesocarp is fibrous.




THE SEED

- It is the mature ovule developed after fertilisation.
- A seed is made up of a **seed coat** and an **embryo**.
- Embryo is made up of a **radicle**, an **embryonal axis** and **one** (e.g. wheat, maize) or **2 cotyledons** (e.g. gram & pea).

Structure of a Dicotyledonous Seed

- The outermost covering of a seed is the seed coat.
- Seed coat has 2 layers: outer **testa** and inner **tegmen**.
- On the seed coat, there is a scar called **hilum** through which the developing seeds are attached to the fruit.
- Above the hilum is a small pore called the **micropyle**.




Structure of dicotyledonous seed

- Within the seed coat is the **embryo**, consisting of an **embryonal axis** and **two cotyledons**.
- The cotyledons are often fleshy and full of reserve food materials. At the two ends of the embryonal axis, the **radicle** and **plumule** are present.
- In some seeds such as castor, the **endosperm** is formed due to double fertilisation. It is a food storing tissue.
- In plants such as bean, gram and pea, the seeds are **non-endospermous** (endosperm is not seen in mature seeds).

Structure of Monocotyledonous Seed

- Generally, monocot seeds are **endospermic** but some are non-endospermic (e.g. orchids).
- In cereals such as maize, the seed coat is membranous and generally fused with the fruit wall.
- The endosperm is bulky and stores food.


- The outer covering of endosperm separates the embryo by a protein layer called **aleurone layer**.
- The embryo is small and situated in a groove at one end of the endosperm. It consists of one large and shield shaped cotyledon known as **scutellum** and a short axis with a **plumule** and a **radicle**.
- The plumule is protected in a sheath called **coleoptile** and radicle is protected in a sheath called **coleorhiza**.




SEMI-TECHNICAL DESCRIPTION OF A TYPICAL FLOWERING PLANT

- The plant is described beginning with its habit, vegetative characters and then floral characters.
- Then a **floral diagram** and a **floral formula** are presented.
- Floral formula is represented by some symbols. They are
Br (bracteates) **K** (calyx) **C** (corolla)
P (perianth) **A** (androecium) **G** (Gynoecium)
G (superior ovary) $\overline{\text{G}}$ (inferior ovary)
♂ (male) ♀ (female) ♀ (bisexual)
⊕ (actinomorphic) % (zygomorphic)
- Fusion is indicated by enclosing the figure within bracket and adhesion by a line drawn above the symbols of the floral parts.

- A floral diagram gives information about the number of parts of a flower, their arrangement and relation.



Floral formula



Floral diagram of mustard plant (Family: *Brassicaceae*)

- Floral formula also shows cohesion and adhesion within parts of whorls and in between whorls.

SOME IMPORTANT FAMILIES

1. Fabaceae


This family was earlier called **Papilionoideae**, a subfamily of family **Leguminosae**. It is distributed all over the world.

Vegetative Characters:

- Trees, shrubs, herbs; root with root nodules.
- **Stem:** erect or climber.
- **Leaves:** alternate, pinnately compound or simple; leaf base, pulvinate; stipulate; venation reticulate.

Floral characters:

- **Inflorescence:** racemose.
- **Flower:** bisexual, zygomorphic.
- **Calyx:** sepals five, gamosepalous; valvate/imbricate aestivation.
- **Corolla:** petals five, polypetalous, papilionaceous, consisting of a posterior standard, two lateral wings, two anterior ones forming a keel (enclosing stamens and pistil), vexillary aestivation.
- **Androecium:** ten, diadelphous, anther dithecous.
- **Gynoecium:** ovary superior, mono carpellary, unilocular with many ovules, style single.
- **Fruit:** legume; seed: one to many, non-endospermic.
- **Floral Formula:** % ♀ $\text{K}_{(5)} \text{C}_{1+2+(2)} \text{A}_{(9)+1} \underline{\text{G}}_1$



Pisum sativum (pea) plant: (a) Flowering twig (b) Flower (c) Petals (d) Reproductive parts (e) L.S. carpel (f) Floral diagram

Economic importance:

- Pulses: E.g. gram, *arhar*, *sem*, *moong*, soyabean.
- Edible oil: E.g. soyabean, groundnut.
- Dye: E.g. Indigofera.
- Fibres: E.g. sun hemp.
- Fodder: E.g. *Sesbania*, *Trifolium*.
- Ornamentals: E.g. lupin, sweet pea.
- Medicine: E.g. *muliathi*.

2. Solanaceae (Potato family)


It is a large family. It is widely distributed in tropics, subtropics and even temperate zones.

Vegetative Characters:

- Plants mostly herbs, shrubs and small trees.
- **Stem:** herbaceous rarely woody, aerial; erect, cylindrical, branched, solid or hollow, hairy or glabrous, underground stem in potato (*Solanum tuberosum*).
- **Leaves:** alternate, simple, rarely pinnately compound, exstipulate; venation reticulate.

Floral Characters:

- **Inflorescence:** Solitary, axillary or cymose as in *Solanum*.
- **Flower:** bisexual, actinomorphic.
- **Calyx:** sepals five, united, persistent, valvate aestivation.
- **Corolla:** petals five, united; valvate aestivation.
- **Androecium:** stamens five, epipetalous.
- **Gynoecium:** bicarpellary obligately placed, syncarpous; ovary superior, bilocular, placenta swollen with many ovules, axile.
- **Fruits:** berry or capsule.
- **Seeds:** many, endospermous
- **Floral Formula:** $\oplus \overline{\text{G}} \text{K}_{(5)} \text{C}_{(5)} \text{A}_{(5)} \underline{\text{G}}_{(2)}$



Solanum nigrum (makoi) plant: (a) Flowering twig (b) Flower (c) L.S. of flower (d) Stamens (e) Carpel (f) Floral diagram

Economic Importance:

- Food: E.g. tomato, brinjal, potato
- Spice: E.g. chilli
- Medicine: E.g. belladonna, *ashwagandha*.
- Fumigatory: E.g. tobacco.
- Ornamentals: E.g. petunia.

3. Lilaceae (Lily family)


A characteristic representative of monocotyledonous plants. It is distributed worldwide.

Vegetative characters:

- Perennial herbs with underground bulbs/corms/ rhizomes.
- **Leaves** mostly basal, alternate, linear, exstipulate with parallel venation.

Floral characters:

- **Inflorescence:** solitary / cymose; often umbellate clusters.
- **Flower:** bisexual; actinomorphic.
- **Perianth** tepal six (3+3), often united into tube; valvate aestivation.
- **Androecium:** stamen six, (3+3).
- **Gynoecium:** tricarpeillary, syncarpous, ovary superior, trilobular with many ovules; axile placentation.
- **Fruit:** capsule, rarely berry.
- **Seed:** endospermous
- **Floral Formula:** $\oplus \varnothing P_{(3+3)} A_{3+3} \underline{G}_{(3)}$




Allium cepa (onion) plant: (a) Plant (b) Inflorescence (c) Flower (d) Floral diagram

Economic Importance:

- Ornamentals: E.g. tulip, *Gloriosa*
- Medicine: E.g. *Aloe*
- Vegetables: E.g. *Asparagus*
- Colchicines: E.g. *Colchicum autumnale*

MODEL QUESTIONS

1. Even though Ginger is seen under the soil, it is not a root, but a stem.
 - a. Give reason. b. Write its functions.
2. While examining a flower, Geetha noticed in its corolla a large outer standard petal, two small wing-like petals and two innermost petals united in to a keel. The aestivation was of vexillary type.
 - a. Identify the family of the plant which produced the above flower.
 - b. Write the floral formula of the family.
3. Identify the types of the arrangement of petals shown in the following diagrams.



4. Pick out the whorled arrangement of leaves from the group given below and write why it is said so? Neem, Nerium, Nepenthes
5. From the following terms relating a flower write its floral formula and family. Bisexual, sepals 5 united, petals 5 united, stamens 5 epipetalous, Carpels 2 superior.
6. Plants growing in swampy areas have special type of roots. Name the roots and their function?
7. From the following group of plants choose the best examples for root, stem and leaf modification. Hibiscus, Nepenthes, Rice, Carrot, Ginger, Calotropis

6. ANATOMY OF FLOWERING PLANTS

- Anatomy is the study of internal structure of plants and other organisms.


THE TISSUES

- A tissue is a group of cells having common origin and function.
- Based on the capability of cell division, plant tissues are 2 groups: **Meristematic** and **Permanent**.

MERISTEMATIC TISSUES (MERISTEMS)

These are the tissues where active cell division and growth occurs. Based on the position, meristems are 3 types:

- o **Apical meristems:** They occur at the tips of roots and shoots and produce primary tissues. It is 2 types:
 - **Root apical meristem:** It occupies the tip of a root.
 - **Shoot apical meristem:** It occupies the distant most region of the stem axis. Some cells 'left behind' from shoot apical meristem, constitute the **axillary bud**. They are present in the axils of leaves and can form a branch or a flower.



Apical meristem: (a) Root (b) Shoot

- o **Intercalary meristems:** They occur between mature tissues. They occur in grasses and regenerate parts removed by the grazing herbivores. Apical and intercalary meristems are **primary meristems** because they appear early in a plant life and contribute to the formation of primary plant body. During that, specific regions of the apical meristem produce **dermal tissues**, **ground tissues** and **vascular tissues**.
- o **Secondary (lateral) meristems:** The meristems that occur in mature regions of roots and shoots. They are cylindrical meristems. They are seen in gymnosperms and dicots. E.g. **Fascicular vascular cambium**, **interfascicular cambium** & **cork cambium**. These are responsible for producing the secondary tissues.

PERMANENT (MATURE) TISSUES

- The cells produced by primary and secondary meristems, become structurally and functionally specialized and lose the ability to divide. They are called **permanent (mature) cells** and constitute the **permanent tissues**.
- They are 2 types: Simple and Complex.


1. Simple Permanent Tissues

- The tissues having all cells similar in structure & function.
- 3 types: Parenchyma, Collenchyma and Sclerenchyma.

a. Parenchyma


- It forms the major component within organs.

- Cells are generally isodiametric. They may be spherical, oval, round, polygonal or elongated.
- Their walls are thin and made up of cellulose.
- Cells are closely packed or have small intercellular spaces.
- **Functions:** Photosynthesis, storage, secretion etc.




b. Collenchyma

- It occurs in layers below the epidermis in dicot plants.
- It is found as a homogeneous layer or in patches.
- It consists of cells with much thickened corners due to deposition of cellulose, hemicellulose and pectin.
- Intercellular spaces are absent.
- Cells are oval, spherical or polygonal and often contain chloroplasts.
- **Functions:** They provide mechanical support to the growing parts such as young stem and petiole of a leaf. The cells that contain chloroplasts assimilate food.



c. Sclerenchyma

- It consists of long, narrow cells with thick and lignified cell walls having a few or numerous pits.
- They are usually dead without protoplasts.
- Based on the form, structure, origin and development, sclerenchyma is 2 types: fibres & sclereids.
- o **Fibres:** These are thick-walled, elongated and pointed cells, generally occurring in groups.
- o **Sclereids:** These are spherical, oval or cylindrical, highly thickened dead cells with very narrow cavities (lumen). These are found in the fruit walls of nuts; pulp of fruits like guava, pear and sapota; seed coats of legumes and leaves of tea.
- **Function:** It provides mechanical support to organs.




2. Complex Permanent Tissues

- These are made of more than one type of cells and they work together as a unit.
- 2 types: Xylem and Phloem.

a. Xylem

- It functions as a conducting tissue for water and minerals from roots to the stem and leaves.
- It also provides mechanical strength to the plant parts.
- It is composed of 4 kinds of elements: **tracheids**, **vessels**, **xylem fibres** and **xylem parenchyma**.

- **Tracheids:** These are elongated tube like dead cells with thick and lignified walls and tapering ends. Protoplasm absent. The inner layers of cell walls have thickenings which vary in form. In flowering plants, tracheids & vessels are the main water transporting elements.



- **Vessel:** It is a long cylindrical tube-like structure made up of many cells (**vessel members**), each with lignified walls and a large central cavity. Protoplasm absent. Vessel members are interconnected through perforations in their common walls. The vessels are a characteristic feature of angiosperms. Gymnosperms lack vessels.

- **Xylem fibres:** They have highly thickened walls and obliterated central lumens. They are septate or aseptate.

- **Xylem parenchyma:** Living and thin-walled cells with cellulosic cell walls. They store food materials (starch or fat) and other substances like tannins. Radial conduction of water takes place by the ray parenchymatous cells.

- Primary xylem is 2 types:

- **Protoxylem:** The first formed primary xylem.
- **Metaxylem:** The later formed primary xylem.

- In stems, the protoxylem lies towards the centre (pith) and the metaxylem lies towards the periphery of the organ. This type of primary xylem is called **endarch**.

- In roots, the protoxylem lies towards periphery and metaxylem lies towards the centre. Such arrangement of primary xylem is called **exarch**.


b. Phloem (Bast)

It transports food materials from leaves to other parts.

In angiosperms, phloem is composed of **sieve tube elements, companion cells, phloem parenchyma & phloem fibres**.

Gymnosperms have albuminous cells and sieve cells. They lack sieve tubes and companion cells.

- **Sieve tube elements:** These are long, tube-like structures, arranged longitudinally and are associated with companion cells. Their end walls are perforated to form the **sieve plates**. A mature sieve element has a peripheral cytoplasm and a large vacuole but lacks a nucleus. The functions of sieve tubes are controlled by the nucleus of companion cells.



The first formed primary phloem (**protophloem**) consists of narrow sieve tubes. The later formed phloem (**metaphloem**) has bigger sieve tubes.

Function: Conduction of food materials from leaves.

- **Companion cells:** Specialized parenchymatous cells closely associated with sieve tube elements. Sieve tube elements & companion cells are connected by pit fields present between their common longitudinal walls.

Function: Maintain the pressure gradient in sieve tubes.

- **Phloem parenchyma:** It is made up of elongated, tapering cylindrical cells which have dense cytoplasm and nucleus. The cell wall is composed of cellulose and has pits through which plasmodesmatal connections exist between the cells. Phloem parenchyma is absent in most of the monocots.

Function: It stores food material and other substances like resins, latex and mucilage.

- **Phloem fibres (bast fibres):** These are made up of sclerenchymatous cells. Generally absent in primary phloem but are found in the secondary phloem. These are much elongated, unbranched and have pointed, needle like apices. Cell wall is quite thick. At maturity, these fibres lose their protoplasm and become dead. Phloem fibres of jute, flax and hemp are used commercially.

Function: Mechanical support & protection to soft tissues.

THE TISSUE SYSTEM

Based on structure and location, tissue systems are 3 types:

- **Epidermal tissue system**
- **Ground (fundamental) tissue system**
- **Vascular (conducting) tissue system**

1. Epidermal Tissue System


- It forms the outer-most covering of the whole plant body.
- It comprises **epidermal cells, stomata and epidermal appendages (trichomes & hairs)**.

Epidermis

- It is the outermost layer of the primary plant body.
- Epidermis is usually single layered.
- It is made up of elongated, compactly arranged parenchymatous cells with a small amount of cytoplasm lining the cell wall and a large vacuole.
- The outside of the epidermis is often covered with a waxy thick layer (**cuticle**). It prevents the loss of water. Cuticle is absent in roots.

Stomata

- These are structures present in the epidermis of leaves.
- Stomata regulate the transpiration and gaseous exchange.
- A stoma is made of two bean-shaped cells (**guard cells**).
- In grasses, the guard cells are dumbbell shaped.



Stomata with bean-shaped guard cells

Stomata with dumb-bell shaped guard cell

- The outer walls of guard cells (away from the stomatal pore) are thin and the inner walls (towards the stomatal pore) are highly thickened.
- The guard cells possess chloroplasts and regulate the opening and closing of stomata.
- Sometimes, a few epidermal cells near the guard cells become specialized in their shape and size. They are known as **subsidiary cells**.

- The stomatal aperture, guard cells and the surrounding subsidiary cells are together called **stomatal apparatus**.

Epidermal appendages

- The cells of epidermis bear many hairs.
- **Root hairs:** Unicellular elongations of the epidermal cells. They help to absorb water and minerals from the soil.
- **Trichomes:** They are the epidermal hairs on the stem. They are usually multicellular, branched or unbranched and soft or stiff. They may be secretory. Trichomes help to prevent water loss due to transpiration.

2. The Ground Tissue System

- All tissues except epidermis and vascular bundles constitute the **ground tissue**.
- It consists of **simple tissues** (parenchyma, collenchyma and sclerenchyma).
- Parenchymatous cells are present in cortex, pericycle, pith and medullary rays, in the primary stems and roots.
- In leaves, the ground tissue consists of thin-walled chloroplast containing cells and is called **mesophyll**.

3. The Vascular Tissue System


It consists of **complex tissues** (xylem and phloem).

Xylem and Phloem together constitute **vascular bundles**. Based on the presence or absence of **cambium**, vascular bundles are 2 types:

- **Open type:** In this, cambium is present between phloem and xylem. So vascular bundles can form secondary xylem and phloem tissues. E.g. **dicotyledonous** stems.
- **Closed type:** In this, cambium is absent. Hence, they do not form secondary tissues. E.g. **monocotyledons**.

Based on the arrangement of xylem and phloem, vascular bundles are 2 types:

- **Radial type:** Xylem and phloem are arranged in an alternate manner on different radii. Seen in roots.
- **Conjoint type:** Xylem and phloem are jointly situated at the same radius of vascular bundles. Seen in stems and leaves. Conjoint vascular bundles usually have phloem located only on the outer side of xylem.




ANATOMY OF DICOTYLEDONOUS & MONOCOTYLEDONOUS PLANTS

Dicotyledonous (Dicot) Root

Transverse section of the sunflower root shows the following tissue organization:


- **Epidermis (epiblema):** The outermost layer. Many cells of epiblema protrude in the form of unicellular root hairs.
- **Cortex:** It consists of several layers of thin-walled parenchyma cells with intercellular spaces.
- **Endodermis:** Innermost layer of the cortex. It comprises a single layer of barrel-shaped cells without intercellular spaces. The tangential as well as radial walls of the endodermal cells have a deposition of water impermeable, waxy material-suberin-in the form of **casparian strips**.
- **Stele:** All tissues on the inner side of the endodermis together constitute stele. They include
 - **Pericycle:** A few layers of thick-walled parenchymatous cells next to endodermis. Initiation of lateral roots and vascular cambium during the secondary growth takes place in these cells.
 - **Pith:** Innermost region of the stele. It is small or inconspicuous.
 - **Conjunctive tissue:** The parenchymatous cells which lie between the xylem and the phloem.
 - **Vascular bundles:** 2-4 xylem & phloem patches. Later, a cambium ring develops between the xylem & phloem.



T.S of Dicot root (Primary)


Monocotyledonous (Monocot) Root

- It has **epidermis, cortex, endodermis, pericycle, vascular bundles and pith**.
- There are usually more than six (**polyarch**) xylem bundles.
- Pith is large and well developed.
- Monocot roots do not undergo any secondary growth.



T.S of Monocot root

Dicotyledonous (Dicot) Stem




T.S. of Dicot stem

- **Epidermis:** Outermost protective layer. Covered with a thin layer of cuticle, it may bear trichomes & few stomata.
- **Cortex:** Multiple layers of the cells arranged in between epidermis and pericycle. It consists of 3 sub-zones:

- **Hypodermis:** Outer zone. It consists of a few layers of collenchymatous cells just below the epidermis. It provides mechanical strength to the young stem.
- **Cortical layers:** Below hypodermis. They consist of rounded thin walled parenchymatous cells with conspicuous intercellular spaces.
- **Endodermis:** Innermost layer. The cells are rich in starch grains. So the layer is also called as the **starch sheath**. **Pericycle** is present on the inner side of the endodermis and above the phloem in the form of semi-lunar patches of sclerenchyma.
- **Stele:** Consists of **pericycle, vascular bundles, medullary rays & pith**.
 - **Medullary rays:** These are few layers of radially placed parenchymatous cells in between vascular bundles.
 - **Vascular bundles:** Large in number. They are arranged in a ring. Ring arrangement is a characteristic of dicot stem. Each vascular bundle is conjoint, open, and with endarch protoxylem.
 - **Pith:** Central portion of the stem. It has many rounded, parenchymatous cells with large intercellular spaces.

Monocotyledonous (Monocot) Stem



- It has a sclerenchymatous **hypodermis**, many scattered **vascular bundles**, each surrounded by a sclerenchymatous **bundle sheath**, and a large, conspicuous parenchymatous **ground tissue**.
- **Vascular bundles** are **conjoint & closed**. Peripheral vascular bundles are smaller than centrally located ones.
- The phloem parenchyma is absent, and water-containing cavities are present within the vascular bundles.


Dicotyledonous (Dorsiventral) Leaf

The vertical section of a dicot leaf through lamina shows 3 main parts: **Epidermis, mesophyll & vascular system**.

- **Epidermis:** It covers both the upper surface (**adaxial epidermis**) and lower surface (**abaxial epidermis**) of the leaf. It has a conspicuous cuticle. Abaxial epidermis generally bears more stomata than the

adaxial epidermis. The latter may even lack stomata.

- **Mesophyll:** The tissue between the upper and the lower epidermis. It is made up of parenchyma. They contain chloroplasts




for photosynthesis. It has 2 types of cells:

- **Palisade parenchyma:** It is adaxially placed. Made up of elongated cells arranged vertically and parallel to each other.
- **Spongy parenchyma:** The oval or round and loosely arranged. It is situated below the palisade cells and extends to the lower epidermis. There are numerous large spaces and air cavities between these cells.
- **Vascular system:** It includes vascular bundles. They can be seen in the veins and midrib.

Size of vascular bundles is dependent on the size of the veins. The veins vary in thickness in the reticulate venation of dicot leaves. Vascular bundles are surrounded by a layer of thick walled **bundle sheath cells**.

Monocotyledonous (Isobilateral) Leaf

The anatomy of monocot leaf is like that of the dicot leaf in many ways. It shows following differences:



- Stomata are present on both surfaces of the epidermis.
- Mesophyll is not differentiated into palisade and spongy parenchyma.
- In grasses, certain adaxial epidermal cells along the veins modify themselves into large, empty, colourless cells. These are called **bulliform cells**. When the bulliform cells have absorbed water and are turgid, the leaf surface is exposed. When they are flaccid due to water stress, they make the leaves curl inwards to minimise water loss.
- Parallel venation is reflected in the near similar sizes of vascular bundles (except in main veins).

SECONDARY GROWTH

- The growth of the roots and stems in length with the help of apical meristem is called the **primary growth**.
- **Secondary growth** is the increase in girth of dicot plants.
- Tissues involved in secondary growth are the two **lateral meristems: Vascular cambium & cork cambium**.

Vascular Cambium

- It is the meristematic layer responsible for cutting off vascular tissues (xylem and phloem).


- In the young stem, it is present in patches as a single layer between xylem & phloem. Later it forms a complete ring.

Formation of cambial ring

- In dicot stems, cells of cambium present between primary xylem & primary phloem is **intrafascicular cambium**.
- Cells of medullary cells, adjoining this intrafascicular cambium become meristematic and form **interfascicular cambium**. Thus, a continuous ring of cambium is formed.

Activity of the cambial ring

- The cambial ring becomes active and cut off new cells, both towards the inner and outer sides. The cells cut off towards pith, mature into **secondary xylem**. The cells cut off towards periphery mature into **secondary phloem**.
- Cambium is more active on the inner side than on the outer. As a result, more secondary xylem is produced than secondary phloem and soon forms a compact mass.
- Primary and secondary phloems get gradually crushed due to the continued formation and accumulation of secondary xylem. However, primary xylem remains intact, in or around the centre. At some places, cambium forms a narrow band of parenchyma, which passes through the secondary xylem and the secondary phloem in the radial directions. These are the **secondary medullary rays**.



Secondary growth in a dicot stem – stages in transverse views

Spring wood and autumn wood

- Many physiological & environmental factors control the activity of cambium.
- In spring season, cambium is very active and produces many xylary elements having vessels with wider cavities. This wood is called **spring wood (early wood)**. It is lighter in colour and has a lower density.
- In winter, cambium is less active and forms fewer xylary elements having narrow vessels. This wood is called **autumn wood (late wood)**. It is darker and has higher density.
- These two kinds of woods that appear as alternate concentric rings constitute an **annual ring**. This is used to estimate the age of tree (Dendrochronology).

Heartwood and sapwood

- **Heartwood:** It is the hard, dead, dark brown-coloured, highly lignified and non-functional central part of the secondary xylem of old trees. The dark colour is due to deposition of organic compounds (tannins, resins, oils, gums, aromatic substances, essential oils etc). These substances make it hard, durable and resistant to the attacks of microorganisms and insects.

Function: It gives mechanical support to stem.

- **Sapwood:** It is the peripheral region of secondary xylem. It is living and lighter in colour. It is involved in the conduction of water and minerals from root to leaf.

Cork Cambium


- As the stem continues to increase in girth due to the activity

of vascular cambium, the outer cortical & epidermis layers get broken. It is to be replaced to provide new protective cell layers. Hence another meristematic tissue called **cork cambium (phellogen)** develops, usually in the cortex.


- Phellogen is a couple of layers thick. It is made of narrow, thin-walled and nearly rectangular cells.
- Phellogen cuts off cells on both sides. The outer cells differentiate into **cork (phellem)** while the inner cells differentiate into **secondary cortex (phelloderm)**. Cells of secondary cortex are parenchymatous.
- The cork is impervious to water due to suberin deposition in the cell wall.
- Phellogen, phellem, and phelloderm are collectively known as **periderm**. Due to activity of cork cambium, pressure builds up on the remaining layers peripheral to phellogen and ultimately these layers die and slough off.
- **Bark** is a non-technical term that refers to all tissues (such as periderm & secondary phloem) exterior to the vascular cambium. It is 2 types:

- o **Early (soft) bark:** It is formed early in the season.
- o **Late (hard) bark:** It is formed towards end of season.

- **Lenticels:** At certain regions, phellogen cuts off closely arranged parenchymatous cells on outer side. These cells rupture epidermis, forming a lens shaped openings called **lenticels**. They occur in most woody trees. **Function:** Lenticels permit gas exchange of between the outer atmosphere and the internal tissue of the stem.



Secondary Growth in Roots



Different stages of the secondary growth in a typical dicot root




- In dicot root, vascular cambium is completely secondary in origin. It originates from the tissue located just below the phloem bundles (a portion of pericycle) above the protoxylem forming a complete and continuous wavy ring. It later becomes circular. Further events are similar to those of a dicotyledon stem.
- Secondary growth also occurs in stems and roots of gymnosperms. However, secondary growth does not occur in monocotyledons.

MODEL QUESTIONS

1. Find out the odd one
 - a. Parenchyma, Xylem, Collenchyma, Sclerenchyma
 - b. Sieve tubes, companion cells, tracheids, bast fibre
2. Analyze the table and arrange them in an appropriate order

A	B	C
Monocot Stem	Bulliform cells	Bean shaped guard cells
Isobilateral leaf	Endarch Xylem	Secondary thickening
Dorsiventral leaf	Closed Vascular bundle	Dumble shaped guard cells
Dicot stem	Palisade and spongy parenchyma	Proto xylem and lacunae

3. Give reasons.
 - a. Conduction of the food through the sieve tube is under the control of companion cell.
 - b. Annual rings are not found in coconut tree.
4. Identify the type of vascular bundle.



5. Vascular bundles of a plant are conjoint, collateral and open.
 - a. In which part of a plant this kind of vascular bundles are seen?
 - b. Is it possible for this part of plant to undergo secondary thickening? Give reasons.
6. Some anatomical characters are given in brackets. Arrange them in three columns under root, stem and leaves. (Conjoint open bundles, mesophyll cells, endarch xylem, radial bundles, bulliform cells on epidermis, casparian thickenings in endodermis, exarch xylem)

7. EVOLUTION

Evolution is an orderly change from one form to another.

Evolutionary Biology is the study of evolutionary history of life forms.

ORIGIN OF LIFE

- **Big Bang Theory** states that universe originated about 20 billion years ago by a singular huge explosion.
- The earth was formed about **4.5 billion years** ago.
- There was no atmosphere on early earth. Water vapour, CH₄, CO₂ & NH₃ released from molten mass covered the surface.
- The UV rays from the sun broke up water into H₂ and O₂.
- Oxygen combined with NH₃ & CH₄ to form water, CO₂ etc.
- The ozone layer was formed. As it cooled, the water vapour fell as rain to form oceans.
- Life appeared almost **four billion years** ago.

THEORIES OF ORIGIN OF LIFE

- 1. Theory of spontaneous generation (Abiogenesis):** It states that, life came out of decaying and rotting matter like straw, mud etc.

Louis Pasteur disproved this theory. He demonstrated that life comes only from pre-existing life.


He showed that life did not come from killed yeast in a closed pre-sterilized flask. But in an opened flask, life (microbes) appeared.

- 2. Biogenesis:** Proposed by **Francisco Redi, Spallanzani & Louis Pasteur**. It states that, life originates from pre-existing life. But it does not explain origin of first life.
- 3. Cosmic theory (Theory of Panspermia):** It states that, the units of life (spores) were transferred to different planets including earth.
- 4. Theory of special creation:** It states that, living things were created by some supernatural power (God).

- 5. Theory of chemical evolution:** Proposed by **Oparin & Haldane**. It states that, the first form of life was originated from non-living inorganic & organic molecules such as CH₄, NH₃, H₂O, sugars, proteins, nucleic acids etc. i.e. *“Abiogenesis first, but biogenesis ever since”*.

Urey-Miller experiment

- **Harold Urey & Stanley Miller** experimentally proved theory of chemical evolution. They created a condition like that of primitive earth (i.e. high temperature,



- volcanic storms, reducing atmosphere with CH₄, NH₃, H₂O, H₂ etc).
- They made electric discharge in a closed flask containing CH₄, NH₃, H₂ and water vapour at 800°C. As a result, some amino acids are formed.
- In similar experiments, others observed formation of sugars, nitrogen bases, pigment and fats.

First **non-cellular forms** of life originated 3 billion years ago. They were **self-replicating metabolic capsule** containing RNA, proteins, Polysaccharides etc.

EVIDENCES FOR EVOLUTION

1. Paleontological evidences

Paleontology is the study of fossils.

Fossils are remnants of life forms found in rocks (earth crust).

They are written documents of evolution.

Significance of fossils:

- To study **phylogeny** (evolutionary history or race history). E.g. Horse evolution.
- To study the **connecting link** between two groups of organisms. E.g. *Archaeopteryx*.
- To study about **extinct animals**. E.g. Dinosaurs.
- To study about **geological period** by analysing fossils in different **sedimentary rock layers**. The study showed that life forms varied over time and certain life forms are restricted to certain geological time spans.

2. Morphological & Anatomical evidences

Comparative anatomy and morphology shows that different forms of animals have some common structural features. This can be explained as follows:

a. Homologous organs

- **Homologous organs** are the organs having fundamentally

similar structure and origin but different functions. This phenomenon is called **Homology**.

- E.g. Human hand, Whale's flippers, Bat's wing & Cheetah's foot. These forelimbs have different functions but similar anatomical structures such as bones (e.g. humerus, radius, ulna, carpals, metacarpals & phalanges).
- Homology is also seen in heart, brain etc.
- **Homology in plants:** E.g. Thorns of *Bougainvillea* and tendrils of *Cucurbita*.
- The origin of homologous organs is due to **Divergent evolution**. It is the evolution by which **related species** become **less similar** to survive and adapt in different environmental condition.
- Homology indicates common ancestry.

b. Analogous organs

These are the organs having similar function but different structure & origin. This phenomenon is called **Analogy**. E.g.

- **Wings of insects** (formed of a thin flap of chitin) and **wings of birds** (modified forelimbs).
- **Eyes of Octopus** (retina from skin) and **mammals** (retina from embryonic brain).


- **Flipper of Penguins and Dolphins.**
- **Sweet potato** (modified root) & **Potato** (modified stem).
- **Trachea of insects** (from ectoderm) and **lungs of vertebrates** (from endoderm).

Origin of analogous organs is due to **Convergent evolution**. It is the evolution by which **unrelated species** become more **similar** to survive and adapt in similar environmental condition.

3. Adaptive radiation (Biogeographical evidences)

Adaptive radiation (evolution by adaptation) is the evolution of different species from an ancestor in a geographical area starting from a point. It is a type of divergent evolution. E.g.

- Darwin's finches in Galapagos Islands.



- Australian marsupials (Marsupial radiation).

- Placental mammals in Australia.

When more than one adaptive radiation is appeared in an isolated geographical area, it results in **convergent evolution**. E.g. Australian Marsupials and Placental mammals.

Placental mammals	Australian Marsupials
Mole	Marsupial mole
Ant eater	Numbat (Ant eater)
Mouse	Marsupial mouse
Lemur	Spotted cuscus
Flying squirrel	Flying phalanger
Bobcat	Tasmanian tiger cat
Wolf	Tasmanian wolf

4. Biochemical evidences

- Organisms show similarities in proteins, genes, other biomolecules & metabolism. It indicates common ancestry.

5. Embryological evidences

- Proposed by **Ernst Haeckel**.
- He observed that all vertebrate embryos have some common features that are absent in adult.
- E.g. all vertebrate embryos (including human) develop vestigial gill slits just behind the head. But it is functional only in fish and not found in other adult vertebrates.
- However, **Karl Ernst von Baer** rejected this proposal. He noted that embryos never pass through the adult stages of other animals.

6. Evidences for evolution by natural selection

Natural selection is the process in which organisms with better favourable & heritable variation are survived and reproduced.

Some evidences are given below:

- **Industrial melanism:** In England, before industrialization (1850s), there were more white-winged moths (*Biston betularia*) on trees than dark winged or melanised moths (*Biston carbonaria*). After industrialization (1920), more dark-winged moths and less white winged moths were developed.

Reason:

Before industrialization: There was white lichens covered the trees. In that background, white winged moths survived but dark winged moths were picked out by predators.

After industrialization: The tree trunks became dark due to industrial smoke and soot. No growth of lichens. So white winged moths did not survive because the predators identified them easily. Dark winged moth survived because of suitable dark background.

- Development of resistant varieties in organisms against **herbicides, pesticides, antibiotics or drugs** etc.

These are the examples for natural selection by **anthropogenic action** (evolution due to human activities).

THEORIES OF BIOLOGICAL EVOLUTION

Lamarckism (Theory of Inheritance of Acquired characters)

It is proposed by Lamarck. It states that evolution of life forms occurred by the inheritance of acquired characters.

Acquired characters are developed by use & disuse of organs.

- **Evolution by use of organs:** E.g. Long neck of giraffe is due to continuous elongation to forage leaves on trees. This acquired character was inherited to succeeding generations.
- **Evolution by disuse:** E.g. Disappearance of limbs in snakes.

This theory was eliminated out because it is proved that the characters are inherited only through genes.

Darwinism (Theory of Natural selection)

- Proposed by **Charles Darwin**.
- It was based on observations during a sea voyage in a sail ship called **H.M.S. Beagle**.
- **Alfred Wallace** (a naturalist worked in **Malay Archipelago**) had also come to similar conclusions.
- Work of **Thomas Malthus** on populations influenced Darwin.

Darwinism is based on 2 key concepts:

- **Branching descent:** It explains that all organisms are modified descendants of previous life forms.
- **Natural selection:** Consider a bacterial colony **A** growing on a given medium. If the medium composition is changed, only a part of the population can survive under new condition. This variant population (**B**) outgrows the others and appears as new species, i.e. **B is better than A under new condition**. Thus, nature selects for fitness.

Natural selection is based on the following facts:

- **Heritable minor variations:** It is either beneficial or harmful to the organisms.
- **Overproduction:** Population size grows exponentially due to maximum reproduction (E.g. bacterial population).
- **Limited natural resources:** Resources are not increased in accordance with the population size.
- **Struggle for existence:** It is the competition among organisms for resources so that population size is limited.
- **Survival of the fittest:** In struggle for existence, organisms with beneficial variations can utilize resources better. Hence, they survive and reproduce. This is called

Survival of the fittest. It leads to a change in population characteristics and new forms appear.

Darwin ignored about origin of variation and mechanism of evolution or speciation.

MECHANISM OF EVOLUTION

- **Hugo de Vries** proposed **Mutation Theory** of evolution.
- He conducted experiments on *Oenothera lamarckiana*

(evening primrose) and believed that evolution takes place through mutation and not by minor variation.

- **Darwinian variation** is minor, slow and directional. It results in **gradual evolution**.
- **Mutational variation** is sudden, random & directionless. Here, speciation is by **saltation** (single step, large mutation).
- Mutation is the origin of variation for evolution.

HARDY-WEINBERG PRINCIPLE

- It states that *allele frequencies in a population are stable and is constant from generation to generation in the absence of disturbing factors.*

- The **gene pool** (total genes and their alleles in a population) remains a constant. This is called **genetic equilibrium (Hardy-Weinberg equilibrium)**.

- Sum total of all the allelic frequencies = 1

- E.g. Consider, in a diploid, **p** & **q** are the frequencies of alleles **A** & **a** respectively.

$$\text{Frequency of AA} = p^2$$

$$\text{Frequency of aa} = q^2$$

$$\text{Frequency of Aa} = 2pq$$

$$\text{Hence } p^2 + 2pq + q^2 = 1 \text{ [binomial expansion of } (p+q)^2]$$

Change of frequency of alleles in a population disturbs Hardy-Weinberg equilibrium. This change is due to evolution.

Factors affecting Hardy-Weinberg equilibrium


- Gene migration:** Gene flow from one population to another. Here gene frequencies change in both populations. Gene flow occurs if migration happens multiple times.
- Genetic drift:** The gene flow by chance causing change in frequency. Sometimes, the change in frequency is so different in the new sample of population that they become a different species. The original drifted population becomes founders and the effect is called **founder effect**.

c. Mutation: It results in formation of new phenotypes. Over few generations, this leads to speciation.

d. Genetic recombination: Reshuffling of gene combinations during crossing over resulting in genetic variation.

e. Natural selection: It is 3 types.

- **Stabilizing selection:** Here, more individuals acquire mean character value and variation is reduced.
- **Directional selection:** Individuals of one extreme (value other than mean character value) are more favoured.
- **Disruptive selection:** Individuals of both extremes (peripheral character value at both ends of the distribution curve) are more favoured.



A BRIEF ACCOUNT OF EVOLUTION

The geological time scale includes 4 eras: **Proterozoic, Palaeozoic, Mesozoic & Cenozoic.**

1. Proterozoic era: 2500 - 541 million yrs ago (mya)

- **2000 mya:** First cellular forms of life appeared.
- Some of the cells had the ability to release O_2 as the light reaction in photosynthesis.
- Single celled organisms became multicellular organisms.

2. Palaeozoic era (540 - 252 mya)

- It has 6 periods: **Cambrian** (540 - 490 mya), **Ordovician** (490 - 443 mya), **Silurian** (425 mya), **Devonian** (405 mya), **Carboniferous** (360 mya) & **Permian** (285 mya).
- **500 mya:** Invertebrates were formed.
- **450 mya:** First land organisms (plants) appeared.
- **400 mya:** Arthropods invaded the land.
- **350 mya:** Jawless fishes were evolved.

Lobefins (stout & strong finned fishes) could move on land and go back to water. They evolved to first amphibians (ancestors of modern day frogs & salamanders).

In 1938, a lobe-fin called **coelacanth** fish was caught in South Africa which was thought to be extinct.

- **320 mya:** Sea weeds and few plants were existed.
- Amphibians evolved to reptiles. They lay thick-shelled eggs (do not dry up in sun).
- **Giant ferns (Pteridophytes)** were present but they all fell to form coal deposits slowly.

3. Mesozoic era (252 - 66 mya)

- Age of reptiles and gymnosperms.
- It has 3 periods: **Triassic (230 mya)**, **Jurassic (208 mya)** & **Cretaceous (144 mya)**.
- **200 mya:** Some of the land reptiles went back into water to evolve into fish-like reptiles (E.g. *Ichthyosaurs*).
- The land reptiles were **dinosaurs** (*Tyrannosaurus rex*, *Triceratops*, *Stegosaurus*, *Brachiosaurus* etc.)
- **T. rex** was the largest dinosaur (20 feet in height, huge fearsome dagger-like teeth).
- Toothed birds were emerged.

4. Cenozoic era (66 - 0 mya)

- Age of Mammals & Angiosperms.
- It has 2 periods: **Tertiary (66 mya) & Quaternary (2 mya)** - Age of man).
- **65 mya:** Dinosaurs suddenly disappeared. Some say climatic changes killed them. Some say most of them evolved into birds.
- First mammals were shrew-like. Their fossils are small sized.
- In South America, there were mammals resembling horse, hippopotamus, bear, rabbit etc. Due to continental drift, when South America joined North America, these animals were overridden by North American fauna.
- Due to continental drift, Australian marsupials survived because of lack of competition from any other mammals.

ORIGIN AND EVOLUTION OF MAN

- **15 mya: *Dryopithecus* & *Ramapithecus*.**
Hairy. Walked like gorillas & chimpanzee.
Dryopithecus: ape-like.
Ramapithecus: man-like.
- **3-4 mya: Man-like primates** walked up right in eastern

Africa. Height up to 4 feet. This belief is based on fossils of man-like bones found in Ethiopia & Tanzania.

- **2 mya: *Australopithecus*.** Lived in East African grass lands. Hunted with stone weapons. Ate fruits.
***Homo habilis*:** First human-like being (hominid).
Brain capacity: 650-800 cc. Did not eat meat.
- **1.5 mya: *Homo erectus* (Java man).** Large brain (900 cc).
Ate meat.
- **1 lakh - 40,000 yrs ago: *Homo neanderthalensis* (Neanderthal man).**
Brain capacity: 1400 cc. Lived in East & Central Asia. Used hides to protect their body. Buried their dead.
- **75,000 - 10,000 yrs ago (ice age): *Homo sapiens* (Modern man).**
Pre-historic cave art developed about 18,000 years ago. E.g. Cave paintings at Bhimbetka rock shelter in Raisen district of Madhya Pradesh.
Agriculture & settlements: 10,000 years ago.

Sequence of Human evolution:

Dryopithecus → *Ramapithecus* → *Australopithecus* → *Homo habilis* → *H. erectus* → *H. neanderthalensis* → *H. sapiens*

Visit: www.bankofbiology.com

MODEL QUESTIONS

1. Match the following:

A	B	C
Charles Darwin	Chemical evolution	Use and disuse of organs
Lamarck	Natural selection	Abiogenic origin of life in ocean
Hugo de Vries	Biogenesis	<i>Oenothera lamarckiana</i>
Louis Pasteur	Inheritance of acquired characters	Survival of the fittest
Oparin & Haldane	Mutation	Disproved theory of spontaneous generation

2. Analyze the relationship between first two words and fill the fourth place.
 - a. Homology: Divergent evolution Analogy:
 - b. *Pisum sativum*: Mendel *Oenothera lamarckiana*:
3. Classify the following points into two categories. Give suitable titles.
Random & directionless, Minor variation, Gradual evolution, Slow & directional,
Large variation, Speciation by saltation
4. A bacterial infection was effectively controlled by using a specific antibiotic for a long time. But now- a- days this antibiotic is not found to be so effective. Give a scientific explanation for this phenomenon based on evolution.
5. Hardy- Weinberg Principle has a great contribution in population genetics.
 - a. State Hardy- Weinberg Principle.
 - b. What are the factors affecting genetic equilibrium?
 - c. What is meant by Founder effect?
6. Select the correct order
 - a. Paleozoic era → Proterozoic era → Mesozoic era → Coenozoic era
 - b. Mesozoic era → Proterozoic era → Coenozoic era → Paleozoic era
 - c. Proterozoic era → Paleozoic era → Mesozoic era → Coenozoic era
 - d. Coenozoic era → Paleozoic era → Mesozoic era → Proterozoic era
7. Prepare a flowchart showing the evolution of man.

7. STRUCTURAL ORGANISATION IN ANIMALS

ANIMAL TISSUES

A group of cells having *same origin, structure and function* are called the **tissues**. Animal tissues are 4 types:

(i) **Epithelial** (ii) **Connective** (iii) **Muscular** (iv) **Neural**

I. EPITHELIAL TISSUE (EPITHELIUM)

- It has a **free surface** that faces **body fluid** or **outside environment**.
- **Covers** or **lines** body or body parts.
- Compactly packed cells with **little intercellular matrix**.
- Epithelial tissues are 2 types: **Simple** and **Compound**.

1. Simple epithelium

It is composed of a **single layer of cells**. It lines body cavities, ducts and tubes. Based on **structural modification of cells**, simple epithelium is 3 types:

o Squamous epithelium:


- Thin layer of flattened cells with irregular boundaries.
- Found in the walls of blood vessels and lung alveoli.
- **Functions:** Form a diffusion boundary.

o Cuboidal (cubical) epithelium:

- Composed of cube-like cells.
- Found in ducts of glands and tubular parts of nephrons.
- **Functions:** Secretion and absorption.
- The epithelium of **proximal convoluted tubule (PCT)** of nephron in the kidney has **microvilli**.

o Columnar epithelium:

- Composed of tall and slender cells.
- Their nuclei are located at the base.
- Free surface may have microvilli.
- Found in the lining of stomach and intestine.
- **Functions:** Secretion and absorption.




Modification of columnar or cuboidal cells

• Ciliated epithelium:

- Cells bearing **cilia** on their free surface.
- Present in the inner surface of hollow organs like bronchioles and fallopian tubes.
- **Functions:** To move particles or mucus in a specific direction over the epithelium.

• Glandular epithelium: For secretion. They are 2 types:


- **Unicellular:** Consists of isolated glandular cells. E.g. Goblet cells of the alimentary canal.
- **Multicellular:** Contains cluster of cells. E.g. salivary glands.



- Based on mode of pouring of secretions, glands are 2 types:
- **Exocrine glands:** Here, secretions are released through **ducts (tubes)**. Exocrine glands secrete **mucus, saliva, earwax, oil, milk, digestive enzymes** etc.
 - **Endocrine glands:** Ductless. They produce **hormones**.

2. Compound epithelium

- Made up of more than one layer (**multi-layered**) of cells.
- Limited role in secretion & absorption.
- They cover dry surface of **skin**, moist surface of **buccal cavity, pharynx, inner lining of ducts of salivary glands and pancreatic ducts**.
- **Function:** Protect against chemical & mechanical stresses.



Cell junctions: The junctions that provide structural and functional links between adjacent cells. They are found in epithelium and other tissues. They are 3 types:

- Tight junctions:** Help to stop substances from leaking across a tissue.
- Adhering junctions:** Perform cementing to keep neighbouring cells together.
- Gap junctions:** Facilitate communication b/w adjoining cells by connecting the cytoplasm for rapid transfer of ions, small molecules and sometimes big molecules.

II. CONNECTIVE TISSUE


- It links and supports other tissues/organs.
- They are most abundant in complex animals.
- All connective tissues except blood have **fibroblast** cells. They secrete structural fibrous proteins called **collagen & elastin**. They give strength, elasticity & flexibility to tissue.
- The cells also secrete modified polysaccharides (**matrix**), which accumulate between cells and fibres.
- Types of connective tissues: **Loose, Dense & Specialised**.

1. Loose Connective Tissues

In this, cells (**fibroblasts, macrophages, mast cells** etc.) and fibres are loosely arranged in a semi-fluid matrix. It is 2 types: Areolar & Adipose.


o Areolar tissue:

- Present beneath the skin.
- It serves as a support framework for epithelium.



o **Adipose tissue:**

- Seen mainly under skin.
- Its cells (**adipocytes**) store fats.
- Excess nutrients which are converted into fats are stored in this tissue.




2. Dense Connective Tissues

In this, fibres and fibroblasts are compactly packed. 2 types:

o **Dense regular connective tissues:** *RTL*


- Show regular pattern of fibres.
- Collagen fibres are present in rows between many parallel bundles of fibres.
- E.g. tendons & ligaments.
 - Tendons:** Attach muscles to bones.
 - Ligaments:** Attach one bone to another.



o **Dense irregular connective tissues:**

Irregular pattern of fibres.


- Fibroblasts & fibres (mostly collagen) are oriented differently.
- This tissue is present in skin.



3. Specialized Connective Tissues


o **Cartilage:**

- o In this, intercellular material (matrix) is solid and pliable (due to **chondroitin salts**) and resists compression.
- o Cartilage cells (**chondrocytes**) are enclosed in small cavities within the matrix secreted by them.
- o Most of the cartilages in vertebrate embryos are replaced by bones in adults.
- o Cartilage is present in the **tip of nose, outer ear, joints in the vertebral column, limbs and hands in adults.**



o **Bone:**

- o It has hard and non-pliable matrix rich in **calcium salts** and **collagen fibres** which give bone its strength.
- o Bone cells (**osteocytes**) are seen in spaces called **lacunae**.
- o **Functions:**
 - It provides structural frame to the body.
 - Support and protect softer tissues and organs.




- Limb bones serve weight-bearing functions.
- Take part in locomotion and movements.
- Blood cells are produced in bone marrow.

o **Blood:**

- o A fluid connective tissue containing **plasma, red blood cells (RBC), white blood cells (WBC)** and **platelets**.
- o Helps in the circulation of various substances.

III. MUSCLE TISSUE

- The tissues made of many **muscle fibres (muscle cells)**.
- Muscle fibres are composed of numerous fine **myofibrils**.
- Muscle fibres can contract (shorten) and relax (lengthen).
- Muscles take part in locomotion and movements.
- Muscles are 3 types: **skeletal, smooth** and **cardiac**.



1. Skeletal (striated or voluntary) muscle

- They are attached to bones. E.g. **Biceps**.
- Striations are present in muscle fibres.
- Muscle fibres are bundled together in a parallel fashion.
- A sheath of tough connective tissue encloses several bundles of muscle fibres.

2. Smooth (non-striated or visceral) muscle

- **Involuntary** and **fusiform** (Fibres taper at both ends).
- No striations.
- Cell junctions hold them together and they are bundled together in a connective tissue sheath.
- They are seen in the wall of internal organs such as the **blood vessels, stomach and intestine**.

3. Cardiac muscle

- **Involuntary** muscle seen only in the **heart**.
- Cell junctions fuse the plasma membranes of cardiac muscle cells and make them stick together.
- Communication (gap) junctions (**intercalated discs**) at some fusion points allow cells to contract as a unit, i.e., when a cell receives signal to contract, other cells also contract.

IV. NEURAL TISSUE

- o Made up of neurons (unit of neural system).
- o Responsible for **control** and **co-ordination** of the body.
- o **Neurons** are excitable cells. They carry impulses.
- o Neurons are protected and supported by **neuroglial cells**.
- o Neuroglia make up more than half the volume of neural tissue.

ORGAN AND ORGAN SYSTEM

- o **Cells → tissues → organs → organ systems.**
- o This organization is essential for better coordinated activities of cells.
- o An organ is made of one or more type of tissues. E.g. Heart has epithelial, connective, muscular & neural tissues.

MORPHOLOGY & ANATOMY OF COCKROACH


- **Morphology:** Study of external features of organisms.
- **Anatomy:** Study of morphology of internal organs.
- **Cockroach (*Periplaneta americana*)** are nocturnal, omnivores and live in damp places.
- **Colour:** Brown or black. Bright yellow, red & green coloured cockroaches are also seen in tropical regions.
- **Size:** ¼ inches to 3 inches (0.6-7.6 cm).

Systematic position

Phylum : Arthropoda
Class : Insecta
Genus : *Periplaneta*
Species : *americana*


MORPHOLOGY OF COCKROACH

- The adults are about **34-53 mm** long.
- Body is covered by a hard brown **chitinous exoskeleton**.
- In each segment, exoskeleton has hardened plates called **sclerites (dorsal tergites & ventral sternites)**. They are joined to each other by a thin and flexible **articular membrane (arthrodial membrane)**.
- The body has 3 regions – **head, thorax** and **abdomen**.



Head

- Triangular head is formed by 6 fused segments.
- It shows great mobility in all directions due to flexible neck.
- Head bears a pair of thread-like **antennae**, a pair of **compound eyes** and **biting & chewing type** mouth parts.



- **Mouthparts:** a **labrum** (upper lip), 2 **mandibles**, 2 **maxillae**, **hypopharynx** (tongue) & a **labium** (lower lip).

Thorax

- It has 3 parts: **prothorax, mesothorax & metathorax**.
- The head is connected to thorax by a **neck** (short extension of the prothorax).
- Each thoracic segment bears a pair of walking legs.
- 2 pairs of wings: **Forewings (2)** and **Hind wings (2)**.

- **Forewings (mesothoracic) or tegmina:** Opaque, dark and leathery and cover the hind wings when at rest.
- **Hind wings (metathoracic):** Transparent, membranous and are used in flight.

Abdomen

- It consists of **10 segments**.
- **In females**, 7th (boat shaped), 8th & 9th sterna form a **brood (genital) pouch**. It contains female **gonopore, spermathecal pores & collateral glands**.
- **In males**, **genital pouch** lies at the hind end of abdomen bounded dorsally by 9th & 10th terga and ventrally by the 9th sternum. It contains **dorsal anus, ventral male genital pore (gonopore)** and **gonapophysis**.
- In both sexes, 10th segment bears a pair of jointed **anal cerci**. Males bear a pair of short, threadlike **anal styles**.

Differences between male & female cockroaches

Male	Female
i. Larger size	Smaller
ii. Wings extend beyond the tip of the abdomen.	Wings do not extend beyond the tip of abdomen.
iii. Narrow abdomen	Broad abdomen
iv. Anal styles present	Absent
v. Brood pouch absent	Present

ANATOMY OF COCKROACH

Digestive system

Alimentary canal has 3 parts: **foregut, mid gut & hindgut**.

- **Foregut:** It is lined by **cuticle**. It includes **Mouth → pharynx → oesophagus → crop (to store food) → gizzard (proventriculus)**.


Gizzard helps in grinding the food. It

has an outer layer of thick circular muscles and thick inner cuticle forming **6 chitinous plates (teeth)**.


- **Mid gut (Mesenteron):** It is **not lined** by cuticle. 6-8 tubules (**hepatic or gastric caecae**) are seen at the junction of foregut & mid gut. They secrete digestive juice. At the junction of mid gut & hindgut, there are **100-150** yellow coloured thin filamentous **Malpighian tubules**.
- **Hindgut:** It is broader than mid gut and lined internally by **cuticle**. Hindgut includes **ileum, colon & rectum**. Rectum opens out through anus.

Circulatory system

- Blood vascular system: **open type**.
- Blood vessels are poorly developed and open into space (**haemocoel**).



- Visceral organs located in the haemocoel are bathed in blood (**haemolymph**).
- **Haemolymph**= **colourless plasma** + **haemocytes**.
- **Heart** consists of elongated **muscular tube** lying along **mid dorsal line** of **thorax** and **abdomen**.
- It has funnel-shaped chambers with **ostia** on either side.
- Blood from sinuses enter heart through ostia and is pumped anteriorly to sinuses again.



Respiratory system

- It consists of a network of **trachea** that open through **10 pairs** of small holes called **spiracles** present on the lateral side of the body.
- The thin branches of **tracheal tubes** are called **tracheoles**. They carry oxygen from the air to all parts.
- The opening of the spiracles is regulated by **sphincters**.
- Gas exchange takes place at the tracheoles by **diffusion**.

Excretory system

- **Uricotelic**. Excretory organ is **Malpighian tubules**.
- Each tubule is lined by glandular and ciliated cells. They absorb nitrogenous wastes and convert them into **uric acid** which is excreted out through the hindgut.
- **Fat body, nephrocytes & urecose glands** also help in excretion.

Nervous system

- It consists of segmentally arranged **ganglia** joined by paired longitudinal connectives on the ventral side.
- 3 ganglia lie in the thorax and 6 in the abdomen.
- The head holds only a bit of nervous system. Remaining part is situated along the ventral part of the body. So, if the head of cockroach is cut off, it will still live for one week.
- The **supra-oesophageal ganglion** (brain) supplies nerves to antennae and compound eyes.
- **Sense organs: Antennae, eyes, maxillary palps, labial palps, anal cerci etc.**
- Sensory receptors of antennae monitor the environment.
- Each compound eye consists of about **2000 hexagonal ommatidia**. Using these, a cockroach can receive several images of an object. This is called **mosaic vision**. It has more sensitivity but less resolution, being common during night (hence called **nocturnal vision**).

Reproductive system


Cockroaches are **dioecious**.

Male reproductive system:

It consists of a pair of **testes**, **seminal vesicles**, **accessory glands & external genitalia**.


- **Testes**: Lie laterally in the **4th -6th abdominal segments**.

Each **testis** → a thin **vas deferens** → **seminal vesicle** → **ejaculatory duct** → **male gonopore**.



- **Seminal vesicles**: To **store sperms**. Sperms are glued together to form bundles called **spermatophores**. They are discharged during copulation.
- **Accessory glands** include a **mushroom gland** (in **6th-7th abdominal segments**) and **phallic gland**. Their secretions nourish the sperms.
- **External genitalia (male gonapophysis or phallomeres)**: Chitinous asymmetrical structures, surrounding the **male gonopore**.

Female reproductive system:



- It consists of 2 large **ovaries**, **oviducts**, **spermatheca**, **genital chamber**, **Collateral glands** etc.
- **Ovaries** lie laterally in the **2nd - 6th abdominal segments**. Each ovary is formed of **8 ovarian tubules (ovarioles)**, containing a chain of developing ova.
- **Oviducts** of each ovary unite into a single **median oviduct (vagina)** which opens into the **genital chamber**.
- A pair of **spermatheca** is present in the **6th segment** which opens into the **genital chamber**.
- Sperms are transferred through spermatophores. Their fertilised eggs are encased in **oothecae**.
- Ootheca is dark reddish to blackish brown capsule, 8 mm long. Females lay **9-10 oothecae**, each contain **14-16 eggs**.
- Development of *P. americana* is **paurometabolous**, (development through **nymphal stage**).
- **Nymphs** look like adults. They moult **13 times** to reach the adult form. The next to last nymphal stage has **wing pads**. Only adult cockroaches have wings.

ECONOMIC IMPORTANCE OF COCKROACH


They are pests because they destroy food and contaminate it with their smelly excreta. They also transmit bacterial diseases like **cholera**, **typhoid**, **tuberculosis** etc.

MODEL QUESTIONS


- Find the relationship between first two words and suggest a suitable word in the fourth position.
 - Squamous epithelium: lining of blood vessels Cuboidal epithelium:
 - Cartilage: chondrocytes Bone:
 - Frog: Kidney Cockroach:
- Odd man out. Justify your answer.
 - Areolar tissue, blood, neuron, tendon
 - Maxilla, Mandible, Labrum, Antennae
 - Mushroom gland, Ommatidium, Ootheca, Oviducts
 - Exocrine, Endocrine, Salivary gland, Ligament
- Identify the following tissues and write down their location.
 - Tissue which store fat
 - Tissue that conducts impulses
 - Tissue that connects muscle and bone
 - Tissue that serves as supporting frame of body
- Identify the following tissue and label A, B, C & D.



- Two types of muscle tissues are diagrammatically represented.



- Identify muscle tissues A and B. Mention their location.
 - Find out any two differences between them.
- Prepare a flowchart using the following terms.
Crop, Pharynx, Ileum, Mouth, Oesophagus, Mesenteron, Anus, Gizzard, Colon
 - Observe the diagram given below:



- Identify the figure.
- Label the parts P, Q, R and S.

- Analyse the following statements. Find out the wrong statements and rewrite them correctly.
 - In cockroach, hepatic or gastric caecae act as the digestive glands.
 - Mushroom gland and phallic gland are concerned with female reproductive system of cockroach.
 - Antennae, eyes, maxillary palps, labial palps etc. are the sensory organs of cockroach.
 - Anal cerci are the structures present only male cockroach and used to identify the sex of cockroach.


MORPHOLOGY & ANATOMY OF EARTHWORM

Systematic position

Phylum : Annelida
Class : Oligochaeta
Genus : *Pheretima*
Species : *posthuma*

- Earthworm is a reddish-brown terrestrial invertebrate that inhabits the upper layer of moist soil.
- During day time, they live in burrows made by boring and swallowing the soil.
- Common Indian earthworms: *Pheretima* and *Lumbricus*.

MORPHOLOGY OF EARTHWORM



- Earthworms have long segmented cylindrical body.
- Number of **segments (metameres)**: about **100-120**.
- **Dorsal surface** has a **dark median mid dorsal line (dorsal blood vessel)** along the longitudinal axis of the body.
- First segment (**peristomium or buccal segment**) bears the mouth. A lobe called **prostomium** covers the mouth.
- Prostomium is sensory in function and is used to force open cracks in the soil into which the earthworm may crawl.
- In a mature worm, segments **14-16** are covered by a dark band of glandular tissue called **clitellum**.
- Body has 3 regions: **preclitellar, clitellar & postclitellar**.
- **4 pairs of spermathecal apertures** are found on ventro-lateral sides of **intersegmental grooves (5th-9th segments)**.
- A single **female genital pore** is present in the mid-ventral line of **14th segment**.
- A pair of **male genital pores** is present on the ventro-lateral sides of the **18th segment**.
- Many minute **nephridiopores** open on the body surface.
- All segments except the first, last and clitellum bear S-shaped **setae**, embedded in the epidermal pits. Setae can be extended or retracted. Their function is locomotion.

ANATOMY OF EARTHWORM


Body wall

It is composed of

- Outermost thin **non-cellular cuticle**.


- **Epidermis**: Made up of a single layer of columnar epithelial cells which contain secretory gland cells.
- **Two muscle layers (circular and longitudinal)**.
- An innermost **coelomic epithelium**.

Digestive system



- The straight **alimentary canal** extends from first to last segment of the body. It has
- **Mouth** → **buccal cavity (1-3 segments)** → **muscular pharynx (4th segment)** → **oesophagus (5-7 segments)** → **muscular gizzard (8-9 segments)** → **stomach (9-14 segments)** → **Intestine (15th segment to last)** → **anus**.
- Gizzard helps to grind soil particles, decaying leaves, etc.
- **Calciferous glands**, present in the stomach, neutralise the **humic acid** present in humus.
- A pair of short and conical **intestinal caecae** project from the intestine on the **26th segment**.
- The intestinal part between **26-35 segments** has an internal median fold of dorsal wall called **typhlosole**. It increases area of absorption.
- The organic rich soil is digested in the digestive tract by digestive enzymes. Digested nutrients are absorbed through intestinal membranes. Their faecal deposits are known as **worm castings**.

Circulatory system (blood vascular system)




- Circulatory system is **closed type** (blood flows through heart and blood vessels).
- Consists of **blood vessels, capillaries** and **heart**.
- Contractions keep blood circulating in one direction.
- **Blood glands** are present on the **4th, 5th & 6th segments**. They produce **phagocytic blood cells** and **haemoglobin** which is dissolved in blood plasma.

Respiratory system

- No specialized system.
- Gas exchange occurs through moist body surface into the blood stream.

Excretory system



Excretory organs are segmentally arranged tubules called **nephridia**. They are 3 types:

- Septal nephridia:** Found on both sides of intersegmental septa (segment 15 to last) that open into intestine.
- Integumentary nephridia:** Attached to lining of body wall (segment 3 to last). They open on body surface.
- Pharyngeal nephridia:** Present in the **4th, 5th & 6th segments** in the form of paired tufts.

Funnel-shaped part of nephridium collects excess fluid from coelom. The funnel connects with a tubular part of nephridium which delivers the wastes into digestive tube.

Nervous system

- Includes segmentally arranged **ganglia** on the ventral paired and fused **nerve cord**.
- The nerve cord in the anterior region (**3rd & 4th segments**) divides and encircles the pharynx and joins the **cerebral ganglia** dorsally to form a **nerve ring**.
- The nerve ring with cerebral ganglia represents the brain.
- **Sensory system:** Includes
 - Light and touch sensitive **receptor cells**. No eyes.
 - **Chemoreceptors (taste receptors):** React to chemical stimuli.

Reproductive system


Earthworm is **hermaphrodite**.

Male reproductive organs:

- **Testes:** 2 pairs. Enclosed in **testis sacs** in **10th & 11th segments**. The sperms from testes shed into testis sacs. From where, they enter **seminal vesicles** for maturation. Mature sperms move back into testis sacs and enter **spermiducal funnels** which are connected to **vasa deferentia (spermatic ducts)**. The vasa deferentia run up to **18th segment** where they join the **prostatic duct**.

The **common prostate** and **spermatic duct** open to the exterior by a pair of **male genital pores** on the ventro-lateral side of the **18th segment**.

- **Accessory glands:** 2 pairs. Found in the **17th & 19th segments** (one pair in each segment).



Female reproductive organs:

- **Spermathecae:** 4 pairs. Located in **6th-9th segments** (one pair in each segment). They receive and store spermatozoa during copulation.
- **Paired ovaries:** Attached at the inter-segmental septum of the **12th and 13th segments**.
- **Ovarian funnels:** Present beneath the ovaries which continue into **oviduct**, join together and open on ventral side as single median female genital pore on **14th segment**.
- During mating, two worms exchange sperms each other. They mate juxtaposing opposite gonadal openings exchanging **spermatophores** (packets of sperms).
- Mature sperm, ova and nutritive fluid are deposited in **cocoons** produced by gland cells of clitellum. Cocoon with fertilized ova slips off and deposit in the soil. After 3 weeks, cocoon produces 2 to 20 baby worms (no larva).

ECONOMIC IMPORTANCE

- Earthworms are known as **'friends of farmers'** because they make burrows in the soil and make it porous which helps in respiration and penetration of the plant roots. This process of increasing fertility of soil is called **vermicomposting**.
- They are used as **bait** in game fishing.

MORPHOLOGY & ANATOMY OF FROG

Systematic position


Phylum : Chordata
Class : Amphibia
Genus : *Rana*
Species : *tigrina*

- *Rana tigrina* is the most common species in India.
- They are **poikilotherms (cold blooded)**.
- They can change colour to hide them from their enemies (**camouflage**). This protective coloration is called **mimicry**.
- During summer and winter, they undergo **aestivation (summer sleep)** and **hibernation (winter sleep)** respectively to protect them from extreme heat and cold.

MORPHOLOGY OF FROG

- Body is divisible into **head & trunk**. Neck and tail absent.
- Skin is moist, smooth and slippery due to the mucus.
- Colour of dorsal side is olive green with dark irregular spots and ventral side is pale yellow.
- The frog never drinks water but absorb it through the skin.
- A **mouth**, paired **nostrils** and bulged **eyes** (covered by nictitating membrane) are present.
- On either side of eyes have a membranous **tympanum (ear)**.
- The **forelimbs (4 digits)** and **hind limbs (5 digits)** help in **swimming, walking, leaping and burrowing**. The hind limbs are larger and muscular than fore limbs.
- Feet have webbed digits that help in swimming.
- Frogs exhibit **sexual dimorphism**. Male frogs have sound producing **vocal sac** and also a **copulatory (nuptial) pad** on the first digit of fore limbs which are absent in female frogs.

ANATOMY OF FROG



Digestive system

- Consists of **alimentary canal** and **digestive glands**.
- The alimentary canal is short because frogs are carnivores and hence the length of intestine is reduced.
- **Mouth** → **buccal cavity** → **pharynx** → **oesophagus** → **stomach** → **intestine** → **rectum** → **cloaca**.
- Liver secretes **bile** that is stored in **gall bladder**. **Pancreas** produces **pancreatic juice** containing digestive enzymes.
- Food is captured by the bilobed tongue.

- **Digestion:** Gastric juice and HCl secreted from gastric wall digest the food. Partially digested food (**chyme**) is passed from stomach to the **duodenum**.

Duodenum receives **bile** and **pancreatic juices** through a **common bile duct**.

Bile emulsifies fat. Pancreatic juice digests carbohydrates and proteins. Digestion completes in the intestine.

- Finger-like **villi** and **microvilli** in intestine absorb digested food. The undigested solid waste moves into the **rectum** and passes out through **cloaca**.

Respiratory system

- **Skin** acts as aquatic respiratory organ (**cutaneous respiration**). Dissolved oxygen in the water is exchanged through the skin by diffusion. During aestivation and hibernation respiration takes place through skin.
- On land, the **buccal cavity, skin** and **lungs (pulmonary respiration)** act as the respiratory organs.
- The lungs are a pair of elongated, pink coloured sac-like structures present in the thorax. Air enters through the nostrils into the buccal cavity and then to lungs.

Circulatory system

- **Closed type.** Includes **Blood vascular system (heart, blood vessels & blood)** and **lymphatic system (lymph, lymph channels & lymph nodes)**.
- Heart is **3-chambered**, (**two atria** and **one ventricle**) and is covered by a membrane called **pericardium**.
- A triangular structure called **sinus venosus** joins the right atrium. It receives blood through major veins (**vena cava**).
- The ventricle opens into a saclike **conus arteriosus** on the ventral side of the heart.
- The blood pumped from the muscular heart is carried to all parts of the body by the **arteries (arterial system)**.
- The **veins** collect blood from different parts of body to the heart and form the **venous system**.
- **Hepatic portal system** (venous connection between liver and intestine) and **renal portal system** (between kidney and lower parts of the body) are present in frogs.
- Blood contains **plasma** and **cells (RBC, WBC & platelets)**. RBCs are nucleated and contain haemoglobin.
- Blood transports nutrients, gases and water to tissues.

Excretory system

- Includes **kidneys (2)**, **ureters (2)**, **cloaca & urinary bladder**.
- **Kidneys** are dark red and bean-shaped. Found posteriorly in the body cavity on both sides of vertebral column. Each kidney is formed of **uriniferous tubules (nephrons)**.
- 2 ureters emerge from the kidneys. In male frogs, the ureters act as **urinogenital duct** which opens into cloaca. In females, ureters & oviduct open separately in cloaca.
- The thin-walled **urinary bladder** is present ventral to the rectum which also opens in the cloaca.
- The frog is a **ureotelic** animal (excretes urea). Nitrogenous wastes are carried by blood into the kidney where it is separated and excreted.

Control and co-ordination

Endocrine system


- The **endocrine glands** secrete **hormones**.
- Endocrine glands: **pituitary, thyroid, parathyroid, thymus, pineal body, pancreatic islets, adrenals & gonads.**

Nervous system


It includes

- **Central nervous system (brain & spinal cord),**
- **Peripheral nervous system (cranial & spinal nerves)**
- **Autonomic nervous system (sympathetic & parasympathetic).**
- There are 10 pairs of **cranial nerves** arising from brain.
- Brain is enclosed in a bony **brain box (cranium).**
- The brain is divided into
 - **Fore-brain:** Includes **olfactory lobes, paired cerebral hemispheres** and **unpaired diencephalon.**
 - **Mid-brain:** Includes a pair of **optic lobes.**
 - **Hind-brain:** Includes **cerebellum & medulla oblongata.**
- **Medulla oblongata** passes out through the **foramen magnum** and continues into **spinal cord**, which is enclosed in the vertebral column.
- Sense organs include organs of
 - **Sensory papillae:** For touch
 - **Taste buds:** For taste
 - **Nasal epithelium:** For smell
 - **Simple eyes:** For vision. Paired and situated in orbit
 - **Tympanum with internal ears:** For hearing and balancing (equilibrium).

Reproductive system



- Male reproductive organs consist of a pair of yellowish ovoid **testes**, which are found adhered to the upper part of kidneys by a double fold of peritoneum (**mesorchium**).
- **Vasa efferentia** (10-12 in number) arise from testes. They enter the kidneys on their side and open into **Bidder's canal**. It communicates with urinogenital duct that comes out of the kidneys and opens into cloaca.
- The **cloaca** is a small, median chamber that is used to pass **faecal matter, urine and sperms** to the exterior.



- The female reproductive organs include a pair of **ovaries**. The ovaries are situated near kidneys and there is no functional connection with kidneys.
- A pair of oviduct arising from the ovaries opens into the cloaca separately.
- A mature female can lay **2500 to 3000 ova** at a time.
- Fertilisation is external and takes place in water.
- **Development** involves a **larval stage** called **tadpole**.
- Tadpole undergoes metamorphosis to form the adult.

ECONOMIC IMPORTANCE

- Frogs are beneficial for mankind because they eat insects and protect the crop.
- Maintain ecological balance by serving as an important link of food chain and food web in the ecosystem.
- In some countries the muscular legs of frog are used as food by man.

8. CELL: THE UNIT OF LIFE

- A cell is the fundamental, structural and functional unit of all living organisms.
- **Robert Hooke:** Discovered cell.
- **Anton Von Leeuwenhoek:** First observed and described a live cell.
- The invention of the **compound & electron microscopes** revealed all the structural details of the cell.

CELL THEORY

- **Matthias Schleiden (1838)** observed that all plants are composed of different kinds of cells.
- **Theodore Schwann (1839)** found that cells have a thin outer layer (plasma membrane). He also found that plant cells have cell wall. He proposed a hypothesis that animals and plants are composed of cells and products of cells.
- Schleiden & Schwann formulated the **cell theory**.
- **Rudolf Virchow (1855)** first explained that cells divide and new cells are formed from pre-existing cells (*Omnis cellula-e cellula*). He modified the cell theory.
- Cell theory states that:

- (i) All living organisms are composed of cells and products of cells.
- (ii) All cells arise from pre-existing cells.

AN OVERVIEW OF CELL

- All cells contain
 - o **Cytoplasm:** A semi-fluid matrix where cellular activities and chemical reactions occur. This keeps the cell in 'living state'.
 - o **Ribosomes:** Non-membrane bound organelles seen in cytoplasm, chloroplasts, mitochondria & on rough ER.
- Cells differ in size, shape and activities.
 - o **Smallest cells:** Mycoplasmas (0.3 μm in length).
 - o **Largest isolated single cell:** Egg of ostrich.
 - o **Longest cells:** E.g. Nerve cell.
 - o Size of bacteria: 3 to 5 μm (Typical: 1 to 2 μm).
 - o Human RBCs are about 7.0 μm in diameter.
- Based on the functions, shape of cells may be disc-like, polygonal, columnar, cuboid, thread like, or irregular.
- Cells are 2 types: **Prokaryotic cells & Eukaryotic cells.**

PROKARYOTIC CELLS

- They have no membrane bound nucleus and organelles.
- They include **bacteria, blue-green algae, mycoplasma & PPLO (Pleuro Pneumonia Like Organisms)**.
- They are generally smaller and multiply more rapidly than the eukaryotic cells.
- They vary in shape & size. E.g. Bacteria have 4 basic shapes: **Bacillus, Coccus, Vibrio and Spirillum.**

- o In distribution of chromosomes to daughter cells.
- o In respiration and secretion processes.
- o To increase the surface area of the plasma membrane and enzymatic content.
- **Chromatophores** are pigment-containing membranous infoldings in some prokaryotes (e.g. cyanobacteria).

Cell organelles in prokaryotic cells

1. Cell Envelope

- It is a chemically complex protective covering.
- It is made of 3 tightly bound layers.
 - o **Glycocalyx:** Outer layer. Its composition and thickness vary in different bacteria. It may be a **slime layer** (loose sheath) or **capsule** (thick & tough).
 - o **Cell wall:** Middle layer. Seen in all prokaryotes except mycoplasma. It gives shape to the cell and provides a structural support to prevent the bacterium from bursting or collapsing.
 - o **Plasma membrane:** Inner layer. It is semi-permeable in nature and interacts with the outside. This is structurally similar to that of the eukaryotes.
- Based on the types of the cell envelopes and response to Gram staining (developed by Gram), bacteria are 2 types:
 - o **Gram positive:** They take up and retain the gram stain.
 - o **Gram negative:** They do not retain the gram stain.

2. Mesosomes & Chromatophores (Membranous structures)

- **Mesosome** is formed by the infoldings of plasma membrane. It includes **vesicles, tubules & lamellae**.
- **Functions:** Mesosomes help
 - o In cell wall formation.
 - o In DNA (chromosome) replication.

3. Nucleoid

- It is formed of non-membranous (naked) circular **genomic DNA** (single chromosome/ Genetic material) & protein.
- Many bacteria have small circular DNA (**plasmid**) outside the genomic DNA. It gives some unique phenotypic characters (e.g. resistance to antibiotics) to bacteria.

4. Flagella

- These are thin filamentous extensions from the cell wall of motile bacteria. Their number and arrangement are varied in different bacteria.
- Bacterial flagellum has 3 parts – **filament, hook and basal body**. The filament is the longest portion and extends from the cell surface to the outside.

5. Pili and Fimbriae

- These are surface structures that have no role in motility.
- **Pili** (sing. Pilus) are elongated tubular structures made of a special protein (**pilin**).
- **Fimbriae** are small bristle like fibres sprouting out of the cell. In some bacteria, they help to attach the bacteria to rocks in streams and to the host tissues.

6. Ribosomes

- They are associated with plasma membrane of prokaryotes.
- They are about 15 nm by 20 nm in size.
- They are made of 2 subunits - **50S & 30S (Svedberg's unit)**. They together form **70S** prokaryotic ribosomes.

(S= sedimentation coefficient; a measure of density & size).

- **Function:** Ribosomes are the site of **translation** (protein synthesis). Several ribosomes may attach to a single mRNA to form a chain called **polyribosomes (polysome)**. Ribosomes translate the mRNA into proteins.


7. Inclusion Bodies

- These are non-membranous, stored reserve material seen freely in the cytoplasm of prokaryotic cells.
- E.g. phosphate granules, cyanophycean granules and glycogen granules, gas vacuoles etc.
- **Gas vacuoles** are found in blue green and purple and green photosynthetic bacteria.

EUKARYOTIC CELLS

- They have well organized **membrane bound nucleus and organelles**.
- Presence of membranes gives clear compartmentalization of cytoplasm.
- Their genetic material is organized into chromosomes.
- They have complex locomotory & cytoskeletal structures.

Plant cell and Animal cell




Cell organelles in eukaryotic cells

1. Cell Membrane

- Chemical studies on human RBCs show that cell membrane is composed of a **lipid bilayer, protein & carbohydrate**.
- Lipids (mainly **phosphoglycerides**) have outer **polar head** and the inner **hydrophobic tails**. So the non-polar tail of saturated hydrocarbons is protected from the aqueous environment.
- Ratio of protein and lipid varies in different cells. E.g. In human RBC, membrane has 52% protein and 40% lipids.
- Based on the ease of extraction, membrane proteins are 2 types:
 - o **Integral proteins:** Partially or totally buried in membrane.

- o **Peripheral proteins:** Lie on the surface of membrane.
- **Fluid mosaic model of cell membrane:** Proposed by **Singer & Nicolson (1972)**. According to this, the quasi-fluid nature of lipid enables lateral movement of proteins within the overall bilayer. This ability to move within the membrane is measured as its fluidity.



Functions:

- o Transport of the molecules. The membrane is selectively permeable to some molecules present on either side of it.
- o Due to the fluid nature, the plasma membrane can help in cell growth, formation of intercellular junctions, secretion, endocytosis, cell division etc.

Types of Transport

1. Passive transport: It is the movement of molecules across the membrane along the concentration gradient (i.e., from higher concentration to the lower) without the expenditure of energy. It is 2 types:

- a. Simple diffusion:** It is the movement of neutral solutes across the membrane.
- b. Osmosis:** It is the movement of water by diffusion across the membrane.

Polar molecules cannot pass through the non-polar lipid bilayer. So they require membrane carrier protein for transport.

2. Active transport: It is the movement of molecules across the membrane against the concentration gradient (i.e. from lower to the higher concentration) with the expenditure of energy (ATP is utilized). E.g. Na⁺/K⁺ pump.

2. Cell Wall

- It is a non-living rigid structure found outer to the plasma membrane of fungi and plants.
- Cell wall of Algae is made of cellulose, galactans, mannans and minerals like CaCO₃. In other plants, it consists of cellulose, hemicellulose, pectins and proteins.
- Cell wall of a young plant cell (**primary wall**) is capable of growth. It gradually diminishes as the cell matures and

the **secondary wall** is formed on the inner side (towards membrane).

- The **middle lamella** is a layer containing calcium pectate which glues the neighbouring cells together. Cell wall and middle lamellae may be traversed by **plasmodesmata**. It connects the cytoplasm of neighbouring cells.

Functions:

- It gives shape to the cell.
- It protects the cell from mechanical damage & infection.
- It helps in cell-to-cell interaction.
- It acts as barrier to undesirable macromolecules.

3. Endomembrane System

- It is a group of membranous organelles having coordinated functions.
- They include endoplasmic reticulum (ER), Golgi complex, lysosomes and vacuoles.

Endoplasmic Reticulum (ER)

- These are a network of tiny tubular structures scattered in the cytoplasm.


ER divides the intracellular space into 2 compartments: **luminal** (inside ER) & **extra luminal** (cytoplasm).

Endoplasmic reticulum is 2 types:

a. **Rough endoplasmic reticulum (RER):**


Bear ribosomes on their surface. RER is frequently observed in the cells actively involved in protein synthesis and secretion. They extend to the outer membrane of the nucleus.

b. **Smooth endoplasmic reticulum (SER):** Ribosomes are absent. SER is the major site for synthesis of lipid. In animal cells lipid-like steroidal hormones are synthesized in SER.



Golgi apparatus

- Densely stained reticular structures near the nucleus.
- First observed by **Camillo Golgi** (1898).
- They consist of flat, disc-shaped sacs (**cisternae**) of 0.5– 1.0 μm diameter. These are stacked parallelly.
- Cisternae are concentrically arranged with convex **cis** (forming) face and concave **trans** (maturing) face. **Cis** & **trans** faces are totally different, but interconnected.



Function of Golgi apparatus:

- o Secretes materials to intra-cellular targets or outside the cell.

Materials to be packaged as vesicles from the ER fuse with the **cis** face and move towards the **trans** face. This is why Golgi apparatus remains in close association with the endoplasmic reticulum.

- o Proteins synthesized by ribosomes on the ER are modified in the cisternae of Golgi apparatus before they are released from its **trans** face.
- o Formation of glycoproteins and glycolipids.

Lysosomes


- These are membrane bound vesicular structures formed by the process of packaging in the Golgi apparatus.
- Lysosomal vesicles contain almost all types of hydrolytic enzymes (hydrolases– lipases, proteases, carbohydrases). They are active at acidic pH. They digest carbohydrates, proteins, lipids and nucleic acids.

Vacuoles

- These are the membrane-bound space found in the cytoplasm. It contains water, sap, excretory product and other materials not useful for the cell.
- Vacuole is bound by a single membrane called **tonoplast**.
- In plant cells, the vacuoles can occupy up to 90% of the volume of the cell.
- In plants, the tonoplast facilitates the transport of ions and other materials against concentration gradients into the vacuole. Hence their concentration is higher in the vacuole than in the cytoplasm.
- In *Amoeba*, the **contractile vacuole** helps for excretion.
- In many cells (e.g. protists), **food vacuoles** are formed by engulfing the food particles.

4. Mitochondria

- Mitochondria are clearly visible only when stained.
- Number, shape and size of mitochondria per cell are variable depending on the physiological activity.
- It is sausage-shaped or cylindrical having a diameter of 0.2-1.0 μm (average 0.5 μm) and length 1.0-4.1 μm.



- A mitochondrion is a double membrane-bound structure with the outer membrane and the inner membrane. It divides lumen into 2 aqueous compartments, i.e., the outer compartment and the inner compartment (**matrix**).
- Inner membrane forms a number of infoldings (**cristae**) towards the matrix. They increase the surface area.
- The two membranes have their own specific enzymes associated with the mitochondrial function.
- Matrix possesses a circular DNA, a few RNA molecules, ribosomes (70S) and components for protein synthesis.
- The mitochondria divide by fission.

- **Function:** Mitochondria are the sites of aerobic respiration. They produce energy in the form of ATP. So they are called '**power houses**' of the cell.


5. Plastids

- Plastids are found in all plant cells and in euglenoides.
- Large sized. Easily observable under the microscope.
- They contain some pigments.
- Based on the type of pigments, plastids are 3 types:
 - Chloroplasts:** Contain **chlorophyll** and **carotenoid** pigments. They trap light energy for photosynthesis.
 - Chromoplasts:** Contain fat soluble **carotenoid** pigments like carotene, xanthophylls etc. This gives a yellow, orange or red colour.
 - Leucoplasts:** These are colourless plastids of varied shapes and sizes with stored nutrients. They include:
 - **Amyloplasts:** Store starch. E.g. potato.
 - **Elaioplasts:** Store oils and fats.
 - **Aleuroplasts:** Store proteins.

Chloroplasts:

- These are double membrane bound organelles mainly found in the **mesophyll** cells of the leaves.


- These are lens-shaped, oval, spherical, discoid or ribbon-like organelles.



- Length: 5-10 μm . Width: 2-4 μm .
- Their number varies from 1 (e.g. *Chlamydomonas*) to 20-40 per cell in the mesophyll.
- Inner membrane of chloroplast is less permeable.
- The space limited by the inner membrane of the chloroplast is called **stroma**. It contains many organized flattened membranous sacs called **thylakoids**.
- Membrane of thylakoids encloses a space called **lumen**.
- **Chlorophyll** pigments are present in the thylakoids.
- Thylakoids are arranged in stacks called **grana** or the intergranal thylakoids.
- There are flat membranous tubules called the **stroma lamellae** connecting the thylakoids of the different grana.
- The stroma contains small, double-stranded circular DNA molecules, ribosomes and enzymes for the synthesis of carbohydrates and proteins.
- The ribosomes of the chloroplasts are smaller (70S) than the cytoplasmic ribosomes (80S).

6. Ribosomes

- They are non-membranous granular structures composed of **ribonucleic acid (RNA) & proteins**.
- It is first observed by **George Palade** (1953).
- Eukaryotic ribosome has 2 subunits- **60S** (large subunit) and **40S** (small subunit). They together form **80S**.



7. Cytoskeleton


- It is a network of filamentous proteinaceous structures

present in the cytoplasm.

- It provides mechanical support, motility, maintenance of the shape of the cell etc.

8. Cilia and Flagella

- They are hair-like outgrowths of the cell membrane.
- **Cilia:** Small structures which work like oars. Causes the movement of the cell or surrounding fluid.
- **Flagella:** Longer. Responsible for cell movement. Flagella of prokaryotes and eukaryotes are structurally different.




- Cilium and flagellum are covered with plasma membrane. Their core (**axoneme**) has many **microtubules** running parallel to the long axis.
- The axoneme has 9 pairs of doublets of radially arranged peripheral microtubules and a pair of central microtubules. This is called **9+2 array**.
- The central tubules are connected by bridges and are enclosed by a **central sheath**. It is connected to one of the tubules of each peripheral doublet by a **radial spoke**. Thus, there are **9 radial spokes**. The peripheral doublets are also interconnected by linkers.
- Cilium and flagellum emerge from centriole-like structure called the **basal bodies**.

9. Centrosome and Centrioles

- **Centrosome** is an organelle usually containing two non-membrane bound cylindrical structures called **centrioles**.
- They are surrounded by **pericentriolar materials**.
- The centrioles lie perpendicular to each other. They are made up of 9 evenly spaced peripheral fibrils of **tubulin**. Each of the peripheral fibril is a triplet. The adjacent triplets are also linked.
- The central part of the centriole is also proteinaceous and called the **hub**, which is connected with tubules of the peripheral triplets by radial **spokes** made of protein.
- The centrioles form the basal body of cilia or flagella, and spindle fibres that give rise to spindle apparatus during cell division in animal cells.

10. Nucleus


- Nucleus was first described by **Robert Brown** (1831).
- The material of the nucleus stained by the basic dyes was given the name **chromatin** by **Flemming**.




- Normally, a cell has only one nucleus. Some cells have more than one. Some mature cells lack nucleus. E.g. mammalian RBC and sieve tube cells of vascular plants.
- The interphase nucleus contains
 - o **Nuclear envelope:** Double layered membrane with a space between (10 - 50 nm) called **perinuclear space**. It is a barrier between the materials present in nucleus & cytoplasm. Outer membrane usually remains continuous with ER and also bears ribosomes on it. Nuclear envelope has minute pores formed by the fusion of its two membranes. These are the passages for the movement of RNA and protein between nucleus and cytoplasm.
 - o **Nuclear matrix (nucleoplasm)**
 - o **Chromatin:** A network of nucleoprotein fibres. It contains DNA and basic proteins (**histones**), non-histone proteins and RNA. During cell division, chromatins condense to form **chromosomes**.
 - o **Nucleolus:** One or more non-membranous spherical bodies. It is continuous with the nucleoplasm. It is a site for ribosomal RNA synthesis.

Chromosomes:

- A human cell has 2 m long thread of DNA distributed among its 46 (23 pairs) chromosomes.
- Every chromosome has a primary constriction (**centromere**). On the sides of centromere, disc shaped structures called **kinetochores** are present.



- Based on position of centromere, chromosomes are 4 types:
 - o **Metacentric chromosome:** Middle centromere forming two equal arms of the chromosome.
 - o **Sub-metacentric chromosome:** Centromere is nearer to one end forming one shorter arm and one longer arm.
 - o **Acrocentric chromosome:** Centromere is close to its end forming one very short and one very long arm.
 - o **Telocentric chromosome:** Terminal centromere.



- Some chromosomes have non-staining secondary constrictions at a constant location. It is called **satellite**.

11. Microbodies

- These are membrane bound minute vesicles that contain various enzymes.
- Present in both plant and animal cells.

Differences between Plant and animal cells

Plant cell	Animal cell
1. Cell wall present	Absent
2. Plastids are present	Absent
3. A large central vacuole	Many small vacuoles
4. Centrioles are absent	Present

COMPARISON BETWEEN PROKARYOTIC AND EUKARYOTIC CELLS

Prokaryotic cells	Eukaryotic cells
1. Generally smaller	Larger
2. Genetic material is in the form of nucleoid	Genetic material is in the form of nucleus
3. Nuclear membrane absent	Present
4. Membrane bound organelles absent	Present
5. Circular DNA	Linear DNA
6. Ribosomes 70 S type	80 S type (70 S in plastids and mitochondria)


MODEL QUESTIONS

1. Match the columns A,B, & C

A	B	C
Mitochondria	Sedimentation coefficient	Spindle fibers
Golgi bodies	Hydrolytic enzyme	Power house
Lysosomes	Axoneme	Cisternae
Ribosomes	Centrioles	Acidic PH
Cilia	Glycoproteins	George Palade
Centrosome	Cristae	9+ 2

2. Bacterial cell envelope is having a complex structure. Name the layers of the envelope.

3. Types of Chromosomes based on the position of centromere are given. Name the Chromosomes.



4. In cells glycoprotein & glycolipids are secreted by a cell organelle.

- a. Name the cell organelle
- b. Neatly draw its diagram

5. Plastids are found in all plant cells


- a. List the three plastids found in plants.
- b. Name the colorless plastids and specify its role.

6. Identify the characters of prokaryotic cells from the following statements

- | | |
|----------------------------------|----------------------|
| a. Endoplasmic reticulum present | b. Ribosome present |
| c. Golgi bodies absent | d. Incipient nucleus |
| e. Yeast is an example | f. Mostly anaerobes |

7. Golgi apparatus remains in close association with the endoplasmic reticulum. Give the reason.

8. Copy the following diagram and label the parts.



9. BIOMOLECULES

Biomolecules are chemical compounds found in living organisms. They include organic and inorganic compounds.

ANALYSIS OF CHEMICAL COMPOSITION IN A TISSUE

- Grind a living tissue (vegetable or piece of liver etc.) in **trichloroacetic acid (Cl₃CCOOH)** to get thick slurry.
- Strain this through a cheesecloth or cotton to get 2 fractions such as **filtrate (acid-soluble pool)** and the **retentate (acid-insoluble fraction)**.
- The filtrate contains **biomolecules** (biomolecules having molecular weight **less than 1000 Dalton**).
- The retentate contains **biomacromolecules** (biomolecules having molecular weight **higher than 1000 Dalton**).

Analysis of inorganic compounds

- Weigh a living tissue (wet weight) and it is dried (dry weight) to evaporate water.
- It is fully burnt to oxidize all carbon compounds to gaseous form (CO₂ & water vapour). It forms ash.

- The ash contains **inorganic elements** (Ca, Mg, Na, K etc.) & **inorganic compounds** (SO₄²⁻, PO₄³⁻, NaCl, CaCO₃ etc.).


Comparison of Elements in Non-living & Living Matter		
Element	% Weight of	
	Earth's crust	Human body
Hydrogen (H)	0.14	0.5
Carbon (C)	0.03	18.5
Oxygen (O)	46.6	65.0
Nitrogen (N)	Very little	3.3
Sulphur (S)	0.03	0.3
Sodium (Na)	2.8	0.2
Calcium (Ca)	3.6	1.5
Magnesium (Mg)	2.1	0.1
Silicon (Si)	27.7	Negligible

BIOMICROMOLECULES (MICROMOLECULES OR BIOMOLECULES)

Molecular weight of micromolecules found in the acid soluble pool ranges from **18 to 800 Dalton (Da)**. The acid soluble pool represents the cytoplasmic composition. They include amino acids, sugars, nitrogen bases, lipids etc.

1. AMINO ACIDS

- They are the compounds formed of an **amino group (-NH₂)**, an **acid group (-COOH)**, H & a variable group (R).
- -NH₂ & -COOH are attached to the same carbon atom (α -carbon). So, they are called α -amino acids.
- They are substituted methanes.



20 types of amino acids are used for protein synthesis. They include


- **Acidic amino acids:** e.g. Glutamic acid, Aspartic acid.
- **Basic amino acids:** e.g. Lysine, Arginine.
- **Neutral amino acids:** e.g. Valine.

Some amino acids are **aromatic**. E.g. tyrosine, phenyl alanine and tryptophan.

Amino acids are 2 types:

- **Essential amino acids:** They cannot be synthesized by the body and should be supplied through diet. E.g. Lysine, leucine, isoleucine, tryptophan etc.
- **Non-essential amino acids:** They can be synthesized by the body. E.g. Glycine, alanine, serine, arginine etc.

In amino acids, -NH₂ & -COOH have ionizable nature. So, structure of amino acids changes in solutions of different pH. If both -NH₂ & -COOH are ionized, it is called **Zwitterion**.



2. LIPIDS

- Water insoluble.
- Contain C, H & O but number of oxygen atoms is less.

Types of lipids


a. Simple lipids: These are formed of **Fatty acids** and **alcohol** such as **glycerol**.

- Fatty acids are lipids with a hydrocarbon chain (R- group) ending in -COOH group. i.e. R-COOH.



E.g. **Palmitic acid** has 16 carbons (CH₃ - (CH₂)₁₄ - COOH) or C₁₅H₃₁-COOH) and **Arachidonic acid** has 20 Carbons.

- Fatty acids are 2 types:
 - **Saturated fatty acids:** They have no double or triple bonds between carbon atoms. E.g. Palmitic acid, Stearic acid (C₁₇H₃₅COOH) etc.
 - **Unsaturated Fatty acids:** They have one or more C=C bonds. E.g. Oleic acid (C₁₇H₃₃COOH), Arachidonic acid (C₁₉H₃₁COOH) etc.

- **Structure of glycerol (trihydroxy propane):**



- Fatty acids are esterified with glycerol through **ester bond** forming **monoglycerides, diglycerides & triglycerides**.




Based on melting point, lipids (triglycerides) are 2 types:


- **Fats:** Higher melting point.
- **Oils:** Lower melting point.

b. Compound lipids: These are the esters of fatty acids and alcohol with additional groups.

E.g. **Phospholipids** (fatty acids + glycerol + phosphorylated organic compound). They are found in cell membranes. **E.g. Lecithin.**




c. Derived lipids: These are the products of hydrolysis of simple lipids and compound lipids. E.g. **Cholesterol.**




3. SUGARS (CARBOHYDRATES)

Sugars are sweet and water-soluble carbohydrates. They are formed of C, H and O in the ratio of 1:2:1.



Glucose (C₆H₁₂O₆)




Ribose (C₅H₁₀O₅)

4. NITROGEN BASES


These are the nitrogen containing cyclic compounds found in nucleic acids. They are 2 types:

a. Purines: Includes **Adenine (A) & Guanine (G).**

b. Pyrimidines: Includes **Cytosine (C), Thymine (T) & Uracil (U).**




Adenine (Purine)




Uracil (Pyrimidine)

Nitrogen base + Sugar → Nucleoside

Adenine	+	Sugar	→	Adenosine
Guanine	+	Sugar	→	Guanosine
Cytosine	+	Sugar	→	Cytidine
Thymine	+	Sugar	→	Thymidine
Uracil	+	Sugar	→	Uridine




Adenosine (A + Sugar)



Uridine (U + Sugar)

Nitrogen base + Sugar + Phosphate → Nucleotide

Adenine	+	Sugar	+	Phosphate	→	Adenylic acid
Guanine	+	Sugar	+	Phosphate	→	Guanylic acid
Cytosine	+	Sugar	+	Phosphate	→	Cytidylic acid
Thymine	+	Sugar	+	Phosphate	→	Thymidylic acid
Uracil	+	Sugar	+	Phosphate	→	Uridylic acid



Adenylic acid

Nucleotides are heterocyclic compounds.

Nucleic acids (DNA & RNA) are made up of nucleotides.

BIOMACROMOLECULES (MACROMOLECULES)

- These are biomolecules having molecular weight greater than 1000 Da. They include

- o Proteins
 - o Polysaccharides
 - o Nucleic acids
- } Molecular weight is 10,000 Da and above.

- **Acid insoluble fraction** (macromolecular fraction) includes macromolecules from cytoplasm and organelles.

- Lipid is not strictly a macromolecule as its molecular weight does not exceed **800 Da**. But it comes under acid insoluble fraction because many lipids are arranged into structures like cell membranes. When a tissue is grinded, cell membranes are broken and form water insoluble vesicles. They cannot be filtered along acid soluble fraction.


Average composition of cells	Water	70-90 %
	Protein	10-15%
	Carbohydrates	3%
	Lipids	2%
	Nucleic acids	5-7%
	Ions	1%

1. PROTEINS

- Proteins are heteropolymer of amino acids.

- They are polypeptides. i.e., linear chains of amino acids linked by **peptide bonds**.

- Peptide bond is formed when -COOH group of one amino acid reacts with -NH₂ group of next amino acid by releasing a molecule of water (dehydration).



Functions of proteins

- o For growth and tissue repair.
- o Transport nutrients across cell membranes. E.g. GLUT-4 enables glucose transport into cell.
- o Acts as intercellular ground substance. E.g. collagen.
- o Acts as antibodies to fight infectious organisms.
- o Acts as receptors. E.g. receptors of smell, taste, hormones.
- o Some are hormones (e.g. Insulin), enzymes (e.g. trypsin), pigments (e.g. hemoglobin) etc.


Most abundant protein in animal world: **Collagen**

Most abundant protein in the biosphere: **Ribulose**


bisphosphate carboxylase - oxygenase (RuBisCO)

Structural levels of protein

- **Primary structure:** It describes the sequence of amino acids, i.e. the positional information in a protein. **Left end** of the chain has first amino acid (N-terminal amino acid). **Right end** has last amino acid (C-terminal amino acid).
- **Secondary structure:** Here, one or more polypeptide chains are folded in the form of a helix. It has only right-handed helices. E.g. Keratin, Fibroin (silk fibre).




- **Tertiary structure:** Here, helical polypeptide chain is further folded like a hollow woolen ball. It gives 3-D view. Tertiary structure is necessary for many biological activities of proteins. E.g. Myoglobin, enzymes.
- **Quaternary structure:** Here, more than one polypeptide chains form tertiary structure and each chain functions as subunits of protein. E.g. Haemoglobin. It has 4 subunits (2 α subunits and 2 β subunits).



2. POLYSACCHARIDES (COMPLEX CARBOHYDRATES)

These are polymers of sugars (monosaccharides). E.g.

- Starch (polymer of glucose)
 - Cellulose (polymer of glucose)
 - Glycogen (polymer of glucose)
 - Inulin (polymer of fructose)
- Homopolymers**
- There are complex polysaccharides formed of **amino-sugars** (e.g. glucosamine, N-acetyl galactosamine etc.).
 - **Chitin** is the homopolymer of **N-acetyl glucosamine**. It is seen in exoskeleton of arthropods and fungal cell wall.
 - **Glycosidic bond in polysaccharides:** It is the bond formed when individual monosaccharides are linked between 2 carbon atoms by dehydration.
 - Starch forms helical **secondary structure**. It can hold I_2 molecules in the helical portion giving **blue colour**.
 - Cellulose has no complex helices and so cannot hold I_2 .
 - **Diagrammatic representation of a portion of glycogen:**




3. NUCLEIC ACIDS (DNA & RNA)

- Nucleic acids are **heteropolymer** of **nucleotides**. i.e., many nucleotides are linked to form **polynucleotide**.
- Nucleic acids are 2 types: **DNA** (Deoxyribonucleic acid) and **RNA** (Ribonucleic acid).

Secondary structure of DNA (Watson - Crick Double Helix Model)

- There are more than a dozen forms of DNA such as A, B, C, D, E, Z etc.
- DNA consists of 2 **polynucleotide strands** arranged antiparallelly as a **double helix**.
- In DNA, a nucleotide consists of **nitrogen base, deoxyribose sugar and phosphate group**.
- Backbone (strands) of DNA is formed by the **sugar-phosphate-sugar chain**.
- Steps are formed of **Nitrogen base pairs**.
- Nitrogen bases include **Adenine (A), Guanine (G), Thymine (T) and Cytosine (C)**. Uracil absent.
- A pairs with T (**A=T**) by **2 hydrogen bonds**.
- G pairs with C (**G=C**) by **3 hydrogen bonds**.
- A phosphate molecule links the **3'-carbon** of the sugar of one nucleotide to the **5'-carbon** of the sugar of the next nucleotide. The bond between the phosphate and -OH group of sugar is an **ester bond**. As there is one such ester bond on either side, it is called **phosphodiester bond**.
- The bond between sugar and nitrogen base is called **N-glycosidic bond**.



In B-DNA:

- One full turn of helical strand has **10 steps (10 base pairs)**.
- Length of one full turn = **34 Å** (i.e. **3.4 Å** for each step).
- At each step, the strand turns **36°** (**360°** for a full turn).

METABOLISM

- All the biochemical reactions taking place inside a living system together constitute **metabolism**. E.g.
 - Removal of CO₂ from amino acids to form amine.
 - Removal of amino group in a nucleotide base.
 - Hydrolysis of a glycosidic bond etc.
- The intermediate products of metabolic reactions are called **Metabolites**.
- Flow of metabolites in metabolic pathway has a definite rate & direction like automobile traffic. This metabolite flow is called **dynamic state of body constituents**.
- Metabolites are 2 types:
 - **Primary metabolites:** They have identifiable functions in physiological processes and necessary for life. E.g. amino acids, sugars, nucleic acids, lipids, vitamins etc.
 - **Secondary metabolites:** They are not directly involved in normal growth, development or reproduction. They are found in plant, fungal and microbial cells. E.g.
 - ▶ **Pigments:** Carotenoids, Anthocyanins etc.
 - ▶ **Alkaloids:** Morphine, Codeine etc.
 - ▶ **Terpenoids:** Monoterpenes, Diterpenes etc.
 - ▶ **Essential oils:** Lemongrass oil etc.
 - ▶ **Toxins:** Abrin, Ricin etc.
 - ▶ **Lectins:** Concanavalin A.
 - ▶ **Drugs:** Vinblastine, curcumin etc.
 - ▶ **Polymeric substances:** Rubber, gums, cellulose etc.
- In metabolism, there is a series of linked multistep chemical reaction called **metabolic pathways**. It is 2 types:

Anabolic (Biosynthetic) pathway	Catabolic pathway
Simpler molecules form complex structures (Constructive process).	Complex molecules become simple structures (destructive process).
It consumes energy.	It releases energy.
E.g. acetic acid becomes cholesterol, assembly of amino acids to protein, photosynthesis etc.	E.g. glucose becomes lactic acid (glycolysis), respiration etc.

- The energy released through catabolism is stored in the form of chemical bonds. When needed, this bond energy is utilized for biosynthetic, osmotic and mechanical works.
- The most important energy currency in living system is the bond energy in **adenosine triphosphate (ATP)**.

THE LIVING STATE

- In organisms, the metabolites are present in different concentrations. E.g. Blood concentration of glucose in a normal person is 4.2 - 6.1 mmol/L. Concentration of hormones is nanograms/mL.
- Systems at equilibrium cannot perform work. As living organisms work continuously, they cannot reach equilibrium. i.e. **“The living state is a non-equilibrium steady-state to be able to perform work”**.
- Living process is a constant effort to prevent falling into equilibrium. This is achieved by energy input obtained from metabolism. So, no living state without metabolism.


ENZYMES

- Enzymes are **biological catalysts** which influence the speed of biochemical reactions.
- *All enzymes are proteins but all proteins are not enzymes.*
- Enzymes are specific. i.e. each enzyme has its own substrate.
- **Ribozymes:** Nucleic acids (RNA) that behave like enzymes.
- Enzymes form tertiary structure (3D) with some crevices (pockets) called **‘active site’** into which the substrate fits.

Chemical Reactions

- Chemical compounds undergo two types of changes:
 - **Physical change:** A change in shape or state of matter without breaking bonds. E.g. ice melts into water, water becomes vapour.
 - **Chemical change (chemical reaction):** In this, bonds are broken and new bonds are formed. It may be organic or inorganic reaction. E.g.
 - Inorganic:** $\text{Ba(OH)}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2\text{H}_2\text{O}$
 - Organic:** $\text{Starch} + \text{H}_2\text{O} \xrightarrow{\text{Hydrolysis}} \text{H}_2\text{CO}_3$
- Rate of a physical or chemical process = Amount of product formed per unit time. i.e. $\frac{\delta p}{\delta t}$
- Rate is called **velocity** if the direction is specified.
- Rates of physical and chemical processes are influenced by factors such as temperature. Generally, rate doubles or decreases by half for every 10°C change in either direction.

- Rate of enzyme catalysed reactions is very high. E.g. **Carbonic anhydrase** is the fastest enzyme. It accelerates the following reaction 10 million times.





- In the absence of enzyme, only 200 molecules of H₂CO₃ are formed in an hour. In the presence of *carbonic anhydrase* about 600,000 molecules are formed per second.
- In a metabolic pathway, each step is catalysed by different enzymes.
 - E.g. In **glycolysis** [Glucose (C₆H₁₂O₆) → 2 Pyruvic acid (C₃H₄O₃)], ten different enzymes take part.

Nature of enzyme action (catalytic cycle)


Enzyme acts with substrate like a **lock & key model** action. It includes the following steps:

- The substrate binds to the **active site** of enzyme (**E+S**).
- This induces some changes in enzyme so that the substrate is tightly bound with active site of enzyme to form **enzyme-substrate complex (ES)**.
- The active site breaks chemical bonds of substrate to form **enzyme- product complex (EP)**.
- The enzyme releases the products and the free enzyme is ready to bind to other molecules of the substrate (**E+P**).



This pathway goes through some unstable *transition state structures*.

How do Enzymes Speed up a chemical Reaction? (Concept of activation energy)




- **Activation energy** is the additional energy required to start a chemical reaction.
- In an exothermic or endothermic reaction, the substrate must go through a much higher energy state. It is called **transition state energy**. Therefore, activation energy is the difference between average energy of substrate and transition state energy.
- If the product (P) is at a lower energy level than the substrate (S), the reaction is an **exothermic reaction (spontaneous reaction)**. It requires no energy (by heating) to form the product.
- In a biochemical reaction, enzymes lower the activation energy. As a result, speed of the reaction increases.

Factors affecting enzyme activity

a) Temperature and pH


- o Enzymes show highest activity at optimum temperature & pH. Activity declines below and above optimum value.
- o At low temperature, enzyme temporarily inactive.
- o At high temperature, enzymes destroy because proteins are denatured by heat.



- o Inorganic catalysts work at high temperature & pressure. But enzymes get damaged at high temperature ($> 40^{\circ}C$).
- o Thermophilic organisms have enzymes which are stable at high temperature (up to $80-90^{\circ}C$).

b) Concentration of substrate

- With the increase in substrate concentration, the velocity of enzyme action rises at first and reaches a *maximum velocity (Vmax)*. This is not exceeded by further rise in




concentration because enzyme molecules are fewer than the substrate molecules i.e. No free enzyme molecules to bind with additional substrate molecules.

c) Presence of Inhibitor

- The binding of specific chemicals (inhibitor) shuts off the enzyme activity. This is called **inhibition**.
- The inhibitor closely similar to the substrate is called **competitive inhibitor**. It competes with the substrate for the binding site of the enzyme. As a result, the substrate cannot bind and the enzyme action declines.

E.g. *Malonate* is similar to the substrate *succinate*. So, it inhibits *succinic dehydrogenase* in the following reaction.



- Competitive inhibitors are used to control bacterial pathogens.

Classification and nomenclature of enzymes

- **Oxido-reductases / Dehydrogenases:** Catalyze oxido-reduction b/w two substrates.
 $S \text{ reduced} + S' \text{ oxidized} \rightarrow S \text{ oxidized} + S' \text{ reduced}$
- **Transferases:** Catalyze transfer of a group (other than hydrogen).
 $S-G + S' \rightarrow S'-G + S$
- **Hydrolases:** Catalyze hydrolysis of ester, ether, peptide, glycosidic, C-C, C-halide or P-N bonds.
- **Lyases:** Catalyze removal of groups by mechanisms other than hydrolysis leaving double bonds.
 $X-C-C-Y \rightarrow X-Y + C=C$
- **Isomerases:** Catalyze inter-conversion of optical geometric or positional isomers.
- **Ligases:** Catalyze the linking of 2 compounds together.
E.g. enzymes catalyzing joining of bonds like C-O, C-S, C-N, P-O etc.

Co-factors





- These are non-protein constituents bound to the enzyme to make the enzyme catalytically active.
- **Apo-enzyme:** Protein portion of the enzyme.
- **Co-factor + Apoenzyme = Holoenzyme.**
When the co-factor is removed from the enzyme, its catalytic activity is lost.

Co-factors are 3 types:

- **Prosthetic group:** Organic. Tightly bound to apoenzyme.
E.g. **Haem**. It is a part of the active site of *peroxidase* and *catalase*. These enzymes catalyze breakdown of H_2O_2 to water & O_2 .
- **Co-enzymes:** Organic. Transient binding to apoenzyme. Many co-enzymes contain vitamins. E.g. **nicotinamide adenine dinucleotide (NAD)** and **NADP** contain **niacin**.
- **Metal ions:** They form **co-ordination bonds** with side chains at active site and one or more co-ordination bonds with the substrate.
E.g. **Zn** is a cofactor for *Carboxypeptidase*.


MODEL QUESTIONS

1. Observe the following diagrams and answer the questions.



- Identify A to D.
- Mention any one difference between A and B.
- Name any two biomolecules obtained by the polymerisation of A.
- Name the compound obtained by the fusion of one 'C' and three 'D'.

2.



- Identify the compound.
- Name the bond produced when another biomolecule of same category combines with this.
- If many such molecules are bonded together, what will be the resultant molecule?

3. Fill in the blanks marked as a, b, c and d.

Base	Nucleoside	Nucleotide
Adenine	Adenosine	Adenylic acid
Guanine	a	Guanylic acid
b	Cytidine	Cytidylic acid
Thymine	c	d

4. Match the following

A	B	C
Amino acid	Ester bond	Triglyceride
Glucose	Hydrogen bond	Protein
Nitrogen bases	Peptide bond	Starch
Fatty acid and glycerol	Glycosidic bond	Nucleic acid


5. Note the relationship between first two words and fill up the fourth place.

- Keratin: Secondary structure Haemoglobin:
- Glycogen: Polysaccharide GLUT 4:

6. Odd man out. Justify your answer.

- Glucose, Lectins, Glycine, Adenine
- Adenosine, Guanosine, Cytosine, Uridine

7. Analyze the graph showing the activity of salivary amylase.



- Which is the optimum temperature for salivary amylase obtained from the graph?
- Why the activity declines below and above the optimum value?

8. Non-protein constituents called cofactors are bound to the enzyme to make the enzyme catalytically active.

- Name the protein portion of the enzyme.
- What happens to the catalytic activity when the cofactor is removed from the enzyme?
- Mention any two kinds of cofactors with examples.

Visit: www.bankofbiology.com

10. CELL CYCLE AND CELL DIVISION


- The growth and reproduction of all organisms depend on the division and enlargement of cells.
- The mechanisms of division and multiplication of cells together constitute **cell reproduction**.

CELL CYCLE

- It is the life period of a cell during which a cell synthesizes DNA (replication), grows & divides into 2 daughter cells.
- Cell growth (cytoplasmic increase) is a continuous process but DNA synthesis occurs only at a specific stage.
- Duration of cell cycle varies in each organism and each cell type. E.g. Duration of a typical eukaryotic cell cycle (e.g. human cell) is about 24 hrs. In Yeasts, it is 90 minutes.

Phases of Cell Cycle

Cell cycle includes 2 basic phases: **Interphase & M Phase**.



1. Interphase (resting phase)

- It is the phase between two successive M phases.
- It includes cell growth and DNA synthesis.
- It lasts more than 95% of the duration of cell cycle.

Interphase has 3 phases:

a. G₁ phase (Gap 1 or Antephase): First growth phase. It is the interval between mitosis and DNA replication.

Main events:

- Continuous growth of cell.
- Cell becomes metabolically active.
- Prepares machinery for the DNA replication.
- Synthesizes RNA and proteins.

b. S (Synthetic) phase:

- In this, DNA replication takes place.
- Amount of DNA per cell doubles. But chromosome number is not increased.
- In animal cells, replication begins in the nucleus, and the centriole duplicates in the cytoplasm.

c. G₂ phase (Gap 2):

- Second growth phase. Cell growth continues.
- Synthesis of RNA and proteins continues.
- Cell is prepared for mitosis.

2. M Phase (Mitosis phase)

- It represents the actual cell division (mitosis).
- In human cell cycle, it lasts for only about an hour.
- M Phase includes **karyokinesis** (nuclear division) and **cytokinesis** (division of cytoplasm).
- Some cells do not show division. E.g. heart cells.
- Many other cells divide only occasionally to replace damaged or dead cells.
- The cells that do not divide further exit G₁ phase and enter an inactive stage called **quiescent stage (G₀)**. Such cells remain metabolically active but do not proliferate.

MITOSIS

- It is the cell division occurring in **somatic cells**.
- It is also called as **equational division** as the number of chromosomes in the parent and progeny cells is same.
- Mitosis is generally seen in **diploid cells**. It also occurs in haploid cells of some lower plants and some social insects.
- It involves major reorganization of all cell components.

The karyokinesis of mitosis has 4 stages: **Prophase, Metaphase, Anaphase & Telophase**.

1. Prophase

- It is the longest phase in mitosis.
- It follows the S and G₂ phases of interphase.
- In the S & G₂ phases, DNA molecules are intertwined.
- **Characteristic events:**
 - Chromosomal materials (chromatin fibres) are untangled and condensed to form mitotic **chromosomes**. They are seen to be composed of **two chromatids** attached together at the **centromere**.

- **Centrosomes** begin to move towards opposite poles of the cell. Each centrosome radiates out microtubules called **asters**. The two asters together with spindle fibres forms **mitotic apparatus**.

- Cells at the end of prophase do not show Golgi complexes, endoplasmic reticulum, nucleolus & nuclear envelope.

2. Metaphase

- The nuclear envelope completely disintegrates. Hence the chromosomes spread through the cytoplasm of the cell.
- Chromosome condensation is completed. They can be observed and studied easily under the microscope. They will have two sister chromatids.
- Chromosomes come to lie at the equator. The plane of alignment of the chromosomes at metaphase is called the **metaphase plate**.
- The **spindle fibres** from both poles are connected to chromatids by their kinetochores in the centromere.

3. Anaphase


- It is the shortest phase in the mitosis.
- Centromere of each chromosome divides longitudinally resulting in the formation of two daughter chromatids (chromosomes of the future daughter nuclei).
- As the spindle fibres contract, the chromatids move from the equator to the opposite poles.

4. Telophase

- Chromosomes cluster at opposite poles and uncoil into chromatin fibres.
- Nuclear envelope develops around the chromosome clusters at each pole. Thus 2 daughter nuclei are formed.
- Nucleolus, Golgi complex and ER reappear.
- The spindle fibres disappear.

Cytokinesis

- It is the division of cytoplasm to form 2 daughter cells. It



starts when telophase is in progress.

- **Cytokinesis in animal cell:** Here, a **cleavage furrow** is appeared in the plasma membrane. It gradually deepens and joins in the centre dividing the cytoplasm into two.
- **Cytokinesis in plant cell:** It is different from the cytokinesis in animal cells due to the presence of cell wall. In plant cells, the vesicles formed from Golgi bodies accumulate at the equator. It grows outward and meets the lateral walls. They fuse together to form the **cell-plate**. It separates the 2 daughter cells. Later, the cell plate becomes the middle lamella.
- During cytokinesis, organelles like mitochondria and plastids get distributed between the daughter cells.
- In some organisms karyokinesis is not followed by cytokinesis. As a result, multinucleate condition (syncytium) arises. E.g. liquid endosperm in coconut.

Significance of Mitosis

- It produces diploid daughter cells with identical genome.
- It helps to retain the same chromosome number in all somatic cells.
- It helps in the body growth of multicellular organisms. Mitosis in the meristematic tissues helps in a continuous growth of plants throughout the life.
- It restores the nucleo-cytoplasmic ratio that disturbed due to cell growth.
- It helps in cell repair & replacement. E.g. cells of the upper layer of the epidermis, lining of the gut & blood cells.

MEIOSIS

- It is the division of diploid germ cells that reduces the chromosome number by half forming haploid daughter cells (gametes). It occurs during gametogenesis.
- It leads to the haploid phase in the life cycle of sexually reproducing organisms. Fertilisation restores diploid phase.

Key features of meiosis

- It involves two cycles (**meiosis I & meiosis II**) but only a single cycle of DNA replication.
- It involves **pairing** of homologous chromosomes and **recombination** between their **non-sister chromatids**.
- Meiosis I begins after replication of parental chromosomes to form identical sister chromatids at the S phase.
- 4 haploid cells are formed at the end of meiosis II.

Meiosis I	Meiosis II
Prophase I	Prophase II
Metaphase I	Metaphase II
Anaphase I	Anaphase II
Telophase I	Telophase II

Meiosis I

Prophase I:

- It is typically longer and more complex.

- It includes 5 phases based on chromosomal behaviour: Leptotene, Zygotene, Pachytene, Diplotene & Diakinesis.

- **Leptotene (Leptonema):** Chromatin fibres become long slender chromosomes. Nucleus enlarges.
- **Zygotene (Zygonema):** Chromosomes become more condensed. Similar chromosomes start pairing together (synapsis) with the help of a complex structure called **synaptonemal complex**. The paired chromosomes are called **homologous chromosomes**. Each pair of homologous chromosomes is called a **bivalent**.
- **Pachytene (Pachynema):** Comparatively longer phase. Bivalent chromosomes split into similar chromatids. This stage is called **tetrads**. During this, **recombination nodules** appear at which **crossing over** occurs. It leads to genetic recombination on homologous chromosomes.

Crossing over: The exchange of genetic material between non-sister chromatids of two homologous chromosomes in presence of an enzyme, **recombinase**.

Recombination is completed by the end of pachytene.

- **Diplotene (Diplonema):** Dissolution of the synaptonemal complex occurs. The recombined homologous

chromosomes of the bivalents separate from each other except at the sites of crossovers. These X-shaped structures are called **chiasmata**. In oocytes of some vertebrates, diplotene lasts for months or years.

- **Diakinesis:** Terminalisation of chiasmata. Chromosomes are fully condensed. The meiotic spindle fibres originate from the poles to prepare the homologous chromosomes for separation. Nucleolus & nuclear envelope disappear.

Metaphase I:

Spindle formation is completed. The chromosomes align on the equatorial plate. The microtubules from the spindle attach to the pair of homologous chromosomes.

Anaphase I:

The homologous chromosomes separate, while sister chromatids remain associated at their centromeres.

Telophase I:

- The nuclear membrane and nucleolus reappear and 2 haploid daughter nuclei are formed. This is called **diad**.
- After this, cytokinesis may or may not occur.
- After a short interphase, it is followed by meiosis II.
- This short stage between the two meiotic divisions is called **interkinesis**. DNA replication does not occur in this phase.

Meiosis II

It resembles the mitosis. It has the following phases:

Prophase II:

It is initiated immediately after cytokinesis. The chromosomes again become compact.

Nucleolus and nuclear membrane disappear in both nuclei.

Metaphase II:

The chromosomes align at the equator and the microtubules from opposite poles of the spindle get attached to the kinetochores of sister chromatids.

Anaphase II:


It begins with the simultaneous splitting of the centromere of each chromosome (which was holding sister chromatids together). Thus they move toward opposite poles of the cell by shortening of microtubules attached to kinetochores.

Telophase II:

The two groups of chromosomes once again get enclosed by a nuclear envelope; cytokinesis follows resulting in the formation of tetrad of cells i.e., 4 haploid daughter cells.

Significance of meiosis

- It conserves the chromosome number of each species.
- It causes genetic variation (due to crossing over) in the population of organisms. It is important for evolution.



MODEL QUESTIONS

1. In a vegetative cell and reproductive cell, chromosomes get separated during Anaphase. Write the difference in the two cells during this stage.
2. Life cycle of a cell is called cell cycle. It consists of four stages such as G1, S, G2 and M.
 - a. Construct a pie diagram showing different stages indicated above
 - b. State the major events occurring in G1, S and G2 phases.
3. Identify the stage of mitosis.
 - Four chromosomes arranged on the equatorial plane.
 - Spindle fibres attached to the centromeres of chromosomes.
 - a. How many daughter cells will produce from mitosis?
 - b. Write the number of chromosomes in each daughter cell
 - c. Compare this stage of mitosis with the same stage in meiosis
4. Crossing over leads to recombination of genetic material between two homologous chromosomes.
 - a. In which stage of meiosis, this phenomenon is seen?
 - b. Give its significance.
5. Interphase lasts for more than 95% of the duration of cell cycle. Justify this statement
6. Cytokinesis differ in plant and animal cell, comment on this statement.
7. Match the following

A		B
Zygotene	-	Chiasmata
Pachytene	-	Terminalisation
Diplotene	-	Recombination Nodules
Diakinesis	-	Bivalent

8. The given diagram is a stage of mitosis



(a) Identify the stage of mitosis

(b) Write any one feature of this stage

13. PHOTOSYNTHESIS

- **Photosynthesis** is a **physico-chemical process** by which green plants use **light energy (solar energy)** to synthesise organic compounds. So they are autotrophs.
- It is the basis of life on earth.

- Ultimately, all living forms depend on sunlight for energy.

Importance of Photosynthesis

- It is the primary source of all food on earth.
- It releases oxygen into the atmosphere.

EXPERIMENTS RELATED WITH PHOTOSYNTHESIS

1. Variegated leaf experiment

- Take a variegated leaf (or leaf partially covered with black paper) that was exposed to light.
- Test the leaves for starch. It shows that photosynthesis occurs only in green parts of the leaves in presence of light.

2. Half-leaf experiment

- A part of a leaf is enclosed in a test tube containing KOH soaked cotton (which absorbs CO₂).
- The other half of leaf is exposed to air.
- Place this setup in light for some time.
- Test the leaf for presence of starch. Exposed part shows positive for starch and portion in the tube shows negative. This proves that CO₂ is required for photosynthesis.

EARLY EXPERIMENTS

Experiments by Joseph Priestley (1770)

- Priestley performed experiments to prove the role of air in the growth of green plants.
- He discovered oxygen in 1774.
- He observed that a candle burning in a closed bell jar gets extinguished. Similarly, a mouse suffocated in closed jar. He concluded that a burning candle or a breathing animal damage the air.
- He placed a mint plant in the same bell jar. He found that the mouse stayed alive and the candle continued to burn.
- He hypothesised that plants restore the air whatever breathing animals and burning candles remove.

Experiments by Jan Ingenhousz (1730-1799)


- He conducted the same experiment by placing in darkness and sunlight.
- He showed that sunlight is essential to the plant for purifying the air fouled by burning candles or animals.
- He repeated this experiment with an aquatic plant. It showed that in bright sunlight, small bubbles were formed around green parts while in the dark they did not.
- Later he identified these bubbles to be of oxygen. Thus he showed that only the green part of plants release O₂.

Experiments by Julius von Sachs (1854)

- He proved that
 - o Glucose is produced when plants grow and it is usually stored as starch.
 - o **Chlorophyll** is located in special bodies (**chloroplasts**).
 - o Glucose is made in the green parts of plants.

Experiments by T.W Engelmann (1843 – 1909)


- He split the light using a prism into its spectral components and illuminated a green alga (*Cladophora*) placed in a suspension of aerobic bacteria.
- The bacteria were used to detect the sites of O₂ evolution.
- He observed that the bacteria accumulated mainly in the region of blue and red light of the split spectrum.
- It was a first described **action spectrum** of photosynthesis. It resembles the absorption spectra of chlorophyll *a* & *b*.
- By the middle of 19th century, it is discovered that plants use light energy to make carbohydrates from CO₂ & H₂O.
- Empirical equation of the process of photosynthesis is




Where, [CH₂O] represents a carbohydrate (e.g. glucose).

Experiments by Cornelius van Niel (1897-1985)

- Van Niel (microbiologist) conducted some studies in purple and green bacteria.
- He demonstrated that photosynthesis is a light-dependent reaction in which hydrogen from an oxidisable compound reduces CO₂ to carbohydrates.




- In plants, H₂O is the hydrogen donor and is oxidised to O₂.
- Purple & green sulphur bacteria use H₂S as H-donor. So the 'oxidation' product is sulphur or sulphate and no O₂ is produced.
- Thus, he inferred that the O₂ evolved by the green plant comes from H₂O, not from CO₂. This was later proved by using radio isotopic techniques.
- Therefore overall correct equation for photosynthesis is:



PHOTOSYNTHESIS: SITE AND PIGMENTS


- Photosynthesis occurs in green leaves & other green parts.
- **Chloroplasts** present in the walls of **mesophyll cells** of leaves. It helps to get optimum quantity of incident light.
- Chloroplast contains a **membranous system**. It consists of **grana**, **stroma lamellae** and **matrix stroma**.
- Each granum is a group of membrane-bound sacs called **thylakoids (lamellae)**. They contain leaf pigments.

- The **membrane system** traps light energy and synthesise ATP and NADPH. It is called **light reactions**.
- In **stroma**, enzymatic reactions synthesise sugar, which in turn forms starch. It is called **dark reactions (carbon reactions)**. It does not mean that they occur in darkness or that they are not light dependent.




PIGMENTS INVOLVED IN PHOTOSYNTHESIS


- Pigments are substances that have ability to absorb light at specific wavelengths.
 - Chromatography shows the following leaf pigments:
 - o **Chlorophyll a** (bright or blue green in chromatogram)
 - o **Chlorophyll b** (yellow green)
 - o **Xanthophylls** (yellow)
 - o **Carotenoids** (yellow to yellow-orange)
- Accessory pigments**
- **Functions of accessory pigments:**
 - o They absorb light at different wavelength and transfer the energy to chlorophyll a.
 - o They protect chlorophyll a from photo-oxidation.
 - The **absorption spectrum & action spectrum** coincide closely showing that photosynthesis is maximum at the **blue & red regions** of the spectrum.
 - The graphs also show that chlorophyll a is the chief pigment associated with photosynthesis.



Graph showing absorption spectrum of chlorophyll a, b & carotenoids




Graph showing action spectrum of photosynthesis



Graph showing action spectrum of photosynthesis superimposed on absorption spectrum of chlorophyll a

Photosystems

- Pigments are organised into two **Photosystems** called **Photosystem I (PSI) & Photosystem II (PSII)**. These are named in the sequence of their discovery.
- Each photosystem has a **chlorophyll a** and **accessory pigments** bound by proteins.
- All pigments (except one molecule of chlorophyll a) form a **light harvesting complex (LHC or antennae)**.
- Single chlorophyll a acts as **reaction centre**.
- In **PS I**, the reaction centre absorbs light at **700 nm**, and so called **P700**.
- In **PS II**, the reaction centre absorbs light at **680 nm**, and so called **P680**.




LIGHT REACTION (PHOTOCHEMICAL PHASE)

- Light reactions include **light absorption, water splitting, oxygen release** and **formation of ATP & NADPH** (high-energy chemical intermediates).

The Electron Transport

- When **PS II** absorbs **red light of 680 nm** wavelength, electrons are excited and transferred to an electron acceptor.
- The electron acceptor passes them to a chain of **electrons transport system** consisting of **cytochromes**.
- This movement of electrons is downhill, in terms of redox potential scale.
- The electrons are transferred to the pigments of **PS I**.
- Simultaneously, electrons in **PS I** are also excited when they receive **red light of 700 nm** and are transferred to another accepter molecule having a greater redox potential.
- These electrons are moved downhill to a molecule of **NADP⁺**. As a result, **NADP⁺** is reduced to **NADPH + H⁺**.
- Transfer of electrons from **PS II** to **PS I** and finally downhill to **NADP⁺** is called the **Z scheme**, due to its zigzag shape. This shape is formed when all the carriers are placed in a sequence on a redox potential scale.



Splitting of Water (Photolysis)

- The **water splitting complex** in **PS II** is located on the inner side of the thylakoid membrane.
 - Water is split into **H⁺, [O]** and electrons.
- $$2\text{H}_2\text{O} \rightarrow 4\text{H}^+ + \text{O}_2 + 4\text{e}^-$$
- So **PS II** can supply electrons continuously by replacing electrons from water splitting.

- Thus PS II provides electrons needed to replace those removed from PS I.
- The protons (H^+) are used to reduce NADP to NADPH.
- Oxygen is liberated as a by-product of photosynthesis.

Photo-phosphorylation


- The synthesis of ATP by cells (in mitochondria & chloroplasts) is called **phosphorylation**.
- **Photo-phosphorylation** is the synthesis of ATP from ADP in chloroplasts in presence of light.
- It occurs in 2 ways: **Non-cyclic** and **Cyclic**.

a) Non-cyclic photo-phosphorylation

- It occurs when the two photosystems work in a series, (first PS II and then PS I) through an electron transport chain as seen in the Z scheme.
- Here, ATP & NADPH + H^+ are synthesised.
- It is a non-cyclic process because the electrons lost by PS II do not come back to it but pass on to NADP⁺.

b) Cyclic photo-phosphorylation

- It occurs in stroma lamellae when only PS I is functional.
- The electron is circulated within the photosystem and the ATP synthesis occurs due to cyclic flow of electrons.
- The lamellae of grana have PS I & PS II. The stroma lamellae membranes lack PS II and *NADP reductase*.
- The electron does not pass on to NADP⁺ but is cycled back to PS I complex through electron transport chain.
- Here, only ATP is synthesised (no NADPH + H^+).
- Cyclic photophosphorylation also occurs when only light of wavelengths beyond 680 nm are available.




Chemiosmotic Hypothesis

- It explains **mechanism of ATP synthesis** in chloroplast.
- **Chemiosmosis:** Movement of ions across a semipermeable membrane. It occurs in chloroplast and mitochondria.
- Chemiosmosis needs a membrane, a proton pump, a proton gradient (across thylakoid membranes) and *ATP synthase*.
- Splitting of water occurs on the inner side of the membrane. So the protons accumulate in the lumen of thylakoids.
- As electrons move through the photosystems, protons are transported across the membrane. It is due to the removal

of protons from the stroma for the following reasons:

- o **Primary electron acceptor** is located towards the outer side of the membrane. It transfers its electron to an **H carrier**. So this molecule removes a proton from the stroma while transporting an electron. When this molecule passes on its electron to the **electron carrier** on the inner side of the membrane, proton is released into the lumen of the membrane.
 - o The *NADP reductase* enzyme is located on the stroma side of the membrane. Along with electrons coming from PS I, protons are necessary to reduce NADP⁺. These protons are also removed from the stroma.
- Hence, protons in stroma are decreased but in lumen, protons are accumulated. It creates a proton gradient across the thylakoid membrane and decrease in pH in the lumen.



ATP synthesis through chemiosmosis

- Breakdown of proton gradient leads to synthesis of ATP by *ATP synthase* enzyme.
- The *ATP synthase* consists of two parts:
 - o **CF₀**: It is embedded in the membrane and forms a trans-membrane channel. It carries out facilitated diffusion of protons across the membrane to the stroma. It results in breakdown of proton gradient.
 - o **CF₁**: It protrudes on the outer surface of the thylakoid membrane. The energy due to breakdown of gradient causes a conformational change in the CF₁ particle. It makes the enzyme to synthesise ATP molecules.
- Energy is used to pump protons across a membrane, to create a gradient or a high concentration of protons within the thylakoid lumen.
- *ATP synthase* has a channel for the diffusion of protons back across the membrane. This releases energy to activate *ATP synthase* that catalyses formation of ATP.

DARK REACTION (BIOSYNTHETIC PHASE) - USE OF ATP & NADPH


- Products of light reaction are ATP, NADPH and O₂.
- **Dark reaction** is the use of ATP and NADPH to drive the processes for the synthesis of food (sugars).
- This phase does not directly depend on the light but is dependent on the products of the light reaction.
- It can be verified as follows: Immediately after light becomes unavailable, the biosynthetic process continues

- for some time, and then stops. If light is available, the synthesis starts again.
- CO₂ combines with H₂O to form (CH₂O)_n or sugars.
- CO₂ assimilation during photosynthesis is 2 types:
 - o **C₃ pathway:** In this, first stable product of CO₂ fixation is a C₃ acid (**3-phosphoglyceric acid - PGA**). **Melvin Calvin** discovered this using ¹⁴C in algal photosynthesis.

- o **C₄ pathway:** In this, first stable product is **oxaloacetic acid (OAA)**, a 4-carbon (C₄) organic acid.

C₃ PATHWAY (CALVIN CYCLE)

- It occurs in **all photosynthetic plants** (C₃ or C₄ pathways).
- It has 3 stages: carboxylation, reduction and regeneration.



1. Carboxylation of RuBP

- RuBP (**ribulose biphosphate** - a 5-carbon ketose sugar) is the primary CO₂ acceptor.
- It is the most crucial step. CO₂ is fixed by **RuBP** to two 3-PGA in presence of the enzyme **RuBP carboxylase**.
- Since this enzyme also has an oxygenation activity it is called **RuBP carboxylase-oxygenase (RuBisCO)**.
- RuBisCO is the most abundant enzyme in the world.

2. Reduction

- It is a series of reactions leading to the glucose formation.
- Here, 2 ATP molecules for phosphorylation and two of NADPH for reduction per CO₂ molecule are used.
- Fixation of 6 CO₂ molecules and 6 turns of the cycle are needed to remove one glucose molecule from the pathway.

3. Regeneration of RuBP

- It is crucial for continuation of the cycle.
- It requires one ATP for phosphorylation to form RuBP.
- Hence for every CO₂ molecule, 3 ATP molecules and 2 NADPH are required.
- It is probably to meet this difference in number of ATP and NADPH used in the dark reaction that the cyclic phosphorylation takes place.
- To make 1 glucose molecule, 6 turns of the cycle are needed.

What does go in and come out of the Calvin cycle?	In	Out
	6 CO ₂	1 glucose
	18 ATP	18 ADP
	12 NADPH	12 NADP


C₄ PATHWAY (HATCH & SLACK PATHWAY)

- It is present in plants adapted to dry tropical regions.
- They also use C₃ pathway as main biosynthetic pathway.
- The large cells around the vascular bundles of the C₄ plants are called **bundle sheath cells**. Such anatomy is called '**Kranz**' anatomy ('Kranz' = 'wreath').

- The bundle sheath cells may form **several layers** around the vascular bundles.
- They have large number of chloroplasts, thick walls impervious to gas exchange and no intercellular spaces.

Steps of Hatch and Slack Pathway

- Primary CO₂ acceptor is **phosphoenol pyruvate (PEP)** - a 3-carbon molecule seen in mesophyll cells. The enzyme for this fixation is **PEP carboxylase (PEPcase)**.
- The mesophyll cells lack **RuBisCO** enzyme.
- The C₄ acid OAA is formed in the mesophyll cells.
- It then forms other 4-carbon acids like malic acid or aspartic acid. They are transported to bundle sheath cells.




- In the bundle sheath cells, C₄ acids are broken down to release CO₂ and a C₃ molecule.
- The C₃ molecule is transported back to mesophyll where it is converted to PEP again.
- The released CO₂ enters the C₃ pathway.
- Bundle sheath cells are rich in **RuBisCO**, but lack **PEPcase**. Thus C₃ pathway is common to C₃ & C₄ plants.

C₄ plants are special because:

- o They have a special type of leaf anatomy (Kranz).
- o They tolerate higher temperatures.
- o They show a response to highlight intensities.
- o They lack photorespiration.
- o They have greater productivity of biomass.

PHOTORESPIRATION

- In Calvin pathway, RuBP combines with CO₂.



- Active site of RuBisCO can bind to CO₂ & O₂ - so the name.
- RuBisCO has a greater affinity for CO₂ than for O₂. This binding is competitive. Relative concentration of O₂ and CO₂ determines which one will bind to the enzyme.
- In C₃ plants, some O₂ bind to RuBisCO. Hence CO₂ fixation is decreased. Here RuBP binds with O₂ to form one

molecule of phosphoglycerate and phosphoglycolate. This pathway is called **photorespiration**.

- In this, there is no synthesis of sugars, ATP and NADPH. Hence **photorespiration is a wasteful process**. Rather it causes the release of CO₂ by using ATP.
- **In C₄ plants, photorespiration does not occur** because they can increase CO₂ concentration at the enzyme site.

This takes place when C₄ acid from the mesophyll is broken down in the bundle cells to release CO₂. This minimises the oxygenase activity of RuBisCO.

- Due to the lack of photorespiration, productivity and yields are better in C₄ plants. Also, these plants show tolerance to higher temperatures.

Differences between C₃ and C₄ plants


C ₃ plants	C ₄ plants
1. Photosynthesis occurs in mesophyll cells.	In mesophyll and bundle sheath cells.
2. Kranz anatomy is absent.	Present.
3. RuBP is the primary CO ₂ acceptor.	PEP is the primary CO ₂ acceptor.
4. 3-PGA, a 3-C compound is the first stable product.	OAA, a 4-C compound is the first stable product.
5. Chloroplasts are of only one type (granal).	Dimorphic (granal in mesophyll and agranal in bundle sheath).
6. Photorespiratory loss is high.	Photorespiration is absent or negligible.
7. High CO ₂ compensation point (25-100 μl. CO ₂ l ⁻¹).	Low CO ₂ compensation point (0-10 μl. CO ₂ l ⁻¹).
8. Optimum temperature for photosynthesis is about 25°C.	About 35°C - 45°C.
9. Photosynthetically less efficient and productivity low.	Photosynthetically more efficient and productivity high.
10. E.g. Rice, wheat, bean, potato.	E.g. Maize, sugarcane, amaranth, sorghum.

FACTORS AFFECTING PHOTOSYNTHESIS

- **Internal (plant) factors:** The number, size, age and orientation of leaves, mesophyll cells and chloroplasts, internal CO₂ concentration and amount of chlorophyll. Plant factors depend on the genes and growth of the plant.
- **External factors:** Sunlight, temperature, CO₂ concentration and water.
- **Blackman's Law of Limiting Factors (1905):** "If a biochemical process is affected by more than one factor, its rate is determined by the factor nearest to its minimal value: it is the factor which directly affects the process if its quantity is changed."
- E.g. a plant with green leaf, optimal light & CO₂ conditions may not photosynthesize if the temperature is very low. If optimal temperature is given, it will start photosynthesis.

Light

- **Light quality, light intensity and duration of exposure to light** influence photosynthesis.
- There is a linear relationship between incident light and CO₂ fixation rates at low light intensities.
- At higher light intensities, the rate does not show further increase because other factors become limiting.
- Light saturation occurs at 10% of the full sunlight. Hence, except for plants in shade or in dense forests, light is rarely a limiting factor in nature.
- High increase in incident light breaks down chlorophyll. It decreases photosynthesis.



Carbon dioxide Concentration

- CO₂ is the major limiting factor for photosynthesis.
- CO₂ concentration is very low in the atmosphere (0.03-0.04%). Increase up to 0.05% cause increase in CO₂ fixation rates. Beyond this level can become damaging over longer periods.
- At low light, C₃ and C₄ plants do not respond to high CO₂. At high light, they show increased rate of photosynthesis.
- C₄ plants show saturation at about 360 μL⁻¹.
- C₃ plants respond to increased CO₂ concentration and saturation is seen only beyond 450 μL⁻¹. Thus, current availability of CO₂ levels is limiting to the C₃ plants.
- Due to response to higher CO₂ concentration, C₃ plants show increased photosynthesis and higher productivity. This fact is used for some greenhouse crops (tomatoes, bell pepper etc). They are grown in CO₂ enriched atmosphere.

Temperature

- Dark reactions, being enzymatic, are temperature controlled. Influence of temperature on Light reactions is very less.
- The C₄ plants respond to higher temperatures and show higher rate of photosynthesis.
- C₃ plants have a much lower temperature optimum.
- The temperature optimum of plants also depends on their habitat. Tropical plants have a higher temperature optimum than the plants adapted to temperate climates.

Water

- Water stress closes the stomata hence reduce the CO₂ availability.
- Water stress also wilts leaves, thus reduce the surface area of the leaves and their metabolic activity.

14. RESPIRATION IN PLANTS

- **Oxidation of food materials** (breaking of C-C bonds of complex molecules) within the cell to release energy for ATP synthesis is called **cellular respiration**.
- This energy is used for absorption, transport, movement, reproduction, breathing etc.
- Ultimate source of food that is respired is photosynthesis.
- The compounds that are oxidized during respiration are called **respiratory substrates**. E.g. Carbohydrates (most common), proteins, fats and organic acids.
- The energy released is not used directly but is used to synthesize ATP. When energy is needed, ATP is broken down. Hence, **ATP** acts as **energy currency** of the cell.

BREATHING IN PLANTS

- For respiration, plants get O₂ and give out CO₂.
- In plants, gas exchange occurs via **stomata & lenticels**.
- Plants need no specialized respiratory organs because
 - Each plant part takes care of its own gas-exchange needs. So **gas transport is very limited**.
 - **Very low gas exchange** as compared to that of animals.
 - Leaves are adapted for maximum **gas exchange during photosynthesis**. During this, O₂ is released within the cell.
 - Most **living cells have contact with air**. They are located close to plant surface. In stems, living cells are organized in thin layers beneath the bark. They also have **lenticels**. In leaves, stems & roots, parenchyma cells are loosely packed that provides interconnected air spaces.
- Complete combustion of glucose yields energy most of which is given out as heat.


$$\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O} + \text{Energy}$$
- This energy is utilized to synthesize other molecules.
- During the glucose catabolism, not all the liberated energy goes out as heat. Glucose is oxidised in several small steps. It enables some steps to couple released energy to ATP synthesis.
- During respiration, oxygen is utilized, and CO₂, water & energy are released.
- Certain organisms are adapted to anaerobic conditions. Some are **facultative** anaerobes. Others are **obligate**.

GLYCOLYSIS (EMP PATHWAY)

- It is the partial oxidation (breakdown) of **glucose** to 2 molecules of **pyruvic acid** (C₃H₄O₃) in the absence of O₂.
- It occurs in cytoplasm of all living organisms.
- Its scheme was given by **Gustav Embden, Otto Meyerhof & J. Parnas**. So it is also known as **EMP pathway**.
- In anaerobes, it is the only process in respiration.
- In plants, glucose is derived from **sucrose** (end product of photosynthesis) or from storage **carbohydrates**. Sucrose is converted to glucose & fructose by an enzyme, **invertase**. These 2 monosaccharides readily enter glycolytic pathway.
- Glucose & fructose are phosphorylated to form glucose-6-phosphate by the enzyme **hexokinase**. It is then isomerised to produce fructose-6-phosphate. Subsequent steps of metabolism of glucose and fructose are same.

Steps of glycolysis:

- It includes 10 steps under the control of different enzymes.
- ATP is utilized at 2 steps:
 - In the conversion of glucose into glucose 6-phosphate.
 - In the conversion of fructose 6-phosphate to fructose 1, 6-diphosphate.
- Fructose 1, 6-diphosphate is split into dihydroxyacetone phosphate (**DHAP**) & 3-phosphoglyceraldehyde (**PGAL**).
- PGAL is oxidised and with inorganic phosphate get converted to 1, 3-bisphosphoglycerate (**BPGA**). During this, 2 redox-equivalents (2 H-atoms) are removed from PGAL and transferred to NAD⁺ forming NADH + H⁺.
- BPGA becomes 3-phosphoglyceric acid (**PGA**) yielding energy. This energy is trapped by the formation of ATP.
- ATP is also formed when PEP converts to **pyruvic acid**.
- In glycolysis, **4 ATP molecules** are directly synthesised from one glucose molecule.



Pyruvic acid (pyruvate) is the key product of glycolysis. Its metabolic fate depends on the cellular need.


In different cells, pyruvic acid is handled in 3 ways:

- Lactic acid fermentation.
- Alcoholic fermentation.
- Aerobic respiration (**Krebs' cycle**).

FERMENTATION (ANAEROBIC RESPIRATION)

- It is the **incomplete oxidation** of glucose under anaerobic condition.
- It occurs in many prokaryotes and unicellular eukaryotes.
- It is 2 types:
 - **Alcoholic fermentation:** Here, the **pyruvic acid** formed from glucose is converted to **CO₂** and **ethanol**. The enzymes, *pyruvic acid decarboxylase* and *alcohol dehydrogenase* catalyse these reactions. E.g. Yeast. Yeasts poison themselves to death when the concentration of alcohol reaches about 13%.
 - **Lactic acid fermentation:** Here, **pyruvic acid** is converted to **lactic acid**. E.g. Some bacteria.
- The reducing agent (**NADH+H⁺**) is reoxidised to **NAD⁺** in both the processes.
- In animals, when oxygen is inadequate during exercise, pyruvic acid in muscle cells is reduced to lactic acid by *lactate dehydrogenase*.
- **Net ATP production** from fermentation of one glucose molecule = **2**. (4 ATP from glycolysis – 2 ATP utilized).

- The steps involved in fermentation are shown below:




Drawbacks of fermentation

- **Energy production is limited.** Less than 7% of the energy in glucose is released and not all of it is trapped as high energy bonds of ATP.
- **Hazardous products** (acid or alcohol) are formed.

AEROBIC RESPIRATION

- It is a **complete oxidation** of organic substances in the presence of oxygen releasing CO₂, water & energy.
- It occurs in **mitochondria**.
- For this, the **pyruvate** (final product of glycolysis) is transported from the cytoplasm into the mitochondria.
- The crucial events in aerobic respiration are:
 - **Complete oxidation of pyruvate** by stepwise removal of all the hydrogen atoms, leaving 3 CO₂ molecules. It takes place in the **matrix of mitochondria**.
 - **Passing on of electrons** removed as part of H-atoms to molecular O₂ with simultaneous synthesis of ATP. It occurs on the **inner membrane of mitochondria**.
- Pyruvate (pyruvic acid) enters mitochondrial matrix and undergoes **oxidative decarboxylation** in presence *pyruvic dehydrogenase*. It needs coenzymes, NAD⁺ & Coenzyme A.
- During this process, 2 NADH molecules are produced from 2 pyruvic acid molecules.



- **Acetyl CoA** then enters **tricarboxylic acid (TCA) cycle**.

Tricarboxylic Acid Cycle (Krebs' cycle or Citric acid cycle)


TCA cycle was first elucidated by **Hans Krebs**.

Steps:

1. Condensation of **acetyl group** with **oxaloacetic acid (OAA)** & water to form **citric acid** in presence of *citrate synthase* enzyme. A **CoA** molecule is released.
2. **Citrate** is isomerised to **isocitrate**.
3. Decarboxylation of isocitrate to **α-ketoglutaric acid**.
4. Decarboxylation of α-ketoglutaric acid to **succinyl-CoA**.
5. Succinyl-CoA is converted to **succinic acid** and a **GTP** molecule is synthesised (substrate level phosphorylation).


In a coupled reaction, GTP is converted to GDP with simultaneous synthesis of ATP from ADP.

6. Oxidation of succinate to **Fumarate** and then to **Malate**.
7. Oxidation of malate to **OAA**.



- At 3 points of TCA cycle, **NAD⁺** is reduced to **NADH + H⁺**. At one point, **FAD⁺** is reduced to **FADH₂**.
- Continued oxidation of acetyl CoA via TCA cycle requires continued replenishment of OAA. It also requires regeneration of **NAD⁺** & **FAD⁺** from **NADH** & **FADH₂**.

Summary equation of Krebs' cycle:




Thus, a **glucose** is broken down to give **6 CO₂**, **8 NADH+H⁺**, **2 FADH₂** and **2 ATP**.

Electron Transport System (ETS) & Oxidative Phosphorylation

- **Electron transport system (ETS)** is the metabolic pathway present in the **inner mitochondrial membrane** through which electron passes from one carrier to another.

- This is to release and utilize energy stored in **NADH+H⁺** and **FADH₂** (formed during TCA cycle) by oxidation.
- The electrons are passed on to O₂ to form H₂O.
- Electrons from NADH are oxidised by an **NADH dehydrogenase (complex I)**.
- Electrons are then transferred to **ubiquinone (UQ)** located within the inner membrane. Ubiquinone also receives reducing equivalents via **FADH₂ (complex II)** that is generated during oxidation of succinate in citric acid cycle.
- The **reduced ubiquinone (ubiquinol or UQH₂)** is then oxidised with the transfer of electrons to **cytochrome c** via **cytochrome bc₁ complex (complex III)**. Cytochrome c is a small protein attached to the outer surface of the inner membrane. It acts as a mobile carrier of electrons between complex III and IV.
- **Complex IV (cytochrome c oxidase)** contains **cytochromes a & a₃**, and 2 copper centres.
- When the electrons pass from one carrier to another via complex I to IV, they are coupled to **ATP synthase (complex V)** for the ATP production.




Number of ATP molecules produced depends on nature of electron donor.

Oxidation of 1 NADH → 3 ATP
 Oxidation of 1 FADH₂ → 2 ATP

- In aerobic respiration, the role of oxygen is limited to the terminal stage. Yet, oxygen is vital since it drives the whole process by removing hydrogen from the system. Oxygen acts as the **final hydrogen acceptor**.
- In respiration, energy of **oxidation-reduction** is utilised for the phosphorylation. So this process is called **oxidative**

phosphorylation. It is not as photophosphorylation (Here, light energy is utilised to produce proton gradient for phosphorylation).

- The energy released during the ETS is utilized to synthesize ATP by **ATP synthase (complex V)**.
- **ATP synthase** has two major components: **F₁ & F₀**.
- **F₁ headpiece (peripheral membrane protein complex):** Site for ATP synthesis from ADP & inorganic phosphate.
- **F₀ (integral membrane protein complex):** It forms a channel through which protons cross the inner membrane. The passage of protons is coupled to the catalytic site of the F₁ component for ATP production.



- For each ATP produced, 2H⁺ passes through F₀ from the inter-membrane space to the matrix down the electrochemical proton gradient.

THE RESPIRATORY BALANCE SHEET

- Net gain of ATP from each glucose molecule is calculated based on the following assumptions:
 - All steps in Glycolysis, TCA cycle & ETS occur sequentially and orderly.
 - The NADH synthesised in glycolysis is transferred into mitochondria and undergoes oxidative phosphorylation.
 - Intermediates in the pathway are not used to synthesise other compounds.
 - Only glucose is being respired. Other alternative substrates are not entered in the pathway at any stages.
- Such assumptions are not valid because,
 - o All pathways work simultaneously and do not take place one after another.
 - o Substrates enter the pathways and are withdrawn from it as and when necessary.
 - o ATP is utilized as and when needed.
 - o Enzymatic rates are controlled by multiple means.
- Such calculations are useful to appreciate the efficiency of the living system in extraction and storing energy.

Net gain of ATP molecules from one glucose molecule

Glycolysis	2 ATP directly	2 ATP
	2 molecules of NADH	6 ATP
Oxidative decarboxylation	2 NADH	6 ATP
	6 NADH	18 ATP
TCA cycle	2 FADH	4 ATP
	2 GTP	2 ATP
	Total	38 ATP


2 ATP molecules are spent for transporting 2 NADH molecules formed during glycolysis to the mitochondria. Hence the **net gain = 36 ATP molecules**.

Comparison b/w fermentation & aerobic respiration

Fermentation	Aerobic respiration
Partial breakdown of glucose.	Complete breakdown of glucose to CO ₂ & H ₂ O.
Net gain of only 2 ATP.	Net gain of 36 ATP.
NADH is oxidised to NAD ⁺ rather slowly.	NADH is oxidised to NAD ⁺ very vigorously.

AMPHIBOLIC PATHWAY

- Glucose is the favoured substrate for respiration. So, all carbohydrates are first converted to glucose for respiration.
- Other substrates are also respired.



- Fats breakdown into glycerol & fatty acids. **Fatty acids** are degraded to **acetyl CoA** and enter the pathway. **Glycerol** is converted to **PGAL** and enters the pathway.
- Proteins are degraded by proteases into amino acids. Each amino acid (after deamination) enters the pathway at some stage in the Krebs' cycle or as pyruvate or acetyl CoA.

Visit: www.bankofbiology.com

- The respiratory pathway is generally considered as a catabolic pathway. But it involves both **anabolism** (synthesis) and **catabolism** (breakdown). So it is better called as an **amphibolic pathway**.

E.g. Fatty acids breakdown to acetyl CoA before entering the respiratory pathway. But when the organism needs to synthesise fatty acids, acetyl CoA withdraw from the respiratory pathway.

Similarly, during breakdown and synthesis of protein, respiratory intermediates are involved.

RESPIRATORY QUOTIENT (RQ) OR RESPIRATORY RATIO

- It is the ratio of the volume of CO₂ evolved to the volume of O₂ consumed in respiration.


$$RQ = \frac{\text{Volume of CO}_2 \text{ evolved}}{\text{Volume of O}_2 \text{ consumed}}$$

- RQ depends upon the type of respiratory substrate.
- **RQ for carbohydrates = 1**, because equal amounts of CO₂ and O₂ are evolved and consumed, respectively.



$$RQ = \frac{6 \text{ CO}_2}{6 \text{ O}_2} = 1.0$$

- **RQ for fats = < 1**. Calculations for a fatty acid, (e.g. tripalmitin) are shown:



$$RQ = \frac{102 \text{ CO}_2}{145 \text{ O}_2} = 0.7$$

- **RQ for proteins = 0.9**.
- In living organisms, respiratory substances are often more than one. Pure proteins or fats are never used as respiratory substrates.

15. PLANT GROWTH AND DEVELOPMENT



All plant cells are descendants of the zygote (fertilized egg). The zygote develops into a mature plant through growth and differentiation forming roots, leaves, branches, flowers, fruits and seeds. Then they eventually die.

GROWTH

- Growth is an irreversible permanent increase in size of an organ or its parts or an individual cell.
- It involves metabolic processes that consume energy.

Plant Growth Generally is Indeterminate

- Plant growth continues throughout the life due to the presence of **meristems**.
- Meristematic cells have capacity to divide & self-perpetuate.
- The growth where new cells are always added to the plant body by the meristem is called **open form of growth**.
- **Primary growth:** It occurs due to **root apical meristem & shoot apical meristem**. It causes the elongation of the plants along the axis.
- **Secondary growth** (In gymnosperms & dicots): It occurs due to **lateral meristems, vascular cambium & cork-cambium**. It causes increase in the girth of organs.





Growth Rates

- It is the increased growth per unit time.
- The growth rate may be **arithmetic or geometrical**.

Arithmetic growth:

- In this, following mitotic division, only one daughter cell continues to divide while the other differentiates & matures.
- On plotting the length of the organ against time, a linear curve is obtained.

Arithmetic growth



Mathematically, it is expressed as $L_t = L_0 + rt$



L_t = length at time 't'

L_0 = length at time 'zero'

r = growth rate / elongation per unit time.

Geometrical growth:

- Here, both daughter cells continue mitotic cell division.
- In most systems, the initial growth is **slow (lag phase)**, then it increases rapidly (**log or exponential phase**).
- If nutrient supply is limited, the growth slows down leading to a **stationary phase**.
- On plotting the parameter of growth against time, we get a typical **sigmoid (S) curve**.
- A sigmoid curve is a characteristic of living organism growing in a natural environment. It is typical for all cells, tissues and organs of a plant.




Growth is Measurable

- At cellular level, growth occurs due to increase in the amount of protoplasm.
- Increase in protoplasm is difficult to measure directly. So growth is measured by parameters like increase in fresh weight, dry weight, length, area, volume & cell number. E.g.
 - o **Cell number:** E.g. A maize root apical meristem can produce more than 17,500 new cells per hour.
 - o **Cell size:** E.g. Cells in a watermelon can increase in size by up to 3,50,000 times.
 - o **Length:** E.g. Growth of a pollen tube.
 - o **Surface area:** E.g. Growth in a dorsi-ventral leaf.

Phases of Growth

3 phases: **meristematic, elongation & maturation**.


- **Meristematic phase:** It occurs in the meristems at the root apex & the shoot apex. Here, cells have rich protoplasm and large nuclei. Cell walls are primary, thin & cellulosic with abundant plasmodesmata.
- **Elongation phase:** It occurs in cells proximal (just next, away from the tip) to the meristematic zone. The cells have increased vacuolation, size and new cell wall deposition.
- **Maturation phase:** It occurs in the cells further away from the apex, i.e., more proximal to the phase of elongation. The cells attain maximal size in terms of wall thickening and protoplasmic modifications.



Exponential growth is expressed as $W_1 = W_0 e^{rt}$

W_1 = final size (weight, height, number etc.)
 W_0 = initial size at the beginning of period
 r = growth rate (relative)
 t = time of growth
 e = base of natural logarithms

- Here, **r** is **relative growth rate**. It is also the measure of ability of plant to produce new plant material (**efficiency index**). Hence, final size W_1 depends on initial size, W_0 .
- Quantitative comparisons between the growth can also be made in 2 ways:
 - (i) **Absolute growth rate**: Measurement & comparison of total growth per unit time.
 - (ii) **Relative growth rate**: Measurement of growth of the given system per unit time expressed on a common basis, e.g., per unit initial parameter.



Diagrammatic comparison of absolute & relative growth rates

Conditions (essential elements) for Growth

- 1. Water:** Essential for cell enlargement. Turgidity of cells helps in extension growth. Water provides medium for enzymatic activities needed for growth.
- 2. Oxygen:** It helps to release metabolic energy for growth.
- 3. Nutrients:** Macro & micro elements are needed for the synthesis of protoplasm and act as source of energy.
- 4. Temperature:** At optimum temperature, growth is maximum. Deviation from this may harm the plants.
- 5. Light & gravity:** Affect certain phases/stages of growth.


DIFFERENTIATION, DEDIFFERENTIATION & REDIFFERENTIATION

- **Differentiation** is the process in which the cells in meristems (root apical & shoot-apical) and cambium differentiate and mature to perform specific functions.
- In this, cell walls & protoplasm undergo major structural changes. The capacity of cell division is lost. E.g. Loss of protoplasm to form a tracheary element. They also develop very strong, elastic, lignocellulosic secondary cell walls to carry water to long distances even under extreme tension.
- Under certain conditions, living differentiated cells regain the capacity of division. This is called **dedifferentiation**. E.g. formation of meristems (interfascicular cambium & cork cambium) from differentiated parenchyma cells.


- The dedifferentiated cells can divide and produce cells that again lose the capacity to divide but mature to perform specific functions. It is called **redifferentiation**.
- Plant growth is open, i.e., it can be indeterminate or determinate. Differentiation in plants is also open, because cells/tissues arising out of the same meristem have different structures at maturity.
- Final structure at maturity of cell/tissue is also determined by the location of the cell. E.g. cells positioned away from root apical meristems differentiate as root-cap cells, while those pushed to the periphery mature as epidermis.

DEVELOPMENT

- It is a process that includes all changes in the life cycle of an organism from seed germination to senescence.
- It is the sum of growth and differentiation.



- Plants follow different pathways in response to environment or phases of life to form different kinds of structures. This ability is called **plasticity**. E.g.
 - **Heterophyly due to phases of life:** E.g. In cotton, coriander and larkspur, the leaves of the juvenile plants and mature plants are different in shape.
 - **Heterophyly due to environment:** E.g. Difference in shapes of leaves produced in air and water (e.g. buttercup).



Factors controlling the development:

- **Intrinsic factors:** Include intracellular (genetic) or intercellular factors (such as plant growth regulators).
- **Extrinsic factors:** Include light, temperature, water, oxygen, nutrition, etc.

PLANT GROWTH REGULATORS (PLANT HORMONES OR PHYTOHORMONES)


Plant growth regulators (PGRs) are small, simple molecules that regulate plant growth.

Based on the functions, PGRs are 2 groups:

- **Plant growth promoters:** For growth promoting activities like cell division & enlargement, tropic growth, pattern formation, flowering, fruiting & seed formation. E.g. auxins, gibberellins and cytokinins.
- **Plant growth inhibitors:** For growth inhibiting activities like dormancy & abscission. Respond to wounds & stresses of biotic and abiotic origin. E.g. abscisic acid & ethylene. (Ethylene fits either of the groups, but it is largely a growth inhibitor).

1. Auxins

- **Charles Darwin** & his son **Francis Darwin** observed that the coleoptiles of canary grass responded to unilateral illumination by growing towards the light source (**phototropism**). It was concluded that the tip of coleoptile caused the bending of the entire coleoptile.




- **F.W. Went** isolated **Auxin** (Greek 'auxein': to grow) from tips of coleoptiles of oat seedlings.
- Auxin was first isolated from human urine.
- Auxins are produced by the growing apices of stems & roots, from where they migrate to regions of their action.

Types of Auxins:

- **Natural:** E.g. **Indole-3-acetic acid (IAA)** and **indole butyric acid (IBA)**. They are isolated from plants.
- **Synthetic:** E.g. **NAA (naphthalene acetic acid)** and **2, 4-D (2, 4-dichlorophenoxyacetic)**.

Functions of auxins:

- Initiate rooting in **stem cuttings** for plant propagation.
- Promote flowering. E.g. in pineapples.
- Prevent fruit and leaf drop at early stages.
- Promote the abscission of older leaves and fruits.
- Induce **parthenocarpy**. E.g., in tomatoes.
- Used as **herbicides**. E.g. 2, 4-D is used to kill dicot weeds. It does not affect mature monocot plants. It is used to prepare weed-free lawns.
- Controls xylem differentiation and helps in cell division.



In higher plants, the growing apical bud inhibits the growth of lateral (axillary) buds. It is known as **apical dominance**. Removal of shoot tips (decapitation) results in the growth of lateral buds. It is applied in tea plantations & hedge-making.

2. Gibberellins

- These are acidic PGR.
- **E. Kurosawa** treated the sterile filtrates of *Gibberella fujikuroi* (a fungus that causes 'bakane' disease or foolish seedling in rice) to healthy rice seedlings. As a result, it showed the symptoms of 'bakane' disease. Later, the active substances were identified as **gibberellic acid**.
- There are more than 100 gibberellins (GA₁, GA₂, GA₃ and so on) in fungi and higher plants.
- **Gibberellic acid (GA₃ or Terpenes)** is one of the first discovered and most intensively studied gibberellins.

Functions:

- They cause an increase in length of axis. So they are used to increase the length of grapes stalks.
- To elongate and improve the shape of fruits such as apple.
- They delay senescence. So the fruits can be left on the tree to extend the market period.
- GA₃ is used to speed up malting process in brewing industry.
- Sugarcane stores sugar in stems. Spraying sugarcane crop with gibberellins increases the length of the stem. It increases the yield by as much as 20 tonnes per acre.
- Spraying juvenile conifers with GAs hastens the maturity period. It leads to early seed production.
- For **bolting** (internode elongation just prior to flowering) in beet, cabbages and many plants with rosette habit.

3. Cytokinins

- **F. Skoog** and co-workers observed that from the internodal segments of tobacco stems, the callus (a mass of undifferentiated cells) proliferated only if the nutrients medium was supplemented with extracts of vascular tissues, yeast extract, coconut milk or DNA.

Skoog & Miller later identified and crystallized the active substance and termed as **kinetin**.

- Cytokinins were discovered as kinetin (N₆-furfurylamino purine - an Adenine derivative) from the autoclaved herring sperm DNA.
- Kinetin does not occur naturally in plants.
- **Zeatin** (from corn-kernels and coconut milk) is the natural substances with cytokinin-like activities.
- There are some synthetic compounds with cell division promoting activity.
- Natural cytokinins are synthesized in regions of rapid cell division (root apices, shoot buds, young fruits etc).

Functions:

- Play a role in cytokinesis.
- Help to produce new leaves, chloroplasts in leaves, lateral shoot growth and adventitious shoot formation.
- Help overcome the apical dominance.

- Promote nutrient mobilization which helps in the delay of leaf senescence.

4. Ethylene (C₂H₄)

- **Cousins** confirmed that ripened oranges released a volatile substance that hastened the ripening of stored bananas. Later this substance was identified as ethylene.
- Ethylene is a simple gaseous PGR.
- It is synthesized in large amounts by tissues undergoing senescence and ripening fruits.

Functions:

- Influences horizontal growth of seedlings, swelling of the axis and apical hook formation in dicot seedlings.
- Promotes senescence and abscission of plant organs especially of leaves and flowers.
- Promotes fruit ripening. It enhances respiration rate during fruit ripening. This is called **respiratory climactic**.
- Breaks seed and bud dormancy, initiates germination in peanut seeds, sprouting of potato tubers.
- Promotes rapid internode/petiole elongation in deep water rice plants. It helps leaves/upper parts of the shoot to remain above water.
- Promotes root growth and root hair formation. It increases absorption surface.
- Used to initiate flowering and for synchronising fruit-set in pineapples. It also induces flowering in mango.
- It is widely used in agriculture.

The most widely used source of ethylene is **ethephon**. Ethephon in an aqueous solution is readily absorbed and transported within the plant and releases ethylene slowly.

Ethephon hastens fruit ripening in tomatoes & apples and accelerates abscission in flowers and fruits (thinning of cotton, cherry, walnut). It promotes female flowers in cucumbers thereby increasing the yield.

5. Abscisic acid (ABA)

- During mid-1960s, it was reported 3 kinds of inhibitors: **inhibitor-B, abscisin II & dormin**. They were chemically identical and now known as **abscisic acid**.
- ABA is the derivatives of carotenoids.
- It regulates abscission and dormancy.

Functions:

- Inhibitor of plant growth and metabolism.
- Inhibits seed germination.
- Stimulates the closure of stomata in the epidermis.
- Increases the tolerance of plants to various kinds of stresses. Therefore, it is also called the **stress hormone**.
- For seed development, maturation & dormancy (it helps to withstand desiccation and other unfavourable factors).

Interactions of PGRs

- PGRs play individualistic or synergistic role. Such roles may be complimentary or antagonistic.
- PGRs interact to affect dormancy in seeds/ buds, abscission, flowering, senescence, vernalisation, apical dominance, seed germination, plant movements etc.
- In most situations, ABA acts as an antagonist to GAs.


Factors influencing the action of PGR:

- **Intrinsic factor:** Genomic control.
- **Extrinsic factors:** Light and Temperature.

ROLE OF LIGHT AND TEMPERATURE ON FLOWERING

1. PHOTOPERIODISM

- It is the response of plants to periods of day/night.
- Some plants require light to induce flowering.
- Based on light duration, plants are 3 groups:
 - **Long day plants:** They require the exposure to light for a period exceeding a well-defined critical duration.
 - **Short day plants:** They require the exposure to light for a period less than the critical duration before the flowering is initiated in them.
 - **Day-neutral plants:** They have no correlation between exposure to light duration and induction of flowering.



- While shoot apices modify into flowering apices, they by themselves cannot perceive photoperiods. The site of perception of light/dark duration is the leaves.

- It has been hypothesised that there is hormone(s) for flowering. When plants get enough photoperiod, the hormone migrates from leaves to shoot apices to induce flowering.

2. VERNALISATION

- It is the phenomenon in which some plants depend quantitatively or qualitatively on exposure to low temperature for flowering.
- It prevents precocious reproductive development late in the growing season, and enables the plant to have sufficient time to reach maturity.

Examples for vernalisation:

1. Some food plants, wheat, barley & rye have two varieties:
 - **Spring varieties:** These are normally planted in the spring and come to flower and produce grain before the end of the growing season.
 - **Winter varieties:** Winter varieties if planted in spring would normally fail to flower or produce mature grain within a span of a flowering season. Hence, they are planted in autumn. They germinate, and over winter come out as small seedlings, resume growth in the spring, and are harvested usually around mid-summer.
2. **Vernalisation in biennial plants:** Biennials are monocarpic plants that normally flower and die in second season. E.g. Sugar beet, cabbages, carrots etc. Subjecting

the growing of a biennial plant to a cold treatment stimulates a subsequent photoperiodic flowering response.

SEED DORMANCY

- Certain seeds fail to germinate even under favourable external conditions. Such seeds are in **dormancy**.
- Dormancy is caused by endogenous conditions within the seed. E.g. Hard seed coat; chemical inhibitors such as ABA, phenolic acids, para-ascorbic acid; and immature embryos.
- Dormancy can be overcome naturally and artificially. E.g.
 - **Breaking of seed coat barrier:** By mechanical abrasions using knives, sandpaper etc. or vigorous shaking. In nature, abrasions are caused by microbial action, and passage through digestive tract of animals.
 - **Removing inhibitory substances:** By subjecting the seeds to chilling conditions or by application of certain chemicals like gibberellic acid and nitrates.
 - Changing the environmental conditions, such as light and temperature.

17. BREATHING AND EXCHANGE OF GASES

Respiration is the oxidation of nutrients in the living cells to release energy for biological work.

Breathing is the exchange of O₂ from the atmosphere with CO₂ produced by the cells.

RESPIRATORY ORGANS


- **General body surface:** E.g. lower invertebrates (sponges,

coelenterates, flatworms etc).

- **Skin or moist cuticle (cutaneous respiration):** E.g. earthworms, leech, amphibians etc.
- **Tracheal tubes:** E.g. insects, centipede, millipede, spider.
- **Gills (Branchial respiration):** E.g. fishes, tadpoles, prawn.
- **Lungs (Pulmonary respiration):** E.g. most vertebrates.

HUMAN RESPIRATORY SYSTEM

It consists of a pair of *air passages (air tract)* and *lungs*.



1. Air passages

- **Conducting part** which transports the atmospheric air into the alveoli, clears it from foreign particles, humidifies and brings the air to body temperature.

External nostrils → nasal passage → nasal chamber (cavity) → pharynx → glottis → larynx → trachea → primary bronchi → secondary bronchi → tertiary bronchi → bronchioles → terminal bronchioles → respiratory bronchiole → alveolar duct.

- Each terminal bronchiole gives rise to many very thin and vascularised *alveoli* (in lungs).

- A cartilaginous *Larynx* (sound box or voice box) helps in sound production.
- During swallowing, *epiglottis* (a thin elastic cartilaginous flap) closes *glottis* to prevent entry of food into larynx.
- Trachea, all bronchi and initial bronchioles are supported by incomplete cartilaginous half rings.

2. Lungs

- Lungs situate in *thoracic chamber* and rest on *diaphragm*.
- Right lung has 3 lobes and left lung has 2 lobes.
- Lungs are covered by double-layered *pleura* (outer **parietal pleura** and inner **visceral pleura**).
- The *pleural fluid* present in between these 2 layers lubricates the surface of the lungs and prevents friction between the membranes.
- **Lungs = Bronchi + bronchioles + alveoli.**
- Alveoli and their ducts form the *respiratory or exchange part* of the respiratory system.
- *Alveoli are the structural and functional units of lungs.*


Steps of respiration

1. Pulmonary ventilation (breathing).
2. Gas exchange between lung alveoli & blood.
3. Gas transport (O₂ transport & CO₂ transport).
4. Gas exchange between blood & tissues.
5. Cellular or tissue respiration.

MECHANISM OF BREATHING (INSPIRATION & EXPIRATION)

a. Inspiration

- **Active** intake of air from atmosphere into lungs.
- During this, the **diaphragm contracts** (flattens) causing an increase in vertical thoracic volume (*antero-posterior axis*).
- Contraction of **external intercostal muscles** (muscles found between ribs) lifts up the ribs and sternum causing an increase in thoracic volume in the *dorso-ventral axis*.
- Increase in thoracic volume reduces thoracic pressure. So, lungs expand. Thus, pulmonary volume increases resulting in decrease of *intra-pulmonary pressure* to less than the atmospheric pressure. So, air moves into lungs.



b. Expiration

- **Passive** expelling of air from the lungs.
- During this, *intercostal muscles & diaphragm* relax causing a decrease in thoracic volume and thereby pulmonary volume. So, air moves out.
- During **forceful expiration**, **abdominal muscles** and **internal inter-costal muscles contract**.

Respiratory volumes and capacities

- **Tidal volume (TV):** Volume of air inspired or expired during a normal respiration. It is about **500 ml**. i.e., **6000-8000 ml** per minute.
- **Inspiratory reserve volume (IRV) or complementary air:** Additional volume of air that can inspire by forceful inspiration. It is **2500-3000 ml**.
- **Expiratory reserve volume (ERV) or supplemental air:** Additional volume of air that can expire by a forceful expiration. It is **1000-1100 ml**.
- **Residual volume (RV):** Volume of air remaining in lungs after a forcible expiration. It is **1100-1200 ml**.

- **Inspiratory capacity (IC):** Total volume of air inspired after a normal expiration (TV + IRV). It is **3000-3500 ml**.
- **Expiratory capacity (EC):** Total volume of air expired after a normal inspiration (TV + ERV). It is **1500-1600 ml**.
- **Functional residual capacity (FRC):** Volume of air remaining in the lungs after a normal expiration (ERV + RV). It is **2100-2300 ml**.
- **Vital capacity (VC):** Volume of air that can breathe in after a forced expiration or Volume of air that can breathe out after a forced inspiration (ERV + TV + IRV).

It is **3500-4500 ml**.

- **Total lung capacity (TLC):** Total volume of air in the lungs after a maximum inspiration. (RV + ERV + TV + IRV or VC + RV). It is **5000-6000 ml**.
- Part of respiratory tract (from nostrils to terminal bronchi) not involved in gaseous exchange is called **dead space**. **Dead air volume** is about **150 ml**.

- **Respiratory cycle**= an inspiration + an expiration
- **Normal respiratory (breathing) rate:** 12-16 times/min
- **Spirometer (respirometer):** To measure respiratory rate.

GAS EXCHANGE

Gas exchange occurs between

1. **Alveoli and blood**
2. **Blood and tissues**

Alveoli are the primary sites of gas exchange.

O₂ & CO₂ are exchanged by simple diffusion. It depends upon the following factors:

- **Pressure/ concentration gradient:** The **Partial pressures** (individual pressure of a gas in a gas mixture) of O₂ and CO₂ (pO₂ and pCO₂) are given below.

Respiratory gas	pO ₂ (in mm Hg)	pCO ₂ (in mm Hg)
Atmospheric air	159	0.3
Alveoli	104	40
Deoxygenated blood	40	45
Oxygenated blood	95	40
Tissues	40	45

pO₂ in alveoli is more (**104 mm Hg**) than that in **blood capillaries (40 mm Hg)**. So O₂ diffuses into capillary blood. pCO₂ in **deoxygenated blood** is more (**45 mm Hg**) than that in alveoli (**40 mm Hg**). So, CO₂ diffuses to alveoli.


- **Solubility of gases:** Solubility of CO₂ is 20-25 times higher than that of O₂. So, the amount of CO₂ that can diffuse through the diffusion membrane per unit difference in partial pressure is higher than that of O₂.

- **Thickness of membranes:** The diffusion membrane is made up of 3 layers:

- a) **Squamous epithelium** of alveoli.
- b) **Endothelium** of alveolar capillaries.
- c) **Basement substance** between them.

Its total thickness is only 0.5 μm. It enables easy gas exchange.

- **Surface area:** Presence of alveoli increases the surface area of lungs. It increases the gas exchange.



GAS TRANSPORT (O₂ TRANSPORT & CO₂ TRANSPORT)


It is the transport of respiratory gases (O₂ & CO₂) from alveoli to the systemic tissues and vice versa.

1. O₂ TRANSPORT

It is the transport of O₂ from lungs to various tissues.

It occurs in 2 ways:

- a. **In physical solution (blood plasma):** About **3%** of O₂ is carried in a dissolved state through plasma.
- b. **As oxyhaemoglobin:** About **97%** of O₂ is transported by **haemoglobin** (red coloured iron containing pigment) on RBC. O₂ binds with haemoglobin (Hb) to form **oxyhaemoglobin**. This is called **oxygenation**. Hb has **4 haem units**. So, each Hb molecule can carry 4 oxygen molecules. Binding of O₂ depends upon pO₂, pCO₂, H⁺ ion concentration (pH) and temperature.




- In the alveoli, high pO₂, low pCO₂, lesser H⁺ ion concentration and lower temperature exist. These factors are favourable for the formation of oxyhaemoglobin.

- In tissues, low pO₂, high pCO₂, high H⁺ ions and high temperature exist. So Hb₄O₈ dissociates to release O₂.
- Every 100 ml of oxygenated blood can deliver around 5 ml of O₂ to the tissues under normal physiological conditions.

Oxygen-haemoglobin dissociation curve

It is a sigmoid curve obtained when percentage saturation of Hb with O₂ is plotted against the pO₂. It is used to study the effect of factors like pCO₂, H⁺ concentration etc., on binding of O₂ with Hb.



2. CO₂ TRANSPORT

It is the transport of CO₂ from tissues to lungs.

In tissues, pCO₂ is high due to catabolism and pO₂ is low. In lungs, pCO₂ is low and pO₂ is high. This favours CO₂ transport from tissues to lungs. It occurs in 3 ways:

- a. **As carbonic acid:** In tissues, **7%** of CO₂ is dissolved in **plasma water** to form **carbonic acid** and carried to lungs.
- b. **As carbamino-haemoglobin:** In tissues, **20-25%** of CO₂ binds to Hb to form **carbamino-haemoglobin**. In alveoli, CO₂ dissociates from carbamino-haemoglobin.
- c. **As bicarbonates:** **70%** of CO₂ transported by this method. RBCs contain an enzyme, *carbonic anhydrase*. (It is slightly present in plasma too).

At tissue site, it facilitates the following reactions:



In alveoli, the above reaction proceeds in opposite direction leading to the formation of CO₂ and H₂O.

Every **100 ml of deoxygenated blood** delivers about **4 ml of CO₂** to the alveoli.

REGULATION OF RESPIRATION

In brain, there are the following **Respiratory centres**:

- **Respiratory rhythm centre (Inspiratory & Expiratory centres):** In **medulla oblongata**. It regulates respiratory rhythms.
- **Pneumotaxic centre:** In **Pons**. It moderates functions of respiratory rhythm centre. Impulse from this centre reduces the duration of inspiration and thereby alter respiratory rate.

- **Chemosensitive area:** Seen adjacent to the rhythm centre. Increase in the concentration of CO₂ and H⁺ activates this centre, which in turn signals rhythm centre. **Receptors in aortic arch & carotid artery** also recognize changes in CO₂ & H⁺ concentration and send signals to rhythm centre. Role of oxygen in the regulation of respiratory rhythm is quite insignificant.

DISORDERS OF RESPIRATORY SYSTEM

1. **Asthma:** Difficulty in breathing causing wheezing due to inflammation of bronchi and bronchioles.
2. **Emphysema:** Damage of alveolar walls. It decreases respiratory surface. Major cause is cigarette smoking.
3. **Occupational respiratory disorders:** Certain industries produce so much dust. So, the defense mechanism of the body cannot cope with the situation. Long exposure causes inflammation leading to **fibrosis** (proliferation of fibrous tissues). It results in lung damage. Workers in such industries should wear protective masks.


Visit: www.bankofbiology.com

MODEL QUESTIONS

1. Draw a flowchart showing the different parts of the air tract.
2. Match the following


A	B	C
IC	TV + ERV	3500-4500 ml
EC	ERV + TV + IRV	2100-2300 ml
FRC	TV + IRV	3000-3500 ml
VC	ERV + RV	1500-1600 ml

3. Note the relationship between first two words and fill up the fourth place.
 - a. TV: 500 ml IRV:..... b. Atmospheric air: 159 mm Hg Alveoli:
4. The given diagram shows the exchange of gases between alveolus and alveolar capillary.



- a. Identify X and Y.
- b. Name the Physical Process involved in gas exchange.
- c. Mention the factors that favour this process.

5. The given graph shows oxygen-haemoglobin dissociation curve.



- a. What is the nature of curve?
- b. Find out the pressure at which Haemoglobin is 50% saturated with O₂?
- c. What are the factors which influence it?

6. Identify the two true statements from the statements given below and rewrite the two false statements correctly.
 - a. Pneumonia is a chronic disorder due to cigarette smoking.
 - b. Carbon dioxide combines with haemoglobin to form carbamino haemoglobin.
 - c. Respiratory rhythm is maintained by the respiratory centre in the heart.
 - d. Alveoli are the primary sites of exchange of gases.

18. BODY FLUIDS AND CIRCULATION

Circulation is the transport of nutrients, oxygen, CO₂ and excretory products to the concerned tissues or organs. For circulation, simple organisms (**sponges, coelenterates etc.**) use water from their surroundings. Complex organisms use body fluids (**blood & lymph**) for circulation.

CIRCULATORY PATHWAYS

Circulatory system is 2 types- **Open** and **Closed**.


- **Open circulatory system:** Here, the blood pumped by the heart passes through large vessels into open spaces or cavities called **sinuses**. E.g. Arthropods and molluscs.
- **Closed circulatory system:** Here, the blood pumped by the heart is always circulated through blood vessels. This is more advantageous as the flow of fluid can be more precisely regulated. E.g. Annelids and chordates.

All vertebrates have a muscular chambered heart.

- **Fishes:** 2-chambered heart (an atrium + a ventricle).
- **Amphibians:** 3-chambered heart (2 atria + a ventricle).
- **Reptiles (except crocodiles):** 3-chambered heart (2 atria + a ventricle). Ventricle is incompletely partitioned.
- **Crocodiles, birds & mammals:** 4-chambered heart.

Types of circulation

- **Single circulation:** In fishes. In this, heart receives impure blood only (*venous heart*).
Deoxygenated blood → to heart → to gills → oxygenated blood → to body parts → deoxygenated blood → to heart.
- **Incomplete double circulation:** In amphibians & reptiles. In this, left atrium gets oxygenated blood from gills/lungs/skin and right atrium gets deoxygenated blood from other body parts. However, they get mixed up in the single ventricle. It pumps out mixed blood.
- **Double circulation:** In birds & mammals. Right atrium gets deoxygenated blood and passes to right ventricle and left atrium gets oxygenated blood and passes to left ventricle. The ventricles pump it out separately without any mixing up.



BLOOD VASCULAR SYSTEM

It includes Heart, Blood & Blood vessels.

1. BLOOD

Formed of **plasma** (55%) & **formed elements** (45%).

A. PLASMA

Straw-coloured, slightly alkaline (pH 7.4) viscous fluid.

Constituents of plasma

- **Water (90-92%):** It is a good solvent.
- **Plasma proteins (6-8 %):** Include
 - **Fibrinogen:** For blood coagulation.
 - **Globulins:** Act as antibodies (for defense of the body).
 - **Albumins:** For osmotic balance. Regulate blood pressure.
- **Glucose, amino acids, lipids & cholesterol.**
- **Inorganic constituents:** Na⁺, Ca²⁺, Mg²⁺, Cl⁻, HCO₃⁻ etc.
- **Gases** like CO₂, O₂, N₂ etc.

Plasma without clotting factors is known as **Serum**.

B. FORMED ELEMENTS (RBC, WBC & PLATELETS)

Red Blood Cells (RBC) or Erythrocytes:

- Biconcave non-nucleated cells. No mitochondria, Golgi complex etc. Red colour is due to **Haemoglobin** (iron containing protein). Normal Hb level is 12-16 g/ 100 ml.
- **Count:** 5 - 5.5 millions/ mm³.
- **Formed in:** Red Bone marrow.
- **Average lifespan:** 120 days. Worn-out RBCs are destroyed in **spleen** (graveyard of RBCs).
- **Function:** CO₂ and O₂ transports.

White Blood Cells (WBC) or Leucocytes:

- Colourless nucleated cells.
- **Count:** 6000-8000 /mm³.
- **Formed in:** Bone marrow, lymph glands, spleen.
- **Average lifespan:** Generally short lived (1- 15 days).
- **Function:** Part of immune system.

Types of WBC: Granulocytes & Agranulocytes

1. Granulocytes

They are 3 types:

- Neutrophils (Heterophils):** 60-65%. Soldier of the body.
Function: Phagocytosis.
- Eosinophils (Acidophils):** 2-3%. Resist infections. Cause allergic reactions.
- Basophils (Cyanophils):** 0.5-1%. Secrete histamine, serotonin, heparin etc. Cause inflammatory reactions.


2. Agranulocytes

They are 2 types:

- Lymphocytes (20-25%):** Smallest WBC with largest nucleus. Includes **B- lymphocytes & T- lymphocytes**. Cause immune responses. Secrete antibodies.
- Monocytes (6-8%):** Largest WBC.
Function: Phagocytosis.

Platelets (Thrombocytes):

- Colourless non-nucleated cell fragments.
- **Count:** 1.5 - 3.5 lakhs/mm³.
- **Formed in:** Megakaryocytes in Bone marrow.
- **Average lifespan:** 7 days.
- **Function:** Blood clotting.



BLOOD COAGULATION

It is a mechanism for **haemostasis** (prevention of blood loss through injuries). At the site of injury, following events occur:

Clumped **platelets & tissues** release **thromboplastin** → It forms **thrombokinase (Prothrombinase)** enzyme → **Thrombokinase** hydrolyses **prothrombin** to **thrombin** enzyme in presence of **Ca²⁺** → **Thrombin** converts soluble **fibrinogen** to insoluble **fibrin** → **Fibrin** threads trap dead & damaged blood cells to form **clot (coagulum)**.

BLOOD GROUPS

Blood groups were discovered by **Carl Land Steiner**.

1. ABO grouping

It is based on presence or absence of 2 surface **antigens** (chemicals that induce immune response) on **RBCs** namely **A & B**. Similarly, **plasma** contains 2 **antibodies** (proteins produced in response to antigens) namely **anti-A & anti-B**.

Blood group	Antigens	Antibodies	Can donate blood to	Can receive blood from (Donor's group)
A	A	Anti-B	A & AB	A, O
B	B	Anti-A	B & AB	B, O
AB	A, B	Nil	AB only	A, B, AB & O
O	Nil	Anti-A & Anti-B	A, B, AB & O	O only

- Antigen A reacts with anti-A. Antigen B reacts with anti-B.
- If bloods with interactive antigens & antibodies are mixed together, it causes **clumping (agglutination)** of RBCs.
- Persons with **O Group** are called **Universal donors** because they can donate blood to persons with any other blood group. Persons with **AB group** are called **Universal recipients** because they can accept blood from all groups.

2. Rh grouping

- **Rhesus (Rh) factor** is another antigen found on RBC.
- **Rh+ve** means the presence of Rh factor and **Rh-ve** means absence of Rh factor. Nearly **80%** of humans are Rh+ve.
- **Anti-Rh antibodies** are not naturally found. So Rh-ve person can receive Rh+ve blood only once but it causes the development of anti-Rh antibodies in his blood. So, a second transfusion of Rh+ve blood causes **agglutination**. Therefore, Rh-group should be matched before transfusion.

Erythroblastosis foetalis

- It is a **Rh incompatibility** between the Rh-ve blood of a pregnant mother and Rh+ve blood of the foetus.
- Rh antigens do not get mixed with maternal blood in first pregnancy because placenta separates the two bloods.
- But during first delivery, the maternal blood may be exposed to small amount of foetal blood (Rh+ve). This induces the formation of Rh antibodies in maternal blood.
- In case of her subsequent pregnancies, the Rh antibodies from the mother leak into the foetal blood (Rh+ve) and destroy the foetal RBCs. This is fatal to the foetus or cause severe **anaemia** and **jaundice** to the baby. This condition is called **Erythroblastosis foetalis**.
- It can be avoided by administering **anti-Rh antibodies** to the mother immediately after the first delivery.

2. BLOOD VESSELS


Blood vessels are 3 types: Arteries, Veins & Capillaries.

- **Arteries:** They carry blood from heart to other tissues. They contain oxygenated blood (except pulmonary artery). Their smaller branches are called **arterioles**. Arteries are 3-layered- inner **tunica intima** (squamous endothelium), middle **tunica media** (smooth muscles & elastic fibres) and outer **tunica externa** (fibrous connective tissue).
- **Veins:** They carry blood towards heart. They contain deoxygenated blood (except pulmonary vein). Their smaller branches are called **venules**. Veins are also 3-layered but tunica media is comparatively thin.
- **Capillaries:** In tissues, arterioles divide into thin walled and single layered vessels. They are called capillaries. They unite into venules.

3. HEART

- It is a mesodermally derived organ located in **mediastinum**.
- It has the size of a clenched fist.
- It is protected by double-layered **pericardium**.
- The **pericardial space** (between pericardial membranes) is filled with **pericardial fluid**. It reduces the friction between the heart walls and surrounding tissues.
- Heart has 4 chambers- two upper **atria (auricles)** and two lower **ventricles**.

- The walls (*cardiac muscles*) of the *ventricles* are much **thicker** than that of the atria.



- The atria are separated by an *inter-atrial septum* and the ventricles are separated by *inter-ventricular septum*.
- In between atrium and ventricle, there is a thick fibrous *atrio-ventricular septum* with an opening.
- A *tricuspid valve* (3 muscular flaps or cusps) guards the opening between right atrium & right ventricle. A *bicuspid (mitral) valve* guards the opening between left atrium and left ventricle. These valves allow the flow of blood only in one direction, i.e. from atria to ventricles.

- Right ventricle has an opening to *pulmonary artery* and left ventricle has an opening to *aorta*. These openings have *semi-lunar valves*. They prevent backward flow of blood.

CONDUCTING SYSTEM OF HEART

- It includes **nodal tissues, bundles & fibres**.
- *Nodal tissues* are specialized cardiac musculature present in heart wall. They are 2 types:
 - o *Sino-atrial node (SAN)* in the right upper corner of the right atrium.
 - o *Atrio-ventricular node (AVN)* in the lower left corner of the right atrium close to the *atrio-ventricular septum*.
- From the AVN, a bundle of fibrous *atrio-ventricular bundle (AV bundle)* passes through *atrio-ventricular septa* and divides into right & left branches. Each branch passes through the ventricular walls of its side. In the ventricular wall, it breaks up into minute fibres (*Purkinje fibres*). These fibres along with the bundles are known as *bundle of His*.
- *Nodal tissues* generate **action potential** without any external stimuli, i.e. it is **autoexcitable**. SAN initiates and maintains contraction of heart by generating action potentials (**70-75/min**). So, it is called the *pacemaker*.

CARDIAC CYCLE

It is the cyclic contraction and relaxation of heart for pumping blood. It involves 3 stages:

- 1. Joint diastole:** It is the relaxed state of all chambers of heart. When the **tricuspid** and **bicuspid valves** open, blood from **pulmonary vein** and **vena cava** flows into **left & right ventricles** respectively through **left and right atria**. **Semilunar valves** are **closed** at this stage.
- 2. Atrial (Auricular) systole:** SAN generates an **action potential**. As a result, both the atria contract. It is called *atrial systole*. This increases the flow of blood into the ventricles by about **30%**.
- 3. Ventricular systole:** The action potential is conducted to ventricular side by AVN & AV bundle from where **bundle of His** transmits it through the *ventricular musculature*. As a result, ventricles contract. It is called *ventricular systole*. During this, the atria undergo diastole. Ventricular systole increases the **ventricular pressure** causing
 - * Closure of **tricuspid** and **bicuspid valves** due to attempted backflow of blood into the atria.
 - * Semilunar valves open. So deoxygenated blood enters the *pulmonary artery* from **right ventricle** and oxygenated blood enters the *aorta* from **left ventricle**.

The ventricles now relax (*ventricular diastole*) and the **ventricular pressure** falls causing

- * The closure of the *semilunar valves* which prevents the backflow of blood into the ventricles.
- * The *tricuspid* and *bicuspid valves* are opened by the pressure in the atria.

The ventricles and atria again undergo joint diastole and the above processes are repeated.


A cardiac cycle is completed in **0.8 seconds**.

- **One heartbeat = a cardiac cycle.**
So, normal heartbeat: **70-75 times/min (average: 72/min)**.
- **Stroke volume:** It is the volume of blood pumped out by each ventricle during a cardiac cycle. It is about **70 ml**.
- **Cardiac output:** It is the volume of blood pumped out by each ventricle per minute, i.e. **stroke volume x heart rate (70 x 72)**. It is about **5000 ml (5 litres)**.
Cardiac output of an athlete is very high.
- **Heart sounds:** During each cardiac cycle, 2 sounds are produced. The first sound (**lub**) is due to the closure of *tricuspid* and *bicuspid valves*. The second sound (**dub**) is due to the closure of the *semilunar valves*.

One heartbeat = a lub + a dub.

ELECTROCARDIOGRAPH (ECG)

- It is an instrument used to obtain *electrocardiogram*.
- Electrocardiogram is the *graphical representation* of the *electrical activity* of the heart during a cardiac cycle.
- To get an ECG, a patient is connected to the machine with 3 *electrical leads* (one to each wrist and to left ankle) that monitor heart activity. For a detailed evaluation of heart's function, multiple leads are attached to the chest region.
- An ECG consists of the following waves:
 - o **P-wave:** Represents the excitation (*depolarization*) of atria during *atrial systole*.
 - o **QRS-complex:** Represents *depolarization* of ventricles during *Ventricular systole*.
 - o **T-wave:** Represents the *repolarisation* of ventricles.



Deviation in the ECG indicates the abnormality or disease. So, ECG has great clinical significance.

DOUBLE CIRCULATION


It is the circulation in which blood flows through the heart twice for completing its circuit.

It includes:

- 1. Pulmonary circulation:** Circulation b/w lungs and heart.
Deoxygenated blood from right ventricle → to **pulmonary artery** → to **lungs** → *oxygenated blood* → to **pulmonary veins** → left atrium.
- 2. Systemic circulation:** Circulation b/w heart and various body parts.
Oxygenated blood from left ventricle → to **aorta** → **arteries** → **arterioles** → **capillaries** → **tissues** → *deoxygenated blood* from tissues → **venules** → **veins** → **vena cava** → to **right atrium**.

Systemic circulation provides nutrients, O₂ and other substances to the tissues and takes CO₂ and other harmful substances away for elimination.

- **Hepatic portal system:** It is a system which includes the *hepatic portal vein* that carries blood from *intestine* to the *liver* before it is delivered to the systemic circulation.
- **Coronary circulatory system:** It is a system of *coronary vessels* that circulate blood to and from *cardiac musculature*.



REGULATION OF CARDIAC ACTIVITY

- Normal activities of heart are auto-regulated by *nodal tissues*. So, it is called **myogenic heart**.
- *Medulla oblongata* regulates cardiac activity through ANS.
- *Sympathetic nerves* of ANS increase the rate of heartbeat, the strength of ventricular contraction and cardiac output.
- *Parasympathetic nerves* of ANS decrease the heartbeat, conduction of action potential and the cardiac output.
- *Adrenal medullary hormones* increase the cardiac output.

LYMPHATIC SYSTEM

- Includes *Lymph, Lymph vessels & Lymph nodes (glands)*.
- As the blood passes through the capillaries in tissues, some water and soluble substances are filtered out from plasma to the intercellular spaces, to form **tissue (interstitial) fluid**. It has same mineral distribution as that in plasma.
- Some tissue fluid enters **lymphatic system** and the tissue fluid in them is called **lymph**. It drains back to major veins.
- Lymph is a colourless fluid containing lymphocytes.

Functions of lymph

- It is the middleman between blood & tissues. Tissue fluid helps to exchange nutrients, gases, etc. b/w blood and cells.
- It carries plasma proteins synthesized in liver to the blood.
- Transports digested fats (through lacteals in the intestinal villi), fat soluble vitamins, hormones etc.
- Filtration of bacteria and foreign particles.
- Lymph nodes produce WBC (lymphocytes) & antibodies.

DISORDERS OF CIRCULATORY SYSTEM

- **Hypertension (High Blood Pressure):** The pressure of circulating blood on the walls of blood vessels is called **blood pressure**. Normal BP is **120/80 mm Hg**. It includes *systolic (pumping) pressure* (120 mm Hg) and *diastolic (resting) pressure* (80 mm Hg).
When the BP is higher than normal, it is called **hypertension**. If an individual repeatedly has the BP of **140/90 or above**, it shows *hypertension*. It leads to *heart diseases* and affects *vital organs* (brain, kidney etc).
- **Coronary Artery Disease (CAD) or Atherosclerosis:** Here, **Ca, fat, cholesterol** and **fibrous tissue** are deposited

in **coronary arteries**. So the lumen of arteries becomes narrower and thereby affects the blood supply.

- **Angina (angina pectoris):** An *acute chest pain* due to *O₂ deficiency* to heart muscles. It occurs due to improper blood flow. It is common among middle-aged and elderly.
- **Heart Failure (congestive heart failure):** It is the inability of heart to pump blood enough to meet the needs of the body. Congestion of the lungs is the main symptom.
- **Cardiac arrest:** Heart stops beating.
- **Heart attack:** Sudden damage of heart muscle due to inadequate blood supply.

Visit: www.bankofbiology.com

MODEL QUESTIONS

1. Complete the table.

Plasma proteins	Albumin	Osmotic balance
	Globulin	(f)
(a)	(b)	Blood clotting
	(c)	Antibody production
	Monocyte	(g)
Granulocytes	(d)	Phagocytosis
	Basophil	(h)
	(e)	Allergic reaction

2. If a person having blood group A is given blood transfusion of blood group B by mistake, what will be its effect?

3. Match the following

A	B	C
Lymph	Stroke volume x heart rate	No antigen
Cardiac output	Tissue fluid	Left ventricle
O group	Aorta	5 litres
Semi-lunar valve	Universal donor	Middle man


4. Rhythmic heartbeat is maintained by a highly specialized excitatory and conductive System. The correct sequence of events will be

- A V node → Bundle of His → S A node → Purkinje fibers
- Purkinje fibers → A V node → S A node → Bundle of His
- A V node → S A node → Bundle of His → Purkinje fibers
- S A node → A V node → Bundle of His → Purkinje fibers

5. Give reason for the following


- Ventricular walls are thicker than the atrial wall.
- Human heart continues to beat even after it is separated from the body for transplantation.

6. Observe the diagram below.



- Name the type of circulation.
- Label A, B, C & D.
- Mention the Significance of this circulation.

7. Observe the diagram given below



- Draw and label P, Q, R and S. What is its clinical significance?
- Why do we call our heart myogenic?

8. Stethoscope is an instrument which is used to detect the sounds of the heart.

- Mention the two sounds of the heart.
- Give the causes of heart sound.

9. The blood pressure of a person is shown as 170/130 mm Hg. What would be his disease? How it affects his body?

19. EXCRETORY PRODUCTS & THEIR ELIMINATION

Excretion is the elimination of metabolic wastes like **ammonia, urea, uric acid** etc. from the tissues.

Types of excretion

- Ammonotelism:** Process of excretion of NH_3 .
Ammonotelic animals: Aquatic invertebrates, aquatic insects, bony fishes, aquatic amphibians etc.
 NH_3 is highly toxic. So, excretion needs excess of water. NH_3 is readily soluble in water and is excreted by diffusion through body surface or gill surfaces (in fishes) as ammonium ions.
 Kidneys do not play any significant role in its removal.
- Ureotelism:** Process of excretion of **urea**.
Ureotelic animals: Cartilaginous fishes, terrestrial & semi-aquatic amphibians (frogs, toads etc.), aquatic & semi-aquatic reptiles (alligators, turtles), mammals etc.
 In liver, NH_3 is converted into less toxic urea. So, it needs only moderate quantity of water for excretion.

Some amount of urea may be retained in the kidney matrix of some animals to maintain a desired osmolarity.

- Uricotelism:** Process of excretion of **uric acid**. It is water insoluble & less toxic. So, water is not needed for excretion.
Uricotelic animals: Insects, some land crustaceans, land snails, terrestrial reptiles & birds.

Ureotelism & uricotelism are needed for water conservation.

Some excretory organs in animals



- Protonephridia (flame cells):** In Flatworms, rotifers, some annelids & cephalochordate (*Amphioxus*). Protonephridia are primarily for osmoregulation.
- Nephridia:** In Annelids. Help in the removal of nitrogenous wastes and osmoregulation.
- Malpighian tubules:** In Insects. Help in the removal of nitrogenous wastes and osmoregulation.
- Antennal or green glands:** In Crustaceans (prawn etc.)
- Kidneys:** In higher animals.

HUMAN EXCRETORY SYSTEM


It includes **kidneys, ureters, urinary bladder & urethra**.

Structure of Kidney

- Reddish brown, bean-shaped structures situated between the levels of last thoracic & 3rd lumbar vertebra.
- Length: **10-12 cm**, width: **5-7 cm**, thickness: **2-3 cm**. Average weight: **120-170 gm**.
- It is enclosed in a tough, 3-layered **fibrous renal capsule**.
- On the concave side of kidney, there is an opening (**hilum** or **hilus**) through which blood vessels, nerves, lymphatic ducts and ureter enter the kidney.
- Hilum leads to funnel shaped cavity called **renal pelvis** with projections called **calyces**.
- A kidney has outer **cortex** & inner **medulla**.
- Medulla has few conical projections called **medullary pyramids (renal pyramids)** projecting into the calyces.
- Cortex extends in between the medullary pyramids as renal columns (**Columns of Bertini**).
- Each kidney has nearly one million tubular **nephrons**.



- o **Glomerulus:** A tuft of capillaries formed by **afferent arteriole** (a fine branch of renal artery). Blood from glomerulus is carried away by **efferent arteriole**.



- o **Renal tubule:** It begins with a double walled cup-like **Bowman's capsule**, which encloses the glomerulus.

Glomerulus + Bowman's capsule = Malpighian body

- The tubule continues with **proximal convoluted tubule (PCT)**, **Henle's loop** & **distal convoluted tubule (DCT)**.
- Henle's loop is hairpin-shaped. It has **descending and ascending limbs**.
- The DCTs of many nephrons open into a **collecting duct**. Collecting duct extends from cortex to inner parts of medulla. They converge and open into the **renal pelvis** through **medullary pyramids** in the **calyces**.
- **Malpighian body (Renal corpuscle)**, **PCT** and **DCT** are situated in **renal cortex**. **Loop of Henle** dips into **medulla**.
- The **efferent arteriole** forms a fine capillary network (**peritubular capillaries**) around the renal tubule. A minute vessel of this network runs parallel to Henle's loop forming a 'U' shaped **vasa recta**.

Types of nephrons

- Cortical nephrons (85%):** In this, the Henle's loop is short and extends only very little into the medulla. Vasa recta is absent or highly reduced.
- Juxtamedullary nephrons (15%):** In this, Henle's loop is long and runs deep into medulla. Vasa recta present.

Nephron


- Nephrons are the structural & functional units of kidney.
- Each nephron has 2 parts: **Glomerulus & Renal tubule**.

URINE FORMATION (PHYSIOLOGY OF KIDNEY)

3 processes: Glomerular filtration, reabsorption & secretion.

1. Glomerular filtration (ultrafiltration)

- The glomerular capillary blood pressure causes filtration of blood through 3 layers, i.e. **endothelium of glomerular blood vessels, epithelium of Bowman's capsule & a basement membrane** between these 2 layers.
- The epithelial cells (**podocytes**) of the Bowman's capsule are arranged in an intricate manner leaving some minute spaces called **filtration slits (slit pores)**.
- Almost all constituents of the blood plasma except the proteins pass onto the lumen of the Bowman's capsule.
- About **1100-1200 ml of blood** is filtered by the kidneys per minute. It constitutes **1/5th** of the blood pumped out by each ventricle of the heart in a minute.
- The amount of glomerular filtrate formed per minute is called **Glomerular filtration rate (GFR)**.
- **Normal GFR = 125 ml/minute, i.e., 180 litres/day.**



2. Reabsorption

- **180 litres** of glomerular filtrate is produced daily. But about **99%** of this is reabsorbed by the renal tubules. So normal volume of urine released is **1.5 litres**.
- From the filtrate, **glucose, amino acids, Na⁺**, etc. are reabsorbed **actively** and **nitrogenous wastes** are absorbed **passively**. Passive reabsorption of water occurs in the initial segments of the nephron.
- **PCT** reabsorbs most of the nutrients, and 70-80% of electrolytes & water. Simple cuboidal brush border epithelium of PCT increases surface area for reabsorption.
- **Loop of Henle** maintains high osmolarity of medullary interstitial fluid. **Descending limb** is permeable to water but almost impermeable to electrolytes. This concentrates the filtrate. In **ascending limb**, minimum reabsorption occurs. It is impermeable to water but allows transport of electrolytes.

So, filtrate gets diluted.


- In **DCT**, conditional reabsorption of **Na⁺** & water takes place.
- **Collecting duct** extends from cortex to inner parts of medulla. It reabsorbs large amount of water to concentrate urine. It also allows passage of small amounts of urea into medullary interstitium to keep up the osmolarity.

3. Tubular Secretion

- Cells of **PCT & DCT** maintain **ionic (Na-K balance)** and **acid-base balance (pH)** of body fluids by selective secretion of **H⁺, K⁺ & NH₃** into the filtrate and absorption of **HCO₃⁻** from it.
- **Collecting duct** also maintains pH and ionic balance of blood by the secretion of **H⁺** and **K⁺** ions.

Mechanism of concentration of the filtrate

- **Henle's loop & vasa recta** help to concentrate the urine.
- The flow of **filtrate** in the 2 **limbs of Henle's loop** and the flow of **blood** through the 2 **limbs of vasa recta** are in opposite directions (**counter current pattern**). This is called **Counter current mechanism**.
- Due to the counter current and proximity between Henle's loop & vasa recta, **osmolarity** increases from cortex (**300 mOsmol⁻¹**) to the **inner medullary interstitium (1200 mOsmol⁻¹)**. This gradient is caused by **NaCl & urea**.
- NaCl is transported by ascending limb of Henle's loop that is exchanged with descending limb of vasa recta. NaCl is returned to interstitium by ascending limb of vasa recta. Similarly, small amount of urea enters the thin segment of the ascending limb of Henle's loop which is transported back to the interstitium by the collecting tubule. Thus electrolytes and urea are retained in the interstitium and maintain a **concentration gradient (interstitial gradient)** in medullary interstitium. It enables easy passage of water from collecting tubule to concentrate the filtrate (urine).
- Thus DCT & collecting duct produce urine **four times concentrated** than the initial filtrate formed (i.e. 300 mOsmol⁻¹ to 1200 mOsmol⁻¹).



MICTURITION

- Gradual filling of urinary bladder causes stretching. As a result, **stretch receptors** on its wall send impulses to CNS. The CNS passes on **motor messages**. It causes the contraction of **smooth muscles** of the bladder and simultaneous relaxation of the **urethral sphincter**. It results in **micturition** (release of urine).
- The neural mechanism causing micturition is called **micturition reflex**.
- An adult human excretes **1 to 1.5 litres** of urine (**25-30 gm urea**) per day.
- Urine is a **light yellow coloured watery fluid**, slightly **acidic (pH-6.0)** and has a characteristic odour.
- Various conditions affect the characteristics of urine.
- Analysis of urine helps in **clinical diagnosis** of many metabolic disorders and malfunctioning of the kidney.

E.g. **Glycosuria** (presence of glucose) and **Ketonuria** (ketone bodies) in urine indicates **diabetes mellitus**.

Role of Lungs, liver & skin in Excretion

- ♦ **Lungs:** Remove **CO₂ (200 mL/minute)** and **water**.
- ♦ **Liver:** Secretes bile containing **bilirubin, biliverdin, cholesterol, degraded steroid hormones, vitamins and drugs**. Most of them pass out along with digestive wastes.
- ♦ **Skin (Sweat glands & sebaceous glands):** Sweat contains water, NaCl, small amounts of urea, lactic acid, etc. Primary function of sweat is to give a **cooling effect** on body surface. **Sebaceous glands** eliminate **sterols, hydrocarbons, waxes etc.** through **sebum**. Sebum provides a protective oily covering for the skin.
- ♦ **Saliva** eliminates small amounts of nitrogenous wastes.

REGULATION OF THE KIDNEY FUNCTION

- It is done by hormonal feedback mechanisms involving the **hypothalamus, JGA** and the **heart**.
- Changes in **blood volume, body fluid volume** and **ionic concentration** activate **Osmoreceptors** in the body.

1. Regulation by ADH (vasopressin)

- When body fluid level decreases, the **osmoreceptors** stimulate **hypothalamus** to release **antidiuretic hormone (ADH)**. It stimulates water reabsorption from **DCT & collecting duct**. Thus, ADH prevents **diuresis** and increases body fluid volume.
- Increase in fluid volume switches off the osmoreceptors and suppresses ADH release to complete the feedback.
- ADH constricts blood vessels resulting in an increase of BP. This increases the glomerular blood flow and GFR.

2. Regulation by JGA (Renin-Angiotensin mechanism)

- **JGA (Juxta glomerular apparatus)** is a sensitive region formed by cellular modification of **DCT** and the **afferent**

arteriole at the location of their contact.

- JGA regulates the **GFR**.
- A fall in glomerular blood flow/glomerular blood pressure/GFR activates the **JG cells** to release **renin**.
- Renin converts **angiotensinogen** in blood to **angiotensin I** and further to **angiotensin II** (a **vasoconstrictor**).
- Angiotensin II performs the following functions:
 - ❖ Increases glomerular blood pressure and thereby GFR.
 - ❖ Activates **adrenal cortex** to release **Aldosterone**.
- Aldosterone causes **reabsorption** of **Na⁺** and **water** from the **distal parts** of the tubule. This also leads to an increase in blood pressure and GFR.

3. Regulation by ANF

- ANF check on the renin- angiotensin mechanism.
- An increase in blood flow to the atria of the heart causes the release of **Atrial Natriuretic Factor (ANF)**.
- ANF causes **vasodilation** (dilation of blood vessels) and thereby decreases the blood pressure.

DISORDERS OF EXCRETORY SYSTEM

- **Uremia:** Accumulation of urea in blood due to malfunction of kidney. It may lead to **kidney failure (renal failure)**.
- **Renal calculi:** Stone or insoluble mass of crystallized salts (oxalates, etc.) formed within the kidney.
- **Glomerulonephritis:** Inflammation of glomeruli.

Hemodialysis

- It is a process of removal of **urea** in patients with uremia.
- The **dialyzing unit** (artificial kidney) contains a coiled **cellophane tube** surrounded by **dialyzing fluid**. It has same composition of plasma except nitrogenous wastes.
- Blood drained from a convenient artery is pumped into **dialyzing unit** after adding anticoagulant like **heparin**.

- The porous **cellophane membrane** of the tube allows the passage of molecules based on concentration gradient.
- As nitrogenous wastes are absent in dialyzing fluid, these substances freely move out, thereby clearing the blood.
- The purified blood is pumped back to the body through a vein after adding **anti-heparin** to it.

Kidney transplantation


- It is the ultimate method in the correction of **acute renal failures**. A functioning kidney is taken from a donor.
- It is better to receive kidney from a close relative to minimize chances of rejection by immune system of host.

MODEL QUESTIONS


1. Terrestrial animals are generally either ureotelic or uricotelic, not ammonotelic. Why?
2. Note the relationship between first two words and fill up the fourth place
 - a) Bony fishes: Ammonotelism Birds:.....
 - b) JG cells: Renin Atria:.....
3. Complete the following sentences
 - a) Reabsorption of water from DCT is facilitated by the hormone.....
 - b) Angiotensin II activates the adrenal cortex to release.....
 - c) In cases of Kidney failure, urea can be removed by the process called.....
4. Match the following

A	B	B
Malpighian body	Urethral sphincter	Oxalates
Uraemia	Glomerulus	Release of urine
Renal calculi	Accumulation of urea	Afferent and efferent arterioles
Micturition	Crystallized salts	Kidney failure

5. Prepare a flowchart of filtrate flow in the nephron using the flow terms.
Collecting duct, PCT, DCT, Ascending limb of Henle's loop, descending limb of Henle's loop, Bowman's capsule
6. "Counter current system plays an important role in concentrating urine." Name any two regions inside the kidney, where the counter current system is seen.
7. Complete the flowchart given below:
(Hint: Angiotensin II, JG cells, Constricts blood vessels, Angiotensinogen, Aldosterone)



8. Observe the diagram



- a) In which season ADH production is higher?
- b) Why the production of ADH varies in different seasons?

20. LOCOMOTION AND MOVEMENT

Locomotion is the voluntary movements resulting in a change in location. All locomotion are movements but all movements are not locomotion. Both are interlinked. E.g.

- In *Paramecium*, cilia help in the movement of food through cytopharynx and in locomotion.
- *Hydra* use tentacles to capture prey and for locomotion.
- Limbs help to change body postures and for locomotion.

Types of movement in human being

- **Amoeboid movement:** By **pseudopodia** formed by streaming of protoplasm as in *Amoeba*. Cytoskeletal

elements like microfilaments also help for this. E.g. Macrophages & leucocytes.

- **Ciliary movement:** By **cilia**. E.g. ciliary movements in trachea (to remove dust particles and foreign substances), and oviducts (for the passage of ova).
- **Muscular movement:** By muscles. E.g. movement of limbs.

Flagellar movement helps in the swimming of spermatozoa, maintenance of water current in the canal system of sponges and in locomotion of Protozoans like *Euglena*.


HUMAN MUSCULAR SYSTEM

- It includes muscles which are mesodermal in origin.
- Muscles constitute 40-50% of the body weight.
- Muscles have excitability, contractility, extensibility & elasticity.
- Based on location, muscles are 3 types:


Skeletal (striated) muscles	Visceral (Non-striated) muscles	Cardiac muscles
Attached to skeleton	In visceral organs	In heart wall
Striations present	Absent	Present
Voluntary	Involuntary	Involuntary
Rich blood supply	Poor blood supply	Rich blood supply
Fatigue muscle	Non-fatigue	Non-fatigue
Multinucleate	Uninucleate	Uninucleate
More mitochondria	Less mitochondria	More mitochondria

STRUCTURE OF STRIATED MUSCLE

- Skeletal muscle is made of **muscle bundles (fascicles)** held together by collagenous connective tissue layer (**fascia**).




- Each fascicle contains many **muscle fibres (muscle cells)**.
- Muscle fibres are lined by **plasma membrane (sarcolemma)** enclosing the **sarcoplasm**.
- Each muscle fibre contains **myofilaments (myofibrils)**.
- Each myofibril has alternate **dark (Anisotropic or A-band)** and **light striations (Isotropic or I-band)**. This is due to the presence of 2 fibrous contractile proteins- thin **Actin filament** and thick **Myosin filament**.
- I-bands contain actin. A-bands contain actin and myosin. They are arranged parallel to each other.
- A-band bears a lighter middle region (**H band**) formed of only myosin. A thin dark line (**M-line**) runs through the centre of **H-zone**.
- I-band is bisected by a dense dark band called **Z-line**. Region between two Z-lines is called **sarcomere**. They are the **functional units of muscle contraction**.




Structure of contractile proteins

- An **actin filament** is made of **2 filamentous (F) actins** which form double helix.
- F-actin is a polymer of monomeric **Globular (G) actins**.



- Actin contains 2 other proteins (**tropomyosin & troponin**).
- Two filaments of **tropomyosin** run along the grooves of the F-actin double helix.
- **Troponin** has 3 subunits. It is seen at regular intervals on tropomyosin. In the resting state, a **subunit of troponin** masks the binding sites for myosin on the actin filaments.
- Each myosin filament is a polymer of many **monomeric proteins** called **Meromyosins**.



- A meromyosin has 2 parts:
 - o **Heavy meromyosin or HMM or cross arm (globular head + short arm):** It projects outwards.
 - o **Light meromyosin or LMM (tail).**
- The globular head is an active **ATPase enzyme** and has **binding sites for ATP** and **active sites for actin**.

MECHANISM OF MUSCLE CONTRACTION

According to **sliding filament theory**, contraction of a muscle fibre occurs by the sliding of thin filaments over thick filaments.


The steps are given below:

- An **impulse** from the **CNS** reaches the **neuromuscular junction (Motor-end plate)** via **motor neuron**.

Neuromuscular junction is the synapse between a motor neuron and the sarcolemma of the muscle fibre.


A motor neuron + muscle fibres = a motor unit.

- Synaptic vesicles release a neurotransmitter **Acetylcholine**. It generates an **action potential** in the sarcolemma that spreads through the muscle fibre. It causes the release of **Ca²⁺ ions** from sarcoplasmic cisternae into **sarcoplasm**.
- Ca** binds with a subunit of **troponin** on actin filaments and unmask the active sites for myosin.



- Using energy from **ATP hydrolysis**, **myosin head** binds to **active sites** on the actin to form **cross bridge**. This pulls actin filaments on both sides towards the centre of **A-band**. Actin filaments partially overlap so that **H-zone** disappears.

- The **Z- line** attached to actins is also pulled inwards. It causes a shortening (**contraction**) of **sarcomere**.
- I-bands** get shortened, whereas **A-bands** retain the length.
- Myosin releases **ADP** and **Pi** and goes back to its relaxed state. A new ATP binds and the cross-bridge is broken.
- The ATP is again hydrolyzed by the myosin head and the above processes are repeated causing further sliding.
- When **Ca²⁺** ions are pumped back to sarcoplasmic cisternae, actin filaments are again masked. As a result, **Z-lines** return to their original position. It results in **relaxation**.



- The reaction time of the fibres varies in different muscles.
- Repeated activation of muscles leads to the accumulation of the **lactic acid** causing **muscle fatigue**. This is due to **anaerobic breakdown** of **glycogen** in muscles.

Red muscle fibres and white muscle fibres

Red (Aerobic) muscles	White muscle
Red coloured due to myoglobin	White coloured due to lesser myoglobin
More mitochondria	Less mitochondria
Aerobic metabolism	Anaerobic metabolism
Slow & sustained contraction	Fast contraction for short period

HUMAN SKELETAL SYSTEM


It consists of a framework of **bones (206)** & few **cartilages**. Human skeletal system has 2 parts: **axial & appendicular**.

1. Axial skeletal system (80 bones)

Includes **bones of head, vertebral column, sternum & ribs**.

a. Bones of Head (29 bones)

It includes skull, Hyoid and Ear ossicles.




- Skull (22):** Include cranial bones and facial bones.
 - Cranial bones (8):** Include *Frontal (1), Parietals (2), Temporals (2), Occipital (1), Sphenoid (1) & Ethmoid (1)*.
 - Facial bones (14):** Include *Nasals (2), Maxillae (2), Zygomatics (2), Lacrimals (2), Palatines (2), Inferior nasals (2), Mandible (1) and Vomer (1)*.

Skull articulates with **First vertebra (atlas)** with the help of **2 occipital condyles (dicondylic skull)**.

- Hyoid bone (1):** U-shaped bone seen below buccal cavity.
- Ear ossicles (3 x 2 = 6):** *Malleus (2), Incus (2) & stapes (2)*.

b. Vertebral column


- Formed of **26 vertebrae**. Includes *Cervical vertebrae (7), Thoracic vertebrae (12), Lumbar vertebrae (5), Sacral vertebrae (1-fused) and Coccygeal vertebrae (1-fused)*.
- Vertebra has a central hollow portion (**neural canal**) through which the **spinal cord** passes.
- Number of cervical vertebrae are 7 in almost all mammals.
- The **vertebral column** protects the spinal cord, supports the head and serves as the point of attachment for the ribs and musculature of the back.



c. Sternum or Breast bone (1)

- Flat bone on the ventral midline of thorax.

d. Ribs (12 pairs)



- **True ribs** (first 7 pairs): They are attached to **thoracic vertebrae** and ventrally connected to sternum with the help of **Hyaline cartilage**.
- **Vertebrochondral (false) ribs** (8th, 9th & 10th pairs): They do not articulate directly with the sternum but join the **7th rib** with the help of **Hyaline cartilage**.
- **Floating ribs** (11th & 12th pairs): They are not connected ventrally (no connection with sternum or other ribs).
- Each rib has 2 articulation surfaces on its dorsal end and is hence called **bicephalic**.

b. Bones of hind-limbs (30 x 2 = 60)

Include *Femur* (thigh bone- 1), *Patella* (knee cap- 1), *Tibia* (1) & *fibula* (1), *Tarsals* (ankle bones-7), *Metatarsals* (5) & *Phalanges* (digits-14).

c. Pectoral girdles (2x2=4)

- Include **clavicle** (2) & **scapula** (2).
- Scapula is a large *triangular* flat bone situated in the dorsal part of the thorax between **the second and 7th ribs**.
- **Scapula (shoulder blade)** has a slightly elevated ridge (*spine*) which projects as a flat, expanded process (*acromion*). The **clavicle (collarbone)** articulates with this.
- Below the acromion is **glenoid cavity** which articulates with the head of **humerus** to form the **shoulder joint**.

d. Pelvic girdles (2)

- Formed of 2 coxal bones. Each coxal bone is formed by the fusion of 3 bones- *Ilium*, *Ischium* & *pubis*.
- At the point of fusion of **Ilium**, **Ischium** and **Pubis** is a cavity (**Acetabulum**) to which the **thigh bone** articulates.
- The 2 halves of the **pelvic girdle** meet ventrally to form **pubic symphysis** containing **fibrous cartilage**.

JOINTS


Joints are points of contact between bones, or between bones and cartilages. 3 types:

1. **Fibrous (immovable) joints:** E.g. sutures b/w skull bones.
2. **Cartilaginous joints (Slightly movable joints):** Bones are joined together with the help of cartilages. E.g. Joints between the adjacent vertebrae.
3. **Synovial (movable) joints:** They have a fluid filled synovial cavity between articulating surfaces of 2 bones.

Types of synovial joint

Joint	Examples
Ball & socket	Shoulder joint & hip joints.
Hinge joint	Knee joint, elbow joint, phalanges joints
Pivot joint	Joints b/w atlas & axis.
Gliding joint	Joints b/w carpals
Saddle joint	Joints b/w carpal & metacarpal of thumb

2. Appendicular skeletal system (126 bones)



a. Bones of fore-limbs (30 x 2 = 60)

Include *Humerus* (1), *Radius* (1), *Ulna* (1), *Carpals* (wrist bones- 8), *Metacarpals* (palm bones-5) & *Phalanges* (digits-14).

DISORDERS OF MUSCULAR & SKELETAL SYSTEMS

- **Myasthenia gravis:** An auto immune disorder that affects neuromuscular junction. It leads to fatigue, weakening and paralysis of skeletal muscles.
- **Muscular dystrophy:** Progressive degeneration of skeletal muscles. Mostly due to genetic disorder.
- **Tetany:** Rapid muscle spasm due to low Ca^{2+} in body fluid.

- **Arthritis:** Inflammation of joints.
- **Osteoporosis:** Age-related disorder characterized by decreased bone mass and increased chances of fractures. Decreased level of estrogen is a common cause.
- **Gout:** Inflammation of joints due to accumulation of uric acid crystals.

Visit: www.bankofbiology.com

MODEL QUESTIONS


1. When a muscle cell is viewed under a microscope, it has the following characters

Striations present, Multi nucleate, Sarcolemma present

- a) Identify the tissue b) Mention the function of the tissue

2. Observe the relaxed unit of a muscle given below.

- a) Label A, B & C.
 b) Redraw the diagram when the muscle unit is maximally contracted.
 c) Repeated activation of the muscle can lead to fatigue. Justify.




3. Arrange the following flow chart in the correct sequence of events:

Generation of action potential in the sarcolemma → Release of Ca^{2+} ions from sarcoplasmic cisternae into sarcoplasm → Active sites of actin are exposed → Impulse from the CNS → Neuromuscular junction → Shortening (contraction) of sarcomere → Synaptic vesicles release Acetylcholine → Actin filaments are pulled towards H-zone → Ca binds with troponin on actin filaments → Myosin head binds to active sites to form cross bridge → H-zone disappears.


4. Suppose a person is suffering from calcium deficiency for a prolonged time. How does it affect muscular contraction?

5. Diagram of a joint is given below:



- a) Identify the joint.
 b) Name three major structural forms of joints.

6. Complete the following chart




7. Match the following

Type of joints	Examples
i. Ball and socket	a) Joints of skull bone
ii. Pivot joint	b) Between carpals and metacarpals of the thumb
iii. Saddle joint	c) Between humerus and pectoral girdle
iv. Gliding joint	d) Between atlas and axis
	e) Between the carpals

21. NEURAL CONTROL AND CO-ORDINATION

- Neural (Nervous) system is a system that controls and coordinates the body activities, conducts and integrates the information and responds to stimuli.
- It includes **brain, spinal cord** and **nerves**.
- It is made up of specialized cells known as **neurons**.



Neuron is the **structural** and **functional unit** of neural system. It is composed of

- **Cell body (cyton):** Contains cytoplasm, cell organelles and **Nissl's granules** (granular bodies).
- **Dendron:** Short fibres projecting from the cyton. Their sub branches (**dendrites**) transmit impulses towards the cyton.

- **Axon:** A long fibre which transmit impulses away from the cell body. The branching of axon is called **axonite**. Each axonite ends as a bulb-like structure called **synaptic knob**.

Types of Neurons

- **Unipolar:** One axon. No Dendron. Found in embryo.
- **Bipolar:** One axon and one dendron. Found in the retina.
- **Multipolar:** One axon and 2 or more dendrons. Most common type. Found in the CNS & PNS.

Types of axon


- **Myelinated axon:** It is enveloped with **Schwann cells** that form a **myelin sheath** around the axon. Found in **spinal & cranial nerves**. The white coloured area, formed of myelinated nerve fibres is called **white matter**. Gaps b/w 2 adjacent myelin sheaths are called **nodes of Ranvier**.
- **Non-myelinated axon:** Schwann cells present but no myelin sheath. The gray coloured area without myelin sheath is called **gray matter**. Found in **autonomous & somatic neural systems**.

GENERATION & CONDUCTION OF NERVE IMPULSES

Impulse transmission is **electrochemical**. It has 3 steps:

1. Maintenance of resting membrane potential

- Neural membrane contains various selectively permeable ion channels.
- In a resting neuron (neuron not conducting impulse), the axonal membrane is more permeable to K^+ ions and nearly impermeable to Na^+ ions. Also, the membrane is impermeable to negatively charged proteins in axoplasm.
- Therefore, concentration of K^+ and -vely charged proteins in axoplasm is high and concentration of Na^+ is low.
- The fluid outside the axon contains low concentration of K^+ and high concentration of Na^+ . This forms an ionic or concentration gradient across resting membrane.
- The ionic gradients are maintained by the active transport of ions by the **Na-K pump**. It transports **3 Na^+** outwards for **2 K^+** into the cell. As a result, the outer surface becomes positively charged and inner surface becomes negatively charged (i.e, polarized).
- The electrical potential difference across the resting plasma membrane is called as the **resting potential**.



2. Action Potential

- When a stimulus is applied, the membrane at the site A becomes permeable to Na^+ . This causes rapid influx of Na^+ and reversal of the polarity at that site (outer negative and inner positive). It is called **depolarization**.

- The electrical potential difference during depolarization across the plasma membrane is called **action potential** (a **nerve impulse**).

3. Propagation of action potential

- At sites ahead (site B), outer surface is positive and inner surface is negative. As a result, a current flows on the inner surface from site A to site B.
- On the outer surface, current flows from site B to site A to complete the circuit. Hence, the polarity is reversed and action potential is generated at site B. i.e., action potential at site A arrives at site B.
- The sequence is repeated along the axon and the impulse is conducted.
- The rise in permeability to Na^+ is extremely short lived. It is quickly followed by a rise in permeability to K^+ .
- Immediately, K^+ diffuses outside the membrane and restores the resting membrane. Thus the fibre becomes ready for further stimulation.

Synaptic transmission of impulses


- Synapse is a functional junction between two neurons.
- It is 2 types: **Electrical & Chemical**.

1. Electrical synapses

- In this, the membranes of pre- and post-synaptic neurons are in close proximity. So impulse transmission is similar to the transmission along an axon.
- Impulse transmission is faster than in chemical synapse.
- Electrical synapses are very rare in human system.

2. Chemical synapses

- In this, there is a fluid filled space (**synaptic cleft**) between the presynaptic neuron and postsynaptic neuron.
- The presynaptic regions have swellings called **Synaptic knob (buttons)**. They contain **synaptic vesicles** filled with **neurotransmitters (acetylcholine or adrenaline)**.



Impulse transmission through chemical synapse:

Impulse reaches at axon terminal → synaptic vesicles bind on plasma membrane → release of neurotransmitter → It diffuses across synaptic cleft → combine with receptors on the post synaptic membrane → opening of ion channels allowing entry of ions → generates action potential.

- This action potential may be excitatory or inhibitory.

HUMAN NERVOUS (NEURAL) SYSTEM


It has 2 parts:

- **Central neural system (CNS):** Brain & spinal cord.
- **Peripheral nervous system (PNS):** All nerves.

CENTRAL NEURAL SYSTEM (CNS)

A. BRAIN

- It is protected in *cranial cavity*.
- It has 3-layered connective tissue membranes called *cranial meninges*.
- Meninges consist of outer *dura mater*, middle *arachnoid mater* and inner *pia mater*.
- The *subarachnoid space* (space between pia mater and arachnoid mater) is filled with **cerebrospinal fluid (CSF)**. The **ventricles** of brain are also filled with CSF.
- Brain has 3 divisions: Forebrain, Midbrain & Hindbrain.



a. Forebrain (Prosencephalon)

It is the anterior part. Consists of *cerebrum & diencephalon*.

Cerebrum

- Largest part. It has 2 **cerebral hemispheres** held together by a tract of nerve fibres (**Corpus callosum**).
- Outer part of cerebrum is called **cerebral cortex**. It has convulsions & depressions and is formed of **gray matter**. Gray colour is due to the presence of neuron cell bodies.
- Inner part of cerebrum is formed of **white matter**.
- Cerebral cortex consists of
 - **Motor area:** Controls voluntary movements of muscles.
 - **Sensory (Somaesthetic) area:** Controls the functioning of sense organs.
 - **Association area:** It is neither clearly sensory nor motor in function. Responsible for intersensory associations, memory and communication.

Integrated activities of different centres of cerebral cortex control intelligence, memory, judgment, learning, thinking and articulate speech.

Diencephalon (Thalamus & Hypothalamus)

- **Thalamus:** It is the structure around which the cerebrum wraps. It is a coordinating centre (relay station) for sensory and motor impulses.
- **Hypothalamus:** Seen below the thalamus. It
 - Regulates temperature, thirst, hunger and emotions.
 - Secretes hypothalamic hormones.
 - Controls pituitary gland.
 - Controls sleep, wakefulness, blood pressure, heart rate.
- The inner parts of cerebral hemispheres and a group of associated deep structures like *amygdala*, *hippocampus*, *hypothalamus*, etc. together constitute **Limbic system (Limbic lobe)**. It regulates sexual behavior, motivations, emotions (excitement, pleasure, rage, fear etc).

b. Midbrain (Mesencephalon)

- It is located between *thalamus/hypothalamus* and *Pons*.
- A canal (**cerebral aqueduct**) passes through the mid brain.
- Mid brain consists of 4 round lobes called **Corpora quadrigemina**. Their anterior pair is the centre of *visual reflexes* and the posterior pair is a centre of *auditory reflex*.

c. Hindbrain (Rhombencephalon)

It consists of **cerebellum, Pons & Medulla oblongata**. Midbrain & hindbrain form the **Brain stem**.

- **Cerebellum ("little cerebrum"):** It has very convoluted surface to accommodate more neurons. It co-ordinates muscular activities and body equilibrium.
- **Pons varoli:** It consists of fibre tracts that interconnect different regions of the brain. It co-ordinates the activities of eye and ear and regulates respiration.
- **Medulla oblongata:** It is connected to spinal cord. It controls respiration, cardiovascular reflexes, gastric secretions, peristalsis etc. It also controls salivation, vomiting, sneezing & coughing.

B. SPINAL CORD

- It is enclosed within the spinal canal of vertebral column.
- It is also protected by meninges.
- Spinal cord has a central canal containing CSF.
- Outer white matter and inner gray matter.

Functions:

- Conduction of impulses to and from the brain.
- Centre of spinal reflexes.

PERIPHERAL NEURAL SYSTEM (PNS)

It includes **cranial nerves** and **spinal nerves**.

Nerve fibres of PNS are 2 types:

- **Afferent (sensory) fibres:** Carry impulses from sense organs to CNS.
 - **Efferent (motor) fibres:** Carry impulses from CNS to muscles and glands.
- PNS has 2 divisions. They are
- **Somatic neural system:** Relays impulses from the CNS to skeletal muscles.
 - **Autonomic neural system (ANS):** Transmits impulses from CNS to involuntary organs & smooth muscles. It includes *sympathetic* & *parasympathetic* nerves. Sympathetic system prepares body to cope with emergencies, stresses & dangers. It increases heartbeat, breathing rate, constricts arteries and elevates BP. Parasympathetic system returns the body to a resting state after stressful situations and slows down heartbeat, dilates arteries, lowers BP etc.

Visceral nervous system is the part of PNS. It includes **nerves, fibres, ganglia & plexus** by which impulses travel from CNS to the viscera and from viscera to CNS.

REFLEX ACTION


It is the **rapid, involuntary** and **unconscious actions** of body in response to a stimulus. E.g.

- ♦ Withdrawal of the hand when it touches a hot object.
- ♦ Touching lips of a nursing baby evokes sucking reflex.

- ♦ Closing of the eyelids when light falls on them.
- ♦ Knee jerk phenomenon.
- ♦ If a child sees or smells a food unknown to him, he does not salivate. But if he sees or smells that food every time before tasting it, he salivates (**conditioned reflex**).

The pathway of impulses in a reflex action is called **Reflex arc**. It consists of

- A **receptor organ**: It receives the stimulus.
- **Sensory (afferent) neuron**: It transmits impulses from sense organ to CNS.
- **Intermediate (connector) neuron**: It connects sensory and motor neurons.
- **Motor (efferent/effector/excitor) neuron**: It conducts impulse from the CNS to effector organ.
- An **effector organ** (muscle/gland): It responds to impulse.




SENSORY RECEPTION & PROCESSING (SENSE ORGANS)

- These are the organs that detect the changes in the environment and convey the information to the CNS.
- It includes **eye, ear, nose, tongue & skin**.

EYE

- Two eyes are located in sockets of the skull called **orbits**.
- The adult human eyeball is nearly spherical.
- Eyeball has three layers: **Sclera, Choroid & Retina**.



a. Sclera

- The external layer formed of a dense connective tissue.
- Anterior transparent portion of sclera is called **cornea**.

b. Choroid

- Bluish middle layer. Contains many blood vessels.
- Choroid is thin over posterior two-thirds of the eyeball, but it is thick in the anterior part to form **ciliary body**.
- Ciliary body continues forward to form a visible pigmented and opaque portion of the eye called the **iris**.
- Iris has a central opening called **pupil**. The diameter of the pupil is regulated by the muscle fibres of iris. This helps to regulate the amount of light entering the eye.
- The eyeball contains a transparent crystalline **lens**. It is held in place by **ligaments** attached to the ciliary body.

c. Retina

- Inner layer. It contains 3 layers of cells – from inner to outer – **ganglion cells, bipolar cells & photoreceptor cells**.
- Photoreceptor cells are 2 types: **rods** and **cones**. They contain **photosensitive proteins (photopigments)**.
- Photopigments are formed of **opsin** (a protein) and **retinal** (an aldehyde of vitamin A).

Cone cells:

- **Function: Daylight (photopic) vision & colour vision.**
- There are 3 types of cones containing photopigments (**photopsin**) that respond to red, green and blue lights.
- The sensations of different colours are produced by combinations of these cones and their photopigments.
- When the cones are stimulated equally, a sensation of white light is produced.

Rod cells:

- **Function: Twilight (scotopic) vision.**
- They contain a purplish-red protein called **rhodopsin** (visual purple). It contains a derivative of **Vitamin A**.
- At the region, slightly above the posterior pole of the eyeball, **optic nerves** leave the eye and retinal blood vessels enter it. Here, photoreceptor cells are absent. It is called **blind spot**.
- Lateral to the blind spot, there is a yellowish pigmented spot called **macula lutea** with a central pit (**fovea**).
- The fovea is a thinned-out portion of the retina where only the cones are densely packed. It is the point of greatest visual acuity (resolution).
- The space between the cornea and lens is called **aqueous chamber**. It contains **aqueous humor** (thin watery fluid).


- The space between the lens and retina is called **vitreous chamber**. It contains **vitreous humor** (a transparent gel).

Mechanism of vision

Light reflected from an object → enters the eye through cornea & lens → focus on retina → dissociation of **retinal** from **opsin** → changes in **membrane permeability** → generates potential differences (impulse) in photoreceptor cells → generates action potentials in ganglion cells through bipolar cells → impulses are transmitted by **optic nerves** to brain (**visual cortex**) → impulses are analyzed and the image is recognized based on memory and experience → vision.

EAR (STATO-ACOUSTIC ORGAN)

- It is the organ for hearing & balancing.
- It has 3 divisions: **External ear, middle ear & inner ear**.



External ear

- Consists of **pinna (ear lobe)** & **auditory meatus (ear canal)**.
- At the opening of ear canal, hairs are seen.
- Ear canal and skin of pinna contains **ceruminous glands** (modified sweat glands). They secrete **wax (cerumen)**.
- Wax and hairs protects ears from foreign objects.
- Ear canal ends in **tympanic membrane (Tympanum or ear drum)**. It is a semi-transparent membrane covered by a thin layer of skin on its outer surface and by mucous membrane on the inside.

Middle ear

- Consists of **tympanic cavity** and **ear ossicles**.
- Tympanic cavity is an air filled space that separates the external and inner ear portions.
- An **auditory tube (Eustachian canal)** connects middle ear to the pharynx. It maintains an equal pressure on either side of the eardrum.
- **Ear ossicles** include 3 small bones namely **Malleus, Incus** and **stapes**. Malleus is attached to tympanum.
- **Stapes** is the **smallest bone** of the body. It is attached to membrane of **oval window (fenestra ovalis)** of inner ear.


Inner ear

- It consists of **bony labyrinth** & **membranous labyrinth**.
- Bony labyrinth is a cavity filled with **perilymph**.
- The membranous labyrinth consists of **cochlea** and **Vestibular apparatus**.

Cochlea (organ of hearing):

- It is a coiled structure having 3 canals - upper **scala vestibula**, middle **scala media** and lower **scala tympani**.
- Scala vestibula & scala media are separated by **Reissner's membrane**.

- Scala media and scala tympani are separated by **basilar membrane**.
- S. vestibula & S. tympani are filled with **perilymph** and scala media is filled with **endolymph**.
- Resting on the basilar membrane and projecting into scala media is complex receptor organ called **Organ of Corti**. It consists of row of **sensory hair cells**. The hairs (stereo cilia) of these cells project upwards and lie in contact with **tectorial membrane**, which projects above them.



Vestibular apparatus:

- It consists of 3 **semicircular canals** and **otolith organ**.
- 2 semicircular canals are vertical and one is horizontal. One end of each canal has a bulging called **ampulla**. Inside it is a lump called **crista ampullaris**. Long cilia of cells of crista are grouped together in a bundle (**cupula**).
- **Otolith organ** consists of **utricle** and **sacculle**.
- Utricle & Sacculle have a projecting ridge called **macula**.
- **Crista** and **Macula** are specific receptors in vestibular apparatus. They contain **sensory hair cells**. They are responsible **equilibrium & posture** of body.

Mechanism of hearing

Pinna collects sound waves → waves reach the **tympanic membrane** via ear canal → tympanic membrane vibrates → vibrations transmit to **ear ossicles** & **oval window** → **perilymph** in the **vestibular canal** vibrates → vibrations reach the **scala tympani** and force the **basilar membrane** to vibrate → hair endings of **sensory hair cells** press against **tectorial membrane** → sensory hair cells are excited → **auditory nerve** carries impulses to **auditory centre** of the brain → hearing.

NOSE

- Organ of **smell (olfaction)**.
- It contains mucus-coated receptors (**olfactory receptors**) made up of **olfactory epithelium**. They receive sense of smell. It contains 3 kinds of cells.
- The neurons of olfactory epithelium extend from the outside environment directly into a pair of broad bean-sized organs, called **olfactory bulb**. These are extensions of the brain's limbic system.

TONGUE

- Organ of **taste (gustation)**.
- 4 primary tastes are **sweet, salt, sour** and **bitter**.

- **Taste buds (Gustatoreceptors + supporting cells)** are seen around the bases of **taste papillae**.

taste buds and a complex flavour is perceived.


Nose & tongue are **chemoreceptors** (detect dissolved chemicals). Senses of taste & smell are functionally similar and interrelated. The brain integrates different input from

SKIN (Cutaneous receptors)

- Largest sense organ.
- It contains receptors for **heat, cold, touch, pain & pressure**.

MODEL QUESTIONS

1. Analyze the concept map given below and fill the gaps appropriately.



2. Note the relationship between first two words and fill up the fourth place

- a) Cone cells: Iodopsin Rod cells:
- b) Unipolar: No Dendron Bipolar:

3. Odd man out. Justify your answer.

- a) Tympanic membrane, Basilar membrane, Arachnoid membrane, Reissner's membrane
- b) Amygdala, Crista, Ampulla, Vestibular apparatus


4. Arrange the following processes in nerve impulse conduction in a sequential order.

- a) Bursting of synaptic vesicle b) Development of action potential
- c) Na⁺ - K⁺ pump starts functioning d) Stimulus received and influx of Na⁺ ions
- e) Maintenance of resting potential f) Binding of neurotransmitter with post synaptic membrane

5. Match the following

A	B	C
Yellow spot	Macula	CSF
Otolith organ	Neurotransmitter	Keenest vision
Synaptic knob	Subarachnoid space	Acetyl choline
Meninges	Cone cells	Equilibrium

6. Observe the following figure.



- a) Identify the diagrammatic representation.
- b) Name P & Q.
- c) Mention the function of P & Q.

7. Prepare a flow chart showing the steps of hearing processes.

Visit: www.bankofbiology.com

22. CHEMICAL CO-ORDINATION AND INTEGRATION


Endocrine system includes **endocrine (ductless) glands** and their secretions (**hormones**).

Hormones are **non-nutrient** chemicals that act as **intercellular messengers** and are produced in trace amounts.

HUMAN ENDOCRINE GLANDS

They include


1. Hypothalamus
2. Pituitary
3. Pineal
4. Thyroid
5. Parathyroid
6. Thymus
7. Adrenals
8. Pancreas (Islets of Langerhans)
9. Gonads (Testis & Ovary)



1. HYPOTHALAMUS

Neurosecretory cells (nuclei) of hypothalamus secrete the following types of hormones:

- **Releasing hormones:** Stimulate secretion of **pituitary hormones**. E.g. **gonadotropin releasing hormone (GnRH)** stimulates pituitary to release **gonadotropins (FSH & LH)**.
- **Inhibiting hormones:** Inhibit secretion of **pituitary hormones**. E.g. **Somatostatin** inhibits release of growth hormone from pituitary.
- **Oxytocin & vasopressin:** These are transported axonally and stored in pituitary. (*See pituitary gland*).



2. PITUITARY GLAND

- It is located in a bony cavity called **sella tursica**.
- It is attached to **hypothalamus** by a stalk.
- It is divided into anterior **Adenohypophysis** & posterior **Neurohypophysis**.

a. Adenohypophysis

It has 2 parts: **Pars distalis** and **Pars intermedia**.

Pars distalis (Anterior pituitary): It produces

- ♦ **Somatotropin (Growth hormone, GH):** For body growth. Its **over-secretion** causes **Gigantism** (abnormal growth). **Hyposecretion** causes **Dwarfism** (stunted growth). **Over-secretion** of GH in adults (mainly in middle age) causes **Acromegaly** (severe disfigurement especially of face). It leads to serious complications and premature death. Early diagnosis of the disease is difficult. It may be undetected for many years.
- ♦ **Prolactin (PRL):** Regulates growth of **mammary glands** and **milk production**.

- ♦ **Thyroid stimulating hormone (TSH):** Stimulates **thyroid gland** to secrete **thyroid hormones**.
- ♦ **Adrenocorticotrophic hormone (ACTH):** Stimulates **adrenal cortex** to synthesise & secrete **steroid hormones (glucocorticoids)**.
- ♦ **Follicle stimulating hormone (FSH):** Stimulates gonadal activity. **In males**, FSH & androgens regulate sperm formation (**spermatogenesis**). **In females**, FSH stimulates growth and development of **ovarian follicles**.
- ♦ **Luteinizing hormone (LH):** Stimulates gonadal activity. **In males**, it stimulates synthesis and secretion of androgens from testis. **In females**, it induces ovulation and maintains the corpus luteum.

Pars intermedia: In human, it is almost merged with pars distalis. It produces **Melanocyte stimulating hormone (MSH)**. It acts on **melanocytes** to regulate skin pigmentation.

b. Neurohypophysis

It stores **Oxytocin & Vasopressin** from hypothalamus.

- Oxytocin:** Contracts **smooth muscles**. In females, it stimulates contraction of uterus during child birth, and milk ejection from the mammary gland.
- Vasopressin or Anti-diuretic hormone (ADH):** Stimulates **reabsorption of water & electrolytes** by **DCT** of kidney and thereby reduces **diuresis** (loss of water through urine). Deficiency of ADH results in diminished ability of the kidney to conserve water. It leads to water loss and dehydration. This is called **Diabetes insipidus**.

3. PINEAL GLAND


Smallest endocrine gland.

It is located on dorsal side of forebrain. Secretes **melatonin**.

Functions of melatonin:

- Regulates **diurnal (24-hour) rhythm** of body. E.g. sleep-wake cycle, body temperature etc.
- Influences metabolism, pigmentation & menstrual cycle.
- Influences defense capability.

4. THYROID GLAND



- Largest endocrine gland.
 - It includes 2 lobes on either side of the trachea. The lobes are interconnected with **isthmus** (a connective tissue).
 - Thyroid gland is made of **follicles & stromal tissues**.
- Follicular cells** secrete the following hormones:

- **Thyroxin (tetraiodothyronine, T₄) & Triiodothyronine (T₃):** Their functions are
 - Regulation of **basal metabolic rate (BMR)**.
 - Physical, mental and sexual development.
 - Support **RBC formation**.
 - Control **metabolism** of carbohydrates, proteins & fats.
 - Maintain **water and electrolyte balance**.
- **Thyrocalcitonin (TCT):** A protein hormone. It regulates (lowers) **blood calcium** levels (Hypocalcaemic hormone).

Iodine is essential for normal hormone synthesis in thyroid.

Hypothyroidism (Goiter):

- Enlargement of thyroid gland due to deficiency of **iodine**.
- In adult women, it causes irregular menstrual cycle.
- Hypothyroidism during pregnancy affects the baby causing stunted growth (cretinism), mental retardation, low intelligence quotient, abnormal skin, deaf-mutism etc.

Hyperthyroidism:

- Abnormal increase of thyroid hormones resulting in adverse effects on the physiological activities.
- It is caused due to development of the nodules or the cancer of thyroid gland.
- **Exophthalmic goiter (Grave's disease):** It is a form of Hyperthyroidism. Symptoms are enlargement of thyroid gland, protruded eyeballs, increased BMR & weight loss.

5. PARATHYROID GLAND

4 parathyroid glands are present on back side of the thyroid gland, one pair each in the two lobes of thyroid gland. They secrete **Parathyroid hormone (PTH)** – a peptide hormone.

Functions of parathyroid hormone:

- Increases **Ca²⁺ level** in blood (**hypercalcaemic hormone**).
- Stimulates the **bone resorption (demineralization)**.
- Stimulates the **reabsorption of Ca²⁺** by the **renal tubules** and increases **Ca²⁺ absorption** from the **digested food**.
- Along with **TCT**, it helps in **calcium balance** in the body.

6. THYMUS GLAND

It is located between lungs behind sternum on the ventral side of aorta. It secretes **Thymosins** (peptide hormones).

Functions of thymosins:

- Differentiation of **T-lymphocytes**. It provides **cell-mediated immunity**.
- Promote **antibody** production for **humoral immunity**.


Thymus is degenerated in old individuals. So, production of thymosins decreases. As a result, immune responses of old persons become weak.

7. ADRENAL GLAND

It has 2 parts: **Adrenal cortex & Adrenal medulla**.

a. Adrenal cortex

It has 3 layers: inner **zona reticularis**, middle **zona fasciculata** & outer **zona glomerulosa**.



It produces the following **corticoid hormones**:

○ **Glucocorticoids** (mainly **cortisol**):

- Involved in **carbohydrate metabolism**.
- Stimulate **gluconeogenesis, lipolysis and proteolysis**.
- Inhibit cellular uptake and utilization of **amino acids**.
- Maintain **cardiovascular** system and **kidney** functions.
- Cortisol stimulates **RBC production**.
- Produces **anti-inflammatory reactions** and suppress immune response.

○ **Mineralocorticoids** (mainly **aldosterone**):

- Regulate the **water** (body fluid volume), **electrolytic balance, osmotic pressure** and **blood pressure**.
- Aldosterone stimulates **reabsorption of Na⁺ & water** from renal tubules and excretion of **K⁺ and PO₄³⁻ ions**.

○ **Androgenic corticoids:** For growth of **axial hair, pubic hair** and **facial hair** during puberty.

Deficiency of corticoid hormones affects carbohydrate metabolism. It causes acute weakness and fatigue. This condition is called **Addison's disease**.

b. Adrenal medulla

- Produces **catecholamine** hormones such as **Adrenaline (epinephrine) & Noradrenaline (norepinephrine)**.
- They are rapidly secreted in response to stress emergency situations so called **emergency hormones (hormones of Fight or Flight)**.
- These increase alertness, pupillary dilation, piloerection (rising of hairs), sweating, heartbeat, heart contraction and respiratory rate. Stimulate glycogenolysis to increase glucose in blood. Also stimulate lipolysis and proteolysis.

8. PANCREAS (ISLETS OF LANGERHANS)

- A **composite (heterocrine)** gland i.e. **exocrine + endocrine**.
- **Islets of Langerhans** are the endocrine part. There are about 1-2 million Islets (1-2% of pancreatic tissue).
- **α cells** and **β cells** in the islets secrete peptide hormones such as **Glucagon** and **Insulin** respectively. They maintain **Glucose homeostasis** in blood.

• **Glucagon:** Hyperglycemic factor. It

- Acts on **hepatocytes** and stimulates **glycogenolysis** resulting in an increased blood sugar (**hyperglycemia**).
- Stimulates **gluconeogenesis**.
- Reduces the cellular glucose uptake and utilization.

• **Insulin:** Hypoglycemic factor. It

- Acts on **hepatocytes** and **adipocytes** to enhance cellular glucose uptake and utilization. So, glucose from blood rapidly moves to hepatocytes and adipocytes. Thus, blood glucose level decreases (**hypoglycemia**).
- Stimulates **glycogenesis** (glucose converts to glycogen).

Prolonged hyperglycemia leads to **Diabetes mellitus** (loss of glucose through urine and formation of harmful compounds like ketone bodies). Treatment is **insulin therapy**.

9. TESTIS (MALE GONAD)

- It is the male primary sex organ and an endocrine gland.
- A pair of testis is present in the **scrotal sac**.
- It is formed of **seminiferous tubules** and **interstitial (stromal) tissues**.

- **Leydig (interstitial) cells** in the inter-tubular spaces produce hormones called **androgens** (mainly **testosterone**).

Functions of androgens:

- Regulate development, maturation and functions of the **accessory sex organs**.
- **Spermatogenesis** (sperm production).
- Stimulate sexual behavior (**libido**), growth of muscles, hairs, aggressiveness, low pitch voice etc.
- Help in anabolism of protein and carbohydrate.

10. OVARY (FEMALE GONAD)

- It is the female primary sex organ.
- A pair of ovaries is located in the abdomen.
- It produces one ovum during each menstrual cycle.

- Ovary is formed of **ovarian follicles** and **stromal tissues**.
- **Ovarian follicles** produce **Estrogen** (a steroid hormone).
- After ovulation, ruptured follicle forms a structure called **Corpus luteum**. It secretes **progesterone** (a steroid hormone).

Functions of Estrogen:

- Growth and activities of female **secondary sex organs**.
- Development of **ovarian follicles & mammary glands**.
- Female **secondary sex characters** (e.g. high pitch voice) and **sexual behavior**.

Functions of Progesterone:

- It supports **pregnancy**.
- It acts on **mammary glands** to stimulate formation of **alveoli** (sacs to store milk) and **milk secretion**.

HORMONES OF HEART, KIDNEY & GASTROINTESTINAL TRACT

1. **Atrial wall of heart:** Produce a peptide hormone called **Atrial natriuretic factor (ANF)**. When BP increases, ANF causes dilation of blood vessels to reduce the BP.
2. **JGA of kidney:** Produces **Erythropoietin** (peptide hormone). Stimulates **erythropoiesis** (formation of RBC).
3. **Gastro-intestinal tract:** Produce peptide hormones. E.g.
 - **Gastrin:** Stimulates **gastric glands** to secrete **HCl** and **pepsinogen**.
 - **Secretin:** Stimulates **exocrine pancreas** to secrete **water** and **bicarbonate ions**.
 - **Cholecystokinin (CCK):** Stimulates secretion of **bile** from gall bladder and pancreatic enzymes from pancreas.

- **Gastric inhibitory peptide (GIP):** Inhibits **gastric secretion**.

Several other **non-endocrine tissues** secrete hormones called **growth factors**. These help for the normal growth of tissues and their repairing or regeneration.

Based on the chemical nature, hormones are various types:


- Peptide, polypeptide, protein hormones:** Insulin, glucagon, pituitary hormones, hypothalamic hormones etc.
- Steroids:** Cortisol, testosterone, estradiol & progesterone.
- Iodothyronines** (thyroid hormones).
- Amino-acid derivatives:** Adrenaline, nor-adrenaline etc.

MECHANISM OF HORMONE ACTION


- Hormones produce their effects by binding to the specific proteins (**hormone receptors**) located in **target tissues**.
- A hormone binds to its specific receptor to form **hormone-receptor complex**.
- It leads to biochemical changes in target tissue and thereby regulates metabolism and physiological functions.

Hormone receptors are 2 types:

- **Membrane-bound receptors:** Some hormones (e.g. protein hormone, FSH) interact with membrane-bound receptors (do not enter the target cell). It generates **second messengers** (e.g. cyclic AMP, IP₃, Ca²⁺). It in turn regulates **cellular metabolism** and causes **physiological effects**.



- **Intracellular receptors** (mostly **nuclear receptors**): Some hormones (e.g. steroid hormones, iodothyronines) interact with intracellular receptors. They mostly regulate **gene expression** or **chromosome function** by the interaction of hormone-receptor complex with the genome. Cumulative biochemical actions result in physiological and developmental effects.



Visit: www.bankofbiology.com

MODEL QUESTIONS

- Note the relationship between first two words and suggest a suitable word for fourth place.
 - Alpha cell: Glucagon Beta cell:
 - Glucocorticoids: Cortisol Mineralocorticoids:
 - Follicular cells: Thyroid Neurosecretory cells:
 - Ovarian follicles: Estrogen Corpus luteum:

- Odd one out. Justify your answer.

- TSH, FSH, MSH, LH
- Cortisol, adrenaline, aldosterone, androgenic corticoids

- Match the following

A	B	C
Thyroid	Insulin	Addison's disease
Pituitary	Cortisol	Goiter
Pancreas	Thyroxine	Gigantism
Adrenal gland	Growth hormone	Diabetes mellitus

- In a 5-year old boy, thymus gland is found to be non-functional. How will it affect his immune system?
- On a hot day, would you expect ADH level in blood to be high or low? Explain.
- Analyze the facts given in three columns, find their relationship and fill the blanks.

Thymus	a) -----	Differentiation of T-lymphocytes
b) -----	Adrenaline	Emergency hormone
Kidney	Erythropoietin	c) -----
Pancreas	d) -----	Decrease blood glucose level

- Make pairs using following terms:

Hypoglycemic factor, Glucagon, TCT, Hypercalcemic factor, Hyperglycemic factor, PTH, Hypocalcemic factor, Insulin

- Anitha saw a snake on her way to school. She was frightened and her heart rate and breathing rate increased.
 - Name the hormones which are dominant at that time in her blood.
 - Which endocrine gland produces the hormone? c) To which organ this endocrine gland is attached?
- Prepare flowcharts showing the mechanism of action of a protein hormone and a steroid hormone on target tissues.